



## A Note on Dieudonne Complete Spaces

Asylbek A. Chekeev<sup>a,b</sup>, Tumar J. Kasymova<sup>a</sup>

<sup>a</sup>Mathematics and Informatics Faculty, Kyrgyz National University, 720033 Bishkek;

<sup>b</sup>Faculty of Science, Kyrgyz-Turkish Manas University, 720038 Bishkek, Kyrgyz Republic

**Abstract.** In this paper, it is established a characterization of  $\tau$ -normal coverings by means of approximation of the Čech complete paracompacta, which are the perfect preimages of complete metric spaces of weight  $\leq \tau$ . In particular, this characterization generalizes to an arbitrary cardinal the result of A. Garsia-Maynez [15].

### 1. Introduction and Preliminaries

All spaces are assumed to be Tychonoff.  $C(X)$  is the set of all real-valued continuous functions on  $X$ . The set  $Z(f) = \{x \in X : f(x) = 0\}$  is called *zero-set* of a function  $f \in C(X)$ . A family  $\mathcal{Z}(X) = \{Z(f) : f \in C(X)\}$  is the set of all zero-sets on  $X$ . A family  $C\mathcal{Z}(X) = \{X \setminus Z(f) : f \in C(X)\}$  is the set of all *cozero-sets* on  $X$ . For any function  $f \in C(X)$  we will assume  $\text{coz} f = X \setminus Z(f)$ . A family  $\mathcal{Z}(X)$  ( $C\mathcal{Z}(X)$ ) forms a base of closed (open) sets of a space  $X$  [12]. A family  $\mathcal{Z}(X)$  is a *separating nest-generated intersection ring* (s.n.-g.i.r.) [21] or a *strong delta normal base* [1], hence it is a *normal base* [10] and the Wallman compactification  $\omega(X, \mathcal{Z}(X))$  coincides with the Stone–Čech compactification  $\beta X$  [1, 12]. The Hewitt–Nachbin–Shirota completion  $\nu X$  [13, 17, 20] is the Wallman realcompactification  $\nu(X, \mathcal{Z}(X))$  [21]. A space  $\beta X$  consists of all  $z$ -ultrafilters ( $\equiv$  maximal centered systems on  $\mathcal{Z}(X)$ ) with the Wallman normal base  $\{\bar{Z} : Z \in \mathcal{Z}(X)\}$ , where  $\bar{Z} = \{p \in \beta X : Z \in p\}$  [1, 12, 21]. The realcompactification  $\nu X$  is a subspace of  $\beta X$  and it consists of all countably centered (CC)  $z$ -ultrafilters ( $\equiv$  maximal countably centered systems on  $\mathcal{Z}(X)$ ) with a base  $\{\bar{Z} \cap \nu X : Z \in \mathcal{Z}(X)\}$ , where  $\bar{Z} \cap \nu X = \{p \in \nu X : Z \in p\}$  [1, 12, 21]. Hence  $\bar{Z} = [Z]_{\beta X}$  and  $\bar{Z} \cap \nu X = [Z]_{\nu X}$  for all  $Z \in \mathcal{Z}(X)$ .

It is known from [7] that points of  $\beta X$  corresponding to the Dieudonne completion  $\delta X$  (by Curzer-Hager) was described as *co-locally finitely additive (co-LFA)  $z$ -ultrafilters*.

Below the important properties of co-LFA  $z$ -ultrafilters are established by Propositions 2.1, 2.3 and Theorem 2.6. Further, it is established a characterization of  $\tau$ -normal coverings (Theorem 2.10), which implies some results of G. Vidossich [22] and A.Di Concilio [8]. Theorems 2.12 and 2.14 generalize to an arbitrary cardinal the result of A.Garsia-Maynez [15]. By using Theorem 2.10, we prove Theorems 2.18, 2.19, 2.20, where the known characterizations of Hewitt-Nachbin-Shirota completions and realcompact spaces are clarified.

Denote by  $\mathbb{N}$  the set of all natural numbers, and by  $\mathbb{R}$  the real line with the ordinary topology. The union and the intersection of a family  $\alpha = \{U_s\}_{s \in S}$  of sets are denoted by  $\cup_{s \in S} U_s$  and  $\cap_{s \in S} U_s$  respectively; in the case of a sequence  $\{U_n\}_{n \in \mathbb{N}}$  of sets we use the symbols  $\cup_{n \in \mathbb{N}} U_n$  and  $\cap_{n \in \mathbb{N}} U_n$ , and in the case of a non-indexed

2010 *Mathematics Subject Classification*. Primary 54D35; Secondary 54D70, 54D80, 54E15

*Keywords*. zero- (cozero-)sets, locally finite cozero covering,  $z$ -ultrafilter, normal base, uniformity, completion

Received: 13 June 2017; Revised: 22 December 2018; Accepted: 28 December 2018

Communicated by Ljubiiša D.R. Kočinac

*Email addresses*: asyl1.ch.top@mail.ru (Asylbek A. Chekeev), tumar2000@mail.ru (Tumar J. Kasymova)

family  $\alpha = \{U\}_{U \in \alpha}$  of sets we write  $\cup \alpha$  and  $\cap \alpha$ . If  $\cup \alpha = X$ , then the family  $\alpha$  is a *covering* of  $X$ . A covering  $\beta$  is a *refinement* of a covering  $\alpha$  if for every  $B \in \beta$  there exists  $A \in \alpha$  such that  $B \subset A$ . For a covering  $\alpha$  of  $X$  the *star* of a set  $D \subset X$  with respect to  $\alpha$  is the set  $\text{St}(D, \alpha) = \{A \in \alpha : A \cap D \neq \emptyset\}$  and  $\alpha(D) = \cup \text{St}(D, \alpha)$ . A covering  $\beta$  is a *strongly star refinement* of a covering  $\alpha$  if covering  $\{\beta(B) : B \in \beta\}$  is a refinement of  $\alpha$ . If  $\alpha = \{U_s\}_{s \in S}$  and  $\gamma = \{V_t\}_{t \in T}$  are two coverings of  $X$ , then  $\alpha$  *screens*  $\gamma$  in case  $S = T$  and  $U_s \subset V_s$  for all  $s \in S$ .

Let  $Y$  be a subspace of a space  $X$ , then  $f|_Y$  is a restriction of a mapping  $f : X \rightarrow Z$  on  $Y$ , and the set  $[Y]_X$  is the closure of  $Y$  in  $X$ . Let  $[Y]_X = X$  and  $U$  be open in  $Y$ . Then  $Ex_X U = X \setminus [Y \setminus U]_X$  is the largest open subset of  $X$  whose intersection with  $Y$  is equal  $U$ . If  $U, V$  are open in  $Y$ , then  $Ex_X(U \cap V) = Ex_X U \cap Ex_X V$ ,  $U \subset V$  if and only if  $Ex_X U \subset Ex_X V$  [9]. If  $\alpha = \{U_s\}_{s \in S}$  is a covering of  $Y$ , then  $Ex_X \alpha = \{Ex_X U_s\}_{s \in S}$  and  $\cup Ex_X \alpha = \cup_{s \in S} Ex_X U_s$ . For a covering  $\alpha$  of a space  $X$  *inner intersection* is the set  $\alpha \wedge Y = \{A \cap Y : A \in \alpha\}$ .

A filter  $\mathcal{F}$  is said to be *countably centered* (CC) if  $\cap_{n \in \mathbb{N}} F_n \neq \emptyset$  for any sequence  $\{F_n\}_{n \in \mathbb{N}}$  of  $\mathcal{F}$ . A filter  $\mathcal{F}$  is a *Cauchy filter* in a uniform space  $uX$  if for any uniform covering  $\alpha \in u$  there exist  $U \in \alpha$  and  $F \in \mathcal{F}$  such that  $F \subset U$ .

If  $\{Z_s\}_{s \in S}$  is a zero-sets family of  $\mathcal{Z}(X)$  such that  $\{X \setminus Z_s\}_{s \in S}$  is locally finite, then  $\cap_{s \in S} Z_s$  is a zero-set. It follows from important Pasyukov Lemma [18].

**Lemma 1.1.** ([18]) *Let  $f_s : X \rightarrow \mathbb{R}_s$  be a system of continuous functions from a space  $X$  into real lines  $\mathbb{R}_s = \mathbb{R}$  ( $s \in S$ ) with marked zero  $0_s = 0$  such that the system  $\alpha = \{\text{coz} f_s = f_s^{-1}(\mathbb{R}_s \setminus \{0_s\})\}_{s \in S}$  is locally finite in  $X$ . Then the diagonal mapping  $f = \Delta_{s \in S} f_s : X \rightarrow H^\tau$ , (where  $H^\tau = \mathcal{M} \prod_{s \in S} (\mathbb{R}_s, 0_s)$  is the Hilbert space of weight  $\tau = |S|$  obtained as the metric product of  $\mathbb{R}_s$  with marked points  $0_s = 0$  [18]) is defined and continuous.*

Everywhere we will follow the denotation  $\mu X$  of [16] for the Dieudonne completion of a space  $X$ .

Standard references for topological spaces are in the books [9], and for uniform spaces are in the books [3, 14]. Information on the normal bases is in [1, 10, 21].

## 2. Main Results

Remind that  $z$ -ultrafilter  $p$  is *co-locally finitely additive* whenever the family  $\text{co}(p) = \{X \setminus Z : Z \in p\}$  is *locally finitely additive*, i.e.  $\cup \mathcal{F} \in \text{co}(p)$ , whenever  $\mathcal{F} \subset \text{co}(p)$  and  $\mathcal{F}$  is locally finite [7].

**Proposition 2.1.** *For a  $z$ -ultrafilter  $p$  the following are equivalent:*

- (1) *The family  $\text{co}(p)$  is locally finitely additive;*
- (2)  *$\cap_{s \in S} Z_s \neq \emptyset$  for any locally finite subfamily  $\{X \setminus Z_s\}_{s \in S}$  of  $\text{co}(p)$ .*

*Proof.* (1)  $\Rightarrow$  (2). Let  $\{X \setminus Z_s\}_{s \in S}$  be a locally finite subfamily of  $\text{co}(p)$ . Then  $\cup_{s \in S} X \setminus Z_s = X \setminus \cap_{s \in S} Z_s \in \text{co}(p)$ . Hence,  $\cap_{s \in S} Z_s \neq \emptyset$  (by Lemma 1.1,  $\cap_{s \in S} Z_s \in \mathcal{Z}(X)$ ).

(2)  $\Rightarrow$  (1). Let  $\cap_{s \in S} Z_s \neq \emptyset$ , where  $\{X \setminus Z_s\}_{s \in S}$  is a locally finite subfamily of  $\text{co}(p)$ . By Lemma 1.1,  $Z = \cap_{s \in S} Z_s$  is a zero-set. Suppose, that  $Z \notin p$ . Then there exists  $Z' \in p$  such that  $Z \cap Z' = \emptyset$ . But the family  $\{X \setminus Z_s\}_{s \in S} \cup \{X \setminus Z\}$  is locally finite, hence  $\cup_{s \in S} X \setminus Z_s \cup \{X \setminus Z\} = X \setminus (\cap_{s \in S} Z_s \cap Z') \in \text{co}(p)$ , i.e.  $Z \cap Z' \neq \emptyset$ . Contradiction.  $\square$

Further, a co-locally finitely additive  $z$ -ultrafilter we will denoted by *co-LFA  $z$ -ultrafilter*.

A family  $\{Z_s\}_{s \in S}$  of subsets of space  $X$  is called *co-locally finite* (*co-LF*) if the family  $\{X \setminus Z_s\}_{s \in S}$  is locally finite in  $X$ .

**Corollary 2.2.** *Every co-LFA  $z$ -ultrafilter  $p$  is closed with respect to the intersections of co-LF subfamilies, i.e.  $\cap_{s \in S} Z_s \in p$  for every co-LF subfamily  $\{Z_s\}_{s \in S}$  of  $p$ .*

*Proof.* Let  $\{Z_s\}_{s \in S}$  be a co-LF subfamily of the  $z$ -ultrafilter  $p$ , and  $Z \in p$  be an arbitrary element. Then  $\cap_{s \in S} Z_s \neq \emptyset$  and the family  $\{Z_s\}_{s \in S} \cup \{Z\}$  is co-LF. Hence  $\cap_{s \in S} Z_s \cap Z \neq \emptyset$ . Let  $q$  be a  $z$ -ultrafilter such that  $p \cup \{\cap_{s \in S} Z_s\} \subset q$ . Then  $p = q$  and  $\cap_{s \in S} Z_s \in p$ .  $\square$

**Proposition 2.3.** *Every co-LFA  $z$ -ultrafilter  $p$  is countably centered, i.e.  $\cap_{n \in \mathbb{N}} Z_n \neq \emptyset$  for any sequence  $\{Z_n\}_{n \in \mathbb{N}}$  of  $p$ .*

*Proof.* Suppose that there is a sequence  $\{Z_n\}_{n \in \mathbb{N}}$  in  $p$  such that  $\bigcap_{n \in \mathbb{N}} Z_n = \emptyset$ . Then the family  $\{X \setminus Z_n\}_{n \in \mathbb{N}}$  is a countable cozero covering of the space  $X$ . There exists a countable locally finite cozero covering  $\{X \setminus Z(f_n)\}_{n \in \mathbb{N}}$  that screens  $\{X \setminus Z_n\}_{n \in \mathbb{N}}$ , i.e.  $X \setminus Z(f_n) \subset X \setminus Z_n$  for any  $n \in \mathbb{N}$  [1, Theorem 11.1]. Then  $Z_n \subset Z(f_n)$  and the family  $\{X \setminus Z(f_n)\}_{n \in \mathbb{N}}$  is a co-LF subfamily of  $p$ . But  $\bigcap_{n \in \mathbb{N}} Z(f_n) = \emptyset$ . Contradiction.  $\square$

**Corollary 2.4.** *Every co-LFA z-ultrafilter  $p$  is closed with respect to the intersections of countable subfamilies, i.e.  $\bigcap_{n \in \mathbb{N}} Z_n \in p$  for every sequence  $\{Z_n\}_{n \in \mathbb{N}}$  in  $p$ .*

*Proof.* It immediately follows from Proposition 2.3, and the proof is similar to the proof of Corollary 2.2.  $\square$

**Corollary 2.5.** *Let  $\mu X$  be a set of all co-LFA z-ultrafilters on  $X$ . Then  $\mu X \subset \nu X \subset \beta X$ .*

*Proof.* It immediately follows from Corollary 2.4, and construction of the Hewitt–Nachbin completion  $\nu X$  and construction of the Stone–Čech compactification  $\beta X$  [12].  $\square$

**Theorem 2.6.** *Let  $p$  be a z-ultrafilter on a space  $X$ . Then the following are equivalent:*

- (1)  $p$  is a co-LFA z-ultrafilter on  $X$ ;
- (2)  $p$  is a Cauchy filter with respect to the fine uniformity  $u_f$  of  $X$ .

*Proof.* (1)  $\Rightarrow$  (2). Let  $p$  be an arbitrary co-LFA z-ultrafilter. A fine uniformity  $u_f$  has a base  $\mathcal{B}$  consisting of all locally finite cozero coverings [9, 14]. Let  $\alpha = \{X \setminus Z(f_s)\}_{s \in S}$  be locally finite. Then  $\bigcap_{s \in S} Z(f_s) = \emptyset$ . Since the family  $\{Z(f_s)\}_{s \in S}$  is co-LF, it is not contained into z-ultrafilter  $p$ . Hence there exists index  $s_0 \in S$  such that  $Z(f_{s_0}) \notin p$ . Therefore, there exists  $Z_{n_0} \in p$  such that  $Z(f_{s_0}) \cap Z_{n_0} = \emptyset$ . Thus,  $Z_{n_0} \subset X \setminus Z(f_{s_0}) \in \alpha$  and  $p$  is a Cauchy filter with respect to the fine uniformity  $u_f$ .

(2)  $\Rightarrow$  (1). Let a z-ultrafilter  $p$  be a Cauchy filter with respect to the fine uniformity  $u_f$ . Suppose that there is a subfamily  $\{Z_s\}_{s \in S}$  of  $p$  such that it is co-LF and  $\bigcap_{s \in S} Z_s = \emptyset$ . Then the family  $\alpha = \{X \setminus Z_s\}_{s \in S}$  is a locally finite cozero covering of  $X$ . Hence  $\alpha \in \mathcal{B}$ . Therefore, there exist an index  $s_0 \in S$  and  $Z_0 \in p$  such that  $Z_0 \subset X \setminus Z_{s_0} \in \alpha$ . Since  $Z_0 \cap Z_{s_0} = \emptyset$ , we have a contradiction.  $\square$

**Corollary 2.7.**  *$X$  is Dieudonne complete if and only if every co-LFA z-ultrafilter converges.*

*Proof.* It follows immediately from Theorem 2.6.  $\square$

**Corollary 2.8.**  *$\mu X$  with topology induced by the Stone–Čech compactification  $\beta X$  is the Dieudonne completion of  $X$  and points of  $\mu X$  are co-LFA z-ultrafilters.*

From [8, 22] all uniform coverings of cardinality  $\leq \tau$  of the fine uniformity  $u_f$  form the compatible uniformity. The natural problem arises: describe open coverings which are refinements of cozero coverings of cardinality  $\leq \tau$ .

**Definition 2.9.** An open covering  $\alpha$  of a space  $X$  is said to be  $\tau$ -normal if it has a cozero refinement  $\beta$  of cardinality  $|\beta| \leq \tau$ .

**Theorem 2.10.** *Let  $\alpha$  be an open covering of a space  $X$  and  $\tau \geq \aleph_0$  be an arbitrary cardinal. The following are equivalent:*

- (1)  $\alpha$  is  $\tau$ -normal;
- (2) There exists  $Y_\alpha$  such that  $X \subset Y_\alpha \subset \bigcup \text{Ex}_{\beta X} \alpha \subset \beta X$  and  $Y_\alpha$  is a perfect preimage of some complete metric space of weight  $\leq \tau$ .

*Proof.* (1)  $\Rightarrow$  (2). Let a locally finite cozero covering  $\beta$  be a refinement of  $\alpha$  and  $|\beta| \leq \tau$ . Then  $\beta = \{\text{coz} f_s\}_{s \in S}$ , where  $\text{coz} f_s = f_s^{-1}(\mathbb{R} \setminus \{0\})$ ,  $f_s \in C(X)$  and  $|S| \leq \tau$ . By Lemma 1.1, the mapping  $f = \Delta_{s \in S} f_s$  continuously maps  $X$  into the Hilbert space  $H^\tau$ . For any  $s \in S$  we have  $f_s = \pi_s \circ f$ , where  $\pi_s = p_s|_{f(X)}$  is the restriction of the natural projection  $p_s : H^\tau \rightarrow \mathbb{R}_s, s \in S$ . We note that  $\text{coz} f_s = f^{-1}(\text{coz} \pi_s)$ . Let  $M = [f(X)]_{H^\tau}$ . Then  $M$  is a

complete metric subspace of  $H^\tau$ . Let  $F : \beta X \rightarrow \beta M$  be the extension of  $f$  on the Stone-Čech compactifications  $\beta X$  and  $\beta M$ ,  $Y_\alpha = F^{-1}(M)$ . Assume  $\varphi = F|_{Y_\alpha}$ . Then  $\varphi : Y_\alpha \rightarrow M$  is a perfect mapping and  $\varphi^{-1}(M) = Y_\alpha$ . Assume  $Ex_M \text{coz} \pi_s = M \setminus [f(X) \setminus \text{coz} \pi_s]_M$ . Then  $\bigcup_{s \in S} f^{-1}(Ex_M \text{coz} \pi_s) = Y_\alpha$  and  $f^{-1}(Ex_M \text{coz} \pi_s) \cap X = \text{coz} f_s$  for all  $s \in S$ . Hence,  $f^{-1}(Ex_M \text{coz} \pi_s) \subset Ex_{\beta X} \text{coz} f_s$ . Then  $X \subset Y_\alpha \subset \bigcup Ex_{\beta X} \beta \subset \bigcup Ex_{\beta X} \alpha \subset \beta X$ .

(2)  $\Rightarrow$  (1). Suppose that an open covering  $\alpha$  satisfies conditions from (2). Then the inner intersection  $Ex_{\beta X} \alpha \wedge Y_\alpha$  is an open covering of paracompactum  $Y_\alpha$ . Since  $Y_\alpha$  is perfectly mapped onto a complete metric space of weight  $\leq \tau$ , then there exists a locally finite cozero covering  $\gamma$  of cardinality  $|\gamma| \leq \tau$  that is a refinement of  $Ex_{\beta X} \alpha \wedge Y_\alpha$  [2, Chapter VI, 42].  $\square$

Applying the fact that every open covering of a paracompactum has a cozero strongly star refinement, we obtain

**Corollary 2.11.** ([8, 22]) *The collection  $(u_f)_\tau$  of all  $\tau$ -normal coverings of a space  $X$  is a compatible uniformity on  $X$ .*

The next theorem characterizes a completion  $\mu_\tau X$  of a space  $X$  with respect to the uniformity  $(u_f)_\tau$ .

**Theorem 2.12.** *The completion  $\mu_\tau X$  of  $X$  with respect to the uniformity  $(u_f)_\tau$  may be viewed as a subspace of  $\beta X$  containing  $X$ , namely, as the intersection of all paracompact  $G_\delta$ -subspaces of  $\beta X$  containing  $X$ , which are perfect preimages of complete metric spaces of weight  $\leq \tau$ .*

*Proof.* The Stone-Čech precompact uniformity  $u_\beta$  [9, 8.5.8] is containing in  $(u_f)_\tau$ , hence the Samuel compactification of  $X$  with respect to the uniformity  $(u_f)_\tau$  is the Stone-Čech compactification of  $\beta X$ . Then for any  $\tau$ -normal covering  $\alpha$  we have  $X \subset Y_\alpha \subset \bigcup Ex_{\beta X} \alpha \subset \beta X$ , where  $Y_\alpha$  is a paracompact  $G_\delta$ -subspace of  $\beta X$  as a perfect preimage of some complete metric space of weight  $\leq \tau$  (by Theorem 2.10). Hence,  $\mu_\tau X = \bigcap \{Y_\alpha : \alpha \in (u_f)_\tau\}$  as  $\mu_\tau X = \bigcap \{\bigcup Ex_{\beta X} \alpha : \alpha \in (u_f)_\tau\}$  [19].  $\square$

**Remark 2.13.** Theorem 2.12 is an extension of a result [15] to an arbitrary cardinal

The next result clarifies some result of [11].

**Theorem 2.14.** *Let  $\mu X \subset Y \subset \beta X$ . Then the following are equivalent:*

- (1)  $\mu X = Y$ ;
- (2) *For any point  $x \in \beta X \setminus Y$  there exist a cardinal  $\tau \geq \aleph_0$ , a cozero covering  $\alpha$  of  $X$  of cardinality  $|\alpha| \leq \tau$  and  $Y_x$  such that  $X \subset Y \subset Y_x \subset \bigcup Ex_{\beta X} \alpha \subset \beta X \setminus \{x\}$  and  $Y_x$  is the perfect preimage of some complete metric space of weight  $\leq \tau$ .*

*Proof.* (1)  $\Rightarrow$  (2). Let  $\mu X = Y$  and  $x \in \beta X \setminus Y$  be an arbitrary point. Then there exists a unique  $z$ -ultrafilter  $p_x$  on  $X$  such that  $\{x\} = \bigcap \{[Z]_{\beta X} : Z \in p_x\}$  [12] and  $p_x$  is not co-LFA. Hence there exists a co-LF subfamily  $\{Z_s\}_{s \in S}$  of  $p_x$  such that  $|S| \leq \tau$  and  $\bigcap_{s \in S} Z_s = \emptyset$ . By Lemma 1.1, the mapping  $f = \bigtriangleup_{s \in S} f_s$  maps  $X$  into the Hilbert space  $H^\tau$ , where  $Z_s = Z(f_s)$ . As in the proof of implication (1)  $\Rightarrow$  (2) from Theorem 2.10, we have  $f_s = \pi_s \circ f$  and  $\text{coz} f_s = f^{-1}(\text{coz} \pi_s)$ . The closure of  $f(X)$  in  $H^\tau$  is a complete metric space of weight  $\leq \tau$ . Let  $F : \beta X \rightarrow \beta M$  be an extension of  $f$  to the Stone-Čech compactifications  $\beta X$  and  $\beta M$ ,  $Y_\alpha = F^{-1}(M)$  and  $Y = F|_{Y_\alpha}$ . Then  $\varphi : Y_\alpha \rightarrow M$  is a perfect mapping and  $\varphi^{-1}(M) = Y_\alpha$ . Suppose  $Ex_M \text{coz} \pi_s = M \setminus [f(X) \setminus \text{coz} \pi_s]_M$ . Then  $\bigcup_{s \in S} f^{-1}(Ex_M \text{coz} \pi_s) = Y_\alpha$  and  $f^{-1}(Ex_M \text{coz} \pi_s) \cap X = \text{coz} f_s$  for all  $s \in S$ . Hence  $f^{-1}(Ex_M \text{coz} \pi_s) \subset Ex_{\beta X} \text{coz} f_s$ . Then  $\mu X = Y \subset Y_\alpha \subset \bigcup Ex_\alpha$ . It is clear,  $x \notin Ex_\alpha \text{coz} f_s = \beta X \setminus [x \setminus \text{coz} f_s]_{\beta X}$  for all  $s \in S$ . So,  $X \subset Y \subset Y_\alpha \subset \bigcup Ex_{\beta X} \alpha \subset \beta X \setminus \{x\}$ .

(2)  $\Rightarrow$  (1). It is clear,  $\mu X = \bigcap \{Y_x : x \in \beta X \setminus Y\} = Y$ .  $\square$

The following statement clarifies the result of [11].

**Corollary 2.15.** *The following are equivalent:*

- (1)  $X$  is Dieudonne complete;
- (2) *For any point  $x \in \beta X \setminus X$  there exists a cardinal  $\tau \geq \aleph_0$ , a cozero covering  $\alpha$  of  $X$  of cardinality  $|\alpha| \leq \tau$  and  $Y_x$  such that  $X \subset Y_x \subset \bigcup Ex_{\beta X} \alpha \subset \beta X \setminus \{x\}$  and  $Y_x$  is the perfect preimage of some complete metric space of weight  $\leq \tau$ .*

*Proof.* It follows immediately from Theorem 2.14, as  $X = \mu X$ .  $\square$

All countable coverings of the fine uniformity  $u_f$  of a space  $X$  form the uniformity  $u_\omega$  [14]. By Shirota [20] it was proved that the completion of the space  $X$  with respect to the uniformity  $u_\omega$  is the realcompactification  $\nu X$  and  $X$  is realcompact if and only if  $X = \nu X$ . The family  $\mathcal{B}_\omega$  of all countable cozero coverings forms a base of the uniformity  $u_\omega$  [20].

**Theorem 2.16.** *Let  $p$  be a  $z$ -ultrafilter on a space  $X$ . The following are equivalent:*

- (1)  $p$  is a CC  $z$ -ultrafilter on  $X$ ;
- (2)  $p$  is a Cauchy filter with respect to the uniformity  $u_\omega$  of  $X$ .

*Proof.* (1)  $\Rightarrow$  (2). Let  $p$  be an arbitrary CC  $z$ -ultrafilter. The uniformity  $u_\omega$  has a base  $\mathcal{B}_\omega$  of all countable cozero coverings [20]. For any  $\alpha = \{X \setminus Z(f_n)\}_{n \in \mathbb{N}}$  of  $\mathcal{B}_\omega$  we have  $\bigcap_{n \in \mathbb{N}} Z(f_n) = \emptyset$ . Then the family  $\{Z(f_n)\}_{n \in \mathbb{N}}$  is not contained in  $z$ -ultrafilter  $p$ . Hence there exists index  $n_0 \in \mathbb{N}$  such that  $Z(f_{n_0}) \notin p$ . Therefore, there exists  $Z_{n_0} \in p$  such that  $Z_{n_0} \cap Z(f_{n_0}) = \emptyset$ . Thus,  $Z_{n_0} \subset X \setminus Z(f_{n_0}) \in \alpha$  and  $p$  is a Cauchy filter with respect to the uniformity  $u_\omega$ .

(2)  $\Rightarrow$  (1). Let  $p$  be a Cauchy filter with respect to the uniformity  $u_\omega$ . Suppose that there is a sequence  $\{Z_n\}_{n \in \mathbb{N}}$  of  $p$  such that  $\bigcap_{n \in \mathbb{N}} Z_n = \emptyset$ . Then the family  $\alpha = \{X \setminus Z_n\}_{n \in \mathbb{N}}$  is a countable cozero covering of the space  $X$ . Hence  $\alpha \in u_\omega$ . Therefore there exist index  $n_0 \in \mathbb{N}$  and  $Z_0 \in p$  such that  $Z_0 \subset X \setminus Z_{n_0} \in \alpha$ . Since  $Z_0 \cap Z_{n_0} = \emptyset$ , we have a contradiction.  $\square$

The next corollary is well known [9].

**Corollary 2.17.**  *$X$  is realcompact if and only if every CC  $z$ -ultrafilter converges.*

*Proof.* It follows from Theorem 2.16.  $\square$

From Theorem 2.10 in the case  $\tau = \aleph_0$  we obtain the next

**Theorem 2.18.** *Let  $\alpha$  be an open covering of a space  $X$ . The following are equivalent:*

- (1)  $\alpha$  is a uniform covering with respect to  $u_\omega$ ;
- (2) There exists  $Y_\alpha$  such that  $X \subset Y_\alpha \subset \bigcup \text{Ex}_{\beta X} \alpha \subset \beta X$  and  $Y_\alpha$  is the perfect preimage of some complete metric space of countable weight.

*Proof.* It is similar to the proof of Theorem 2.10, assuming  $\tau = \aleph_0$ .  $\square$

The next theorem characterizes the completion  $\nu X$  of a space  $X$  with respect to the uniformity  $u_\omega$ .

**Theorem 2.19.** *The completion  $\nu X$  of  $X$  with respect to the uniformity  $u_\omega$  may be viewed as a subspace of  $\beta X$  containing  $X$ , namely, as the intersection of all Lindelöf  $G_\delta$ -subspaces of  $\beta X$  containing  $X$ , which are perfect preimages of complete metric spaces of countable weight.*

*Proof.* The Stone-Čech precompact uniformity  $u_\beta$  [9, 8.5.8] is contained in  $u_\omega$ , hence the Samuel compactification of  $X$  with respect to the uniformity  $u_\omega$  is the Stone-Čech compactification of  $\beta X$ . Then for any countable normal covering  $\alpha$  we have  $X \subset Y_\alpha \subset \bigcup \text{Ex}_{\beta X} \alpha \subset \beta X$ , where  $Y_\alpha$  is a Lindelöf  $G_\delta$ -subspace of  $\beta X$  as the perfect preimage of some complete metric space of countable weight (Theorem 2.10). Hence,  $\nu X = \bigcap \{Y_\alpha : \alpha \in u_\omega\}$  as  $\nu X = \bigcap \{\bigcup \text{Ex}_{\beta X} \alpha : \alpha \in u_\omega\}$  [19].  $\square$

**Theorem 2.20.** *Let  $\nu X \subset Y \subset \beta X$ . Then the following are equivalent:*

- (1)  $\nu X = Y$ ;
- (2) For any point  $x \in \beta X \setminus Y$  there exist countable cozero covering  $\alpha$  of  $X$  and  $Y_x$  such that  $X \subset Y \subset Y_x \subset \bigcup \text{Ex}_{\beta X} \alpha \subset \beta X \setminus \{x\}$  and  $Y_x$  is the perfect preimage of some complete metric space of countable weight.

*Proof.* It is similar to the proof of Theorem 2.14, assuming  $\tau = \aleph_0$ .  $\square$

**Corollary 2.21.** *The following are equivalent:*

- (1)  $X$  is realcompact;
- (2) For any  $x \in \beta X \setminus X$  there exist a countable cozero covering  $\alpha$  and a Lindelöf  $G_\delta$ -subspace  $Y_x$  of  $\beta X$  such that  $X \subset Y_x \subset \bigcup Ex_{\beta X} \alpha \subset \beta X \setminus X$ .

For every space  $X$  with a uniformity  $u$  the set  $\mathcal{Z}_u$  of all zero-sets of uniformly continuous real-valued functions of a uniform space  $uX$  form s.n.-g.i.r. [21] and the characterizations of the Wallman  $\beta$ -like compactification  $\omega(X, \mathcal{Z}_u) = \beta_u X$  and realcompactification  $v(X, \mathcal{Z}_u) = v_u X$  are given in [4–6]. A characterization of the Wallman-Dieudonne completion  $\mu_u X$  is given in [6]. Thus, the next problem arises:

**Problem 2.22.** *Let  $\mathcal{Z} \subset \mathcal{Z}(X)$  be an arbitrary s.n.-g.i.r. on a space  $X$  and  $\mu(X, \mathcal{Z})$  be the set of all co-LFA-ultrafilters on  $\mathcal{Z}$ . It is clear that  $\mu(X, \mathcal{Z}) \subset \omega(X, \mathcal{Z})$ . Is the following true:*

- (1)  $\mu(X, \mathcal{Z}) \subset v(X, \mathcal{Z})$ , where  $v(X, \mathcal{Z})$  is the Wallman realcompactification ?
- (2)  $\mu(X, \mathcal{Z})$  is Dieudonne complete in the induced topology from the compactification  $\omega(X, \mathcal{Z})$ ?

## Acknowledgement

The authors wish to thank the referees for corrections of the original draft of the paper and for many helpful remarks.

## References

- [1] A.R. Alo, H.L. Shapiro, Normal Topological Spaces, Cambridge University Press, 1974, 306 p.
- [2] A.V. Arhangel'skii, V.I. Ponomarev, Fundamentals of General Topology in Problems and Exercises, Nauka, Moscow, 1974. 424 p. (in Russian).
- [3] A.A. Borubaev, A.A. Chekeev, Uniform Spaces, Bishkek, 2003, 245 p. (in Russian).
- [4] A.A. Chekeev, Uniformities for Wallman compactifications and realcompactifications, Topol. Appl. 201 (2016) 145–156.
- [5] A.A. Chekeev, B.Z. Rakhmankulov, A.I. Chanbaeva, On  $C_i$ - and  $C_u$ -embedded uniform spaces, TWMS J. Pure Appl. Math. 9 (2018) 173–189
- [6] A.A. Chekeev, T.J. Kasymova, Ultrafilter-completeness on zero-sets of uniformly continuous functions, Topol. Appl. 252 (2019) 27–41.
- [7] H. Curzer, A.W. Hager, On the topological completion, Proc. Amer. Math. Soc., 56(4), (1976), 365–370.
- [8] A. Di Concilio, Uniform properties and hyperspace topologies for  $\mathbf{N}$ -uniformities, Topol. Appl. 44 (1992) 115–123.
- [9] R. Engelking, General Topology, Heldermann, Berlin, 1989, 626 p.
- [10] O. Frink, Compactifications and seminormal spaces, Amer. J. Math. 86 (1964) 602–607.
- [11] Z. Frolik, On approximation and uniform approximation of spaces, Proc. Japan. Acad. 37 (1961) 530–532.
- [12] L. Gillman, M. Jerison, Rings of Continuous Functions, Princeton, 1960, 303 p.
- [13] E. Hewitt, Rings of real-valued continuous functions, I, Trans. Amer. Math. Soc. 64 (1948) 45–99.
- [14] J.R. Isbell, Uniform Spaces: Mathematical Survey, Providence, 1964, 175 p.
- [15] A. Garsia-Mainez, Uniform Spaces, II, Encyclopedia of General Topology, Elsevier, North-Holland (2004) 264–265.
- [16] K. Morita, Topological completions and  $M$ -spaces, Sci. Rept. Kyoiku Daigaku (A) 10 (1970) 271–288.
- [17] L. Nachbin, On the continuity of positive linear transformations, Proc. Internat. Congress Math., Cambridge, Mass. 1950, vol. I, Providence 1952, 464–465.
- [18] B.A. Pasyukov, Factorization theorems in dimension theory, Uspehi Mat. Nauk 36:3 (1981) 147–176 (in Russian).
- [19] Yu. Smirnov, On completeness of proximity spaces, Trudy Mosk. Math. Soc. 3 (1954) 271–306. (in Russian)
- [20] T. Shirota, A class of topological spaces, Osaka Math. J. 4 (1952) 23–40.
- [21] A.K. Steiner, E.F. Steiner, Nest generated intersection rings in Tychonoff spaces, Trans. Amer. Math. Soc. 148 (1970) 589–601.
- [22] G. Vidossich, A note on cardinal reflections in the category of uniform spaces, Proc. Amer. Math. Soc. 23 (1969) 55–58.