

SOME RELATIONS IN THE GENERALIZED KÄHLERIAN SPACES OF THE SECOND KIND

Mića S. Stanković *, Ljubica S. Velimirović, Milan Lj. Zlatanović

Abstract

Starting from the definition of generalized Riemannian space (GR_N) [1], in which a non-symmetric basic tensor g_{ij} is introduced, in the present paper a generalized Kählerian space GK_N^2 of the second kind is defined, as a GR_N with almost complex structure F_i^h , that is covariantly constant with respect to the second kind of covariant derivative (equation (2.3)).

Several theorems are proved. These theorems are generalizations of the corresponding theorems relating to K_N . The relations between F_i^h and four curvature tensors from GR_N are obtained.

1 Introduction

A generalized Riemannian space GR_N in the sense of Eisenhart's definition [1] is a differentiable N -dimensional manifold, equipped with a non-symmetric basic tensor g_{ij} . Connection coefficients of this space are generalized Cristoffel's symbols of the second kind. Generally, $\Gamma_{jk}^i \neq \Gamma_{kj}^i$.

In a generalized Riemannian space one can define four kinds of covariant derivatives [3], [4]. For example, for a tensor a_j^i in GR_N we have

$$\begin{aligned} a_{j|_1}^i &= a_{j,m}^i + \Gamma_{pm}^i a_j^p - \Gamma_{jm}^p a_p^i, & a_{j|_2}^i &= a_{j,m}^i + \Gamma_{mp}^i a_j^p - \Gamma_{mj}^p a_p^i, \\ a_{j|_3}^i &= a_{j,m}^i + \Gamma_{pm}^i a_j^p - \Gamma_{mj}^p a_p^i, & a_{j|_4}^i &= a_{j,m}^i + \Gamma_{mp}^i a_j^p - \Gamma_{jm}^p a_p^i. \end{aligned}$$

*Corresponding author

2000 *Mathematics Subject Classifications*. 53A45, 53B05.

Key words and Phrases. Generalized Riemannian space, Kählerian space, generalized Kählerian space, generalized Kählerian space of the second kind.

The authors were supported by Project 144032D MNTR Serbia

In the case of the space GR_N we have five independent curvature tensors [5]:

$$\begin{aligned} R_1^i{}_{jmn} &= \Gamma_{jm,n}^i - \Gamma_{jn,m}^i + \Gamma_{jm}^p \Gamma_{pn}^i - \Gamma_{jn}^p \Gamma_{pm}^i, \\ R_2^i{}_{jmn} &= \Gamma_{mj,n}^i - \Gamma_{nj,m}^i + \Gamma_{mj}^p \Gamma_{np}^i - \Gamma_{nj}^p \Gamma_{mp}^i, \\ R_3^i{}_{jmn} &= \Gamma_{jm,n}^i - \Gamma_{nj,m}^i + \Gamma_{jm}^p \Gamma_{np}^i - \Gamma_{nj}^p \Gamma_{pm}^i + \Gamma_{nm}^p (\Gamma_{pj}^i - \Gamma_{jp}^i), \\ R_4^i{}_{jmn} &= \Gamma_{jm,n}^i - \Gamma_{nj,m}^i + \Gamma_{jm}^p \Gamma_{np}^i - \Gamma_{nj}^p \Gamma_{pm}^i + \Gamma_{mn}^p (\Gamma_{pj}^i - \Gamma_{jp}^i), \\ R_5^i{}_{jmn} &= \frac{1}{2} (\Gamma_{jm,n}^i + \Gamma_{mj,n}^i - \Gamma_{jn,m}^i - \Gamma_{nj,m}^i + \Gamma_{jm}^p \Gamma_{pn}^i + \Gamma_{mj}^p \Gamma_{np}^i \\ &\quad - \Gamma_{jn}^p \Gamma_{mp}^i - \Gamma_{nj}^p \Gamma_{pm}^i). \end{aligned}$$

Kählerian spaces and their mappings were investigated by many authors, for example K. Yano [14], [15], M. Prvanović [12], T. Otsuki [11], N. S. Sinyukov [13], J. Mikeš [2] and many others.

In [6], [7], [8] we defined a generalized Kählerian space GK_N as a generalized N -dimensional Riemannian space with a (non-symmetric) metric tensor

$$g_{ij} = g_{\underline{ij}} + g_{\check{ij}},$$

where $g_{\underline{ij}}$ is symmetric part, and $g_{\check{ij}}$ anty-symmetric one of the metric tensor.

The lowering and the raising of indices one defines by the tensors $g_{\underline{ij}}$ and $g^{\check{ij}}$ respectively, where $g^{\check{ij}}$ is defined by the equation

$$g_{\underline{ij}} g^{\check{jk}} = \delta_i^k, \quad i, k = 1, \dots, m.$$

and δ_i^k is Kronecker symbol. Therefore, since the matrix $(g^{\check{ij}})$ is inverse for $(g_{\underline{ij}})$ it is necessary to be

$$g = \det(g_{\underline{ij}}) \neq 0.$$

There exist an almost complex structure F_j^i such that

$$\begin{aligned} (1.1) \quad & F_p^h(x) F_i^p(x) = -\delta_i^h, \\ (1.2) \quad & g_{\underline{pq}} F_i^p F_j^q = g_{\underline{ij}}, \quad g^{\check{ij}} = g^{\underline{pq}} F_p^i F_q^j, \\ (1.3) \quad & F_{i|j}^h = 0, \quad (\theta = 1, 2), \end{aligned}$$

where $|_{\theta}$ denotes the covariant derivative of the kind θ with respect to the metric tensor g_{ij} .

In [9] we defined a generalized Kählerian space of the first kind GK_N^1 if there exists an almost complex structure $F_j^i(x)$, such that

$$\begin{aligned} (1.4) \quad & F_p^h(x) F_i^p(x) = -\delta_i^h, \\ (1.5) \quad & g_{\underline{pq}} F_i^p F_j^q = g_{\underline{ij}}, \quad g^{\check{ij}} = g^{\underline{pq}} F_p^i F_q^j, \\ (1.6) \quad & F_{i|j}^h = 0, \end{aligned}$$

where $\underset{1}{|}$ denotes the covariant derivative of the first kind with respect to the metric tensor g_{ij} .

2 Generalized Kählerian spaces of the second kind

A generalized N -dimensional Riemannian space with (non-symmetric) metric tensor g_{ij} is a generalized Kählerian space of the second kind GK_{2N} if there exists an almost complex structure $F_j^i(x)$ such that

$$(2.1) \quad F_p^h(x)F_i^p(x) = -\delta_i^h,$$

$$(2.2) \quad g_{pq}F_i^pF_j^q = g_{ij}, \quad g^{ij} = g^{pq}F_p^iF_q^j,$$

$$(2.3) \quad F_{i|j}^h = 0,$$

where $\underset{2}{|}$ denotes the covariant derivative of the first kind with respect to the metric tensor g_{ij} . From (2.2), using (2.1), we get

$$(2.4) \quad F_{ij} = -F_{ji}, \quad F^{ij} = -F^{ji},$$

where we denote

$$(2.5) \quad F_{ji} = F_j^p g_{pi}, \quad F^{ji} = F_p^j g^{pi}.$$

From here we prove the following theorems.

Theorem 2.1. *For the almost complex structure F_j^i of a generalized Kählerian space of the second kind the relations*

$$(2.6) \quad \begin{aligned} F_{i|j}^h &= 2(F_i^p \Gamma_{pj}^h + F_p^h \Gamma_{ji}^p), \\ F_{i|j}^h &= 2F_i^p \Gamma_{pj}^h, \\ F_{i|j}^h &= 2F_p^h \Gamma_{ji}^p \end{aligned}$$

are valid, where Γ_{ij}^h is the torsion tensor.

Proof. We get the relations (2.6) by using the condition (2.3) □

Let us denote $\overline{F}_{ij}^h = F_i^p \Gamma_{jp}^h$ and $\overline{\overline{F}}_{ij}^h = F_p^h \Gamma_{ji}^p$. Then we have

Theorem 2.2. For the curvature tensors R_θ^h , ($\theta = 1, \dots, 4$) of a generalized Kählerian space of the second kind the relations

$$\begin{aligned}
(2.7) \quad & F_i^p R_2^h{}_{pjk} = F_p^h R_2^p{}_{ijk}, \\
& F_i^p R_1^h{}_{pijk} - F_p^h R_1^p{}_{ijk} - 4\Gamma_{jk}^p (F_i^q \Gamma_{qp}^h + F_q^h \Gamma_{pi}^q) \\
& = 2(\bar{F}_{ik|j}^h - \bar{F}_{ij|k}^h + \bar{\bar{F}}_{ik|j}^h - \bar{\bar{F}}_{ij|k}^h), \\
& F_i^p R_3^h{}_{pjk} - F_p^h R_3^p{}_{ijk} = -2(\bar{F}_{ij|k}^h + \bar{\bar{F}}_{ij|k}^h), \\
& F_i^p R_4^h{}_{pjk} + F_p^h R_3^p{}_{ijk} = 2(\bar{\bar{F}}_{ik|j}^h - \bar{F}_{ij|k}^h)
\end{aligned}$$

are valid.

Proof. From (2.6) by using the covariant derivative of the first kind we have

$$(2.8) \quad F_{i|jk}^h = -2\bar{F}_{ij|k}^h - 2\bar{\bar{F}}_{ij|k}^h,$$

and also

$$(2.9) \quad F_{i|kj}^h = -2\bar{F}_{ik|j}^h - 2\bar{\bar{F}}_{ik|j}^h.$$

Now, from (2.8) and (2.9) we obtain

$$(2.10) \quad F_{i|jk}^h - F_{i|kj}^h = 2(\bar{F}_{ik|j}^h - \bar{F}_{ij|k}^h + \bar{\bar{F}}_{ik|j}^h - \bar{\bar{F}}_{ij|k}^h).$$

Using the Ricci identity [5], we get from (2.10)

$$\begin{aligned}
(2.11) \quad & F_i^p R_1^h{}_{pjk} - F_p^h R_1^p{}_{ijk} - 2\Gamma_{jk}^p F_{i|p}^h \\
& = 2(\bar{F}_{ik|j}^h - \bar{F}_{ij|k}^h + \bar{\bar{F}}_{ik|j}^h - \bar{\bar{F}}_{ij|k}^h),
\end{aligned}$$

and from here the second equality (2.7) is valid.

The first equality we get directly from the Ricci identity obtained by virtue of the second kind of covariant derivative and using (2.3).

By the same procedure like in the previous two cases, it is easy to prove the third and the fourth equation. \square

If we denote with $(;)$ covariant derivative wrt the symmetric connection, then the next theorem follows.

Theorem 2.3. For the Ricci tensor R_{ij} , given by g_{ij} the relation

$$(2.12) \quad R_{hk} = F_h^p F_k^q R_{pq} - g^{pq} F_h^s (\mathcal{D}_{s.pqk} + \mathcal{D}_{k.pqs})$$

is valid, where

$$(2.13) \quad \begin{aligned} \mathcal{D}_{ijk}^h &= F_{i;k}^p \Gamma_{jp}^h - F_{i;j}^p \Gamma_{kp}^h + F_i^p (\Gamma_{jp;k}^h - \Gamma_{kp;j}^h) \\ &+ F_{p;k}^h \Gamma_{ij}^p - F_{p;j}^h \Gamma_{ik}^p + F_p^h (\Gamma_{ij;k}^p - \Gamma_{ik;j}^p), \end{aligned}$$

and $R_{hk} = R_{hkp}^p$, $\mathcal{D}_{h.ijk} = g_{ph} \mathcal{D}_{ijk}^p$.

Proof. From (2.3) and (2.6) we get

$$(2.14) \quad F_{i;j}^h = -(F_i^p \Gamma_{jp}^h + F_p^h \Gamma_{ij}^p).$$

The integrability conditions of the equation (2.14) are given by

$$(2.15) \quad F_{i;jk}^h - F_{i;kj}^h = -\mathcal{D}_{ijk}^h.$$

Using the Ricci identity, from (2.15) we obtain

$$(2.16) \quad F_p^h R_{ijk}^p - F_i^p R_{pjk}^h = -\mathcal{D}_{ijk}^h.$$

Here R_{ijk}^h is the curvature tensor with respect to the symmetric affine connection Γ_{ij}^h . Composition with F_r^i in (2.16) gives

$$(2.17) \quad F_p^h F_i^q R_{qjk}^p + R_{ijk}^h = -F_i^p \mathcal{D}_{pjk}^h.$$

Now, from (2.17) by composition with g_{hr} we get

$$(2.18) \quad F_{ph} F_i^q R_{qjk}^p + R_{hijk} = -F_i^p \mathcal{D}_{h.pjk}.$$

From here we obtain

$$(2.19) \quad -F_p^h F_i^q R_{pqjk} + R_{hijk} = -F_i^p \mathcal{D}_{h.pjk}.$$

From (2.19) by composition with F_r^i we have

$$(2.20) \quad F_h^p R_{pijk} - F_i^p R_{phjk} = \mathcal{D}_{h.ijk}.$$

Using composition with g^{ij} in (2.20) we obtain

$$(2.21) \quad F_h^p R_{pk} - F_q^p R_{ph.k} = g^{pq} \mathcal{D}_{h.pqk}.$$

The symmetrization in (2.21) with respect to h, k gives the relation (2.12). \square

Theorem 2.4. *The Ricci tensors $R_{\theta jm}$ ($\theta = 1, \dots, 5$) of the space GK_N satisfy the relations*

$$(2.22 a, b, c) \quad \begin{aligned} R_{\alpha}^{(pq)} F_j^p F_m^q &= R_{\alpha}^{(jm)} - 2\Gamma_{r\check{q}}^p \Gamma_{p\check{s}}^q F_j^r F_m^s \\ &\quad + 2\Gamma_{j\check{q}}^p \Gamma_{p\check{m}}^q + 2g^{pq} F_h^s (\mathcal{D}_{s.pqk} + \mathcal{D}_{k.pqs}), \quad \alpha = 1, 2, 3, \end{aligned}$$

$$(2.22 d) \quad \begin{aligned} R_4^{(pq)} F_j^p F_m^q &= R_4^{(jm)} + 6\Gamma_{r\check{q}}^p \Gamma_{p\check{s}}^q F_j^r F_m^s \\ &\quad - 6\Gamma_{j\check{q}}^p \Gamma_{p\check{m}}^q + 2g^{pq} F_h^s (\mathcal{D}_{s.pqk} + \mathcal{D}_{k.pqs}), \end{aligned}$$

$$(2.22 e) \quad \begin{aligned} R_5^{(pq)} F_j^p F_m^q &= R_5^{(jm)} + 2\Gamma_{r\check{q}}^p \Gamma_{p\check{s}}^q F_j^r F_m^s \\ &\quad - 2\Gamma_{j\check{q}}^p \Gamma_{p\check{m}}^q + 2g^{pq} F_h^s (\mathcal{D}_{s.pqk} + \mathcal{D}_{k.pqs}), \end{aligned}$$

where (jm) denotes the symmetrization without division with respect to the indices j, m .

Proof. (a) We can express the tensor $R_1^i{}_{jmn}$ in the form [5]:

$$R_1^i{}_{jmn} = R^i{}_{jmn} + \Gamma_{j\check{m};n}^i - \Gamma_{j\check{n};m}^i + \Gamma_{j\check{m}}^p \Gamma_{p\check{n}}^i - \Gamma_{j\check{n}}^p \Gamma_{p\check{m}}^i.$$

By contraction with respect to the indices i, n , and by symmetrization with respect to j, m , we get

$$(2.23) \quad R_1^{(jm)} = R_{(jm)} - 2\Gamma_{j\check{q}}^p \Gamma_{p\check{m}}^q.$$

From (2.12) and (2.23) we have (2.22a).

(b) The tensor $R_2^i{}_{jmn}$ can be expressed in the form [5]:

$$R_2^i{}_{jmn} = R^i{}_{jmn} - \Gamma_{j\check{m};n}^i + \Gamma_{j\check{n};m}^i - \Gamma_{j\check{m}}^p \Gamma_{p\check{n}}^i + \Gamma_{j\check{n}}^p \Gamma_{p\check{m}}^i.$$

By contraction with respect to i, n , and then by symmetrization with respect to j, m , we get

$$R_2^{(jm)} = R_{(jm)} - 2\Gamma_{j\check{q}}^p \Gamma_{p\check{m}}^q,$$

from where, using (2.12), we get the relation (2.22b).

(c) For the tensor $R_3^i{}_{jmn}$ we have [5]:

$$R_3^i{}_{jmn} = R^i{}_{jmn} + \Gamma_{j\check{m};n}^i + \Gamma_{j\check{n};m}^i - \Gamma_{j\check{m}}^p \Gamma_{p\check{n}}^i + \Gamma_{j\check{n}}^p \Gamma_{p\check{m}}^i - 2\Gamma_{m\check{n}}^p \Gamma_{p\check{j}}^i.$$

Contracting with respect to i, n , and then symmetrizing in relation to j, m , we get

$$R_3^{(jm)} = R_{(jm)} - 2\Gamma_{jq}^p \Gamma_{pm}^q,$$

from where, using (2.12), we can see that the relation (2.22c) is valid.

(d) The tensor $R_4^i{}_{jmn}$ can be expressed in the form [5]:

$$R_4^i{}_{jmn} = R^i{}_{jmn} + \Gamma_{jm;n}^i + \Gamma_{jn;m}^i - \Gamma_{jm}^p \Gamma_{pn}^i + \Gamma_{jn}^p \Gamma_{pm}^i + 2\Gamma_{mn}^p \Gamma_{pj}^i.$$

Contracting with respect to i, n , and symmetrizing with respect to j, m , we get

$$R_4^{(jm)} = R_{(jm)} + 6\Gamma_{jq}^p \Gamma_{pm}^q.$$

Using (2.12) we get the relation (2.22d).

(e) The tensor $R_5^i{}_{jmn}$ satisfies the relation [5]:

$$R_5^i{}_{jmn} = R^i{}_{jmn} + \Gamma_{jm}^p \Gamma_{pn}^i + \Gamma_{jn}^p \Gamma_{pm}^i.$$

Contracting with respect the indices i, n , and than symmetrizing with respect to j, m , we get

$$R_5^{(jm)} = R_{(jm)} + 2\Gamma_{jq}^p \Gamma_{pm}^q,$$

from where, using (2.12), we get (2.22e). □

References

- [1] L. P. Eisenhart, *Generalized Riemannian spaces I*, Proc. Nat. Acad. Sci. USA, **37** (1951), 311–315.
- [2] J. Mikeš, G. A. Starko, *K-koncircular vector fields and holomorphically projective mappings on Kählerian spaces*, Rend. del Circolo di Palermo, **46** (1997), 123–127.
- [3] S. M. Minčić, *Ricci identities in the space of non-symmetric affine connection*, Matematički Vesnik, **10**(25), (1973), 161–172.
- [4] S. M. Minčić, *New commutation formulas in the non-symmetric affine connection space*, Publ. Inst. Math. (Beograd) (N. S), **22**(36), (1977), 189–199.
- [5] S. M. Minčić, *Independent curvature tensors and pseudotensors of spaces with non-symmetric affine connection*, Coll. Math. Soc. János Bolyai, **31**, (1979), 445–460.
- [6] S. M. Minčić, M. S. Stanković, Lj. S. Velimirović, *Generalized Kählerian spaces*, Filomat, **15**, (2001), 167–174.
- [7] M. S. Stanković, S. M. Minčić, Lj. S. Velimirović, *On Holomorphically Projective Mappings of Generalized Kählerian Spaces*, Matematički vesnik **54**, (2002), 195–202.

- [8] M. S. Stanković, S. M. Minčić, Lj. S. Velimirović, *On equitorsion Holomorphically projective mappings of generalized Kählerian spaces*, Czech. Math. Jour., **54(129)** (2004), 701–715.
- [9] M. S. Stanković, M. Lj. Zlatanović, Lj. S. Velimirović, *Equitorsion holomorphically projective mappings of generalized Kählerian spaces of the first kind*, Czech. Math. Jour., accepted for publication.
- [10] M. S. Stanković, Lj. S. Velimirović, S. M. Minčić, M. Lj. Zlatanović, *Equitorsion conform mappings of generalized Riemannian spaces*, Matematički vesnik, **61**, (2009), 119–129.
- [11] T. Otsuki, Y. Tasiro, *On curves in Kählerian spaces*, Math. J. Okayama Univ., **4** No1, (1954), 57–78
- [12] M. Prvanović, *A note on holomorphically projective transformations of the Kähler space*, Tensor, N. S. Vol. **35**, (1981), 99–104.
- [13] N. S. Sinyukov, *Geodesic Mappings of Riemannian Spaces*, Nauka, Moscow, 1979 (in Russian).
- [14] K. Yano, *Differential Geometry of Complex and Almost Complex Spaces*, Pergamon Press, New York, 1965.
- [15] K. Yano, *On complex conformal connections*, Kodai Math. Sem. Rep., **26**, (1975), 137–151.

Mića S. Stanković,

University of Niš, Faculty of Science and Mathematics,
E-mail: stmica@ptt.rs

Ljubica S. Velimirović,

University of Niš, Faculty of Science and Mathematics,
E-mail: vljubica@pmf.ni.ac.yu

Milan Lj. Zlatanović,

University of Niš, Faculty of Science and Mathematics,
E-mail: zlatmilan@yahoo.com