



Prime Ends in Theory of Mappings with Finite Distortion in the Plane

Denis Kovtonyuk^a, Igor Petkov^a, Vladimir Ryazanov^a

^a*Institute of Applied Mathematics and Mechanics, National Academy of Sciences of Ukraine*

Abstract. In the present paper, it is studied the boundary behavior of the so-called lower Q -homeomorphisms in the plane that are a natural generalization of the quasiconformal mappings. In particular, it was found a series of effective conditions on the function $Q(z)$ for a homeomorphic extension of the given mappings to the boundary by prime ends. The developed theory is applied to mappings with finite distortion by Iwaniec, also to solutions of the Beltrami equations, as well as to finitely bi-Lipschitz mappings that a far-reaching extension of the known classes of isometric and quasiisometric mappings.

1. Introduction

The problem of the boundary behavior is one of the central topics of the theory of quasiconformal mappings and their generalizations. During the last years, it is intensively studied various classes of mappings with finite distortion in a natural way generalizing conformal, quasiconformal and quasiregular mappings, see, e.g., many references in the monographs [16, 18, 20] and [49]. In this case, as it was earlier, the main geometric approach in the modern mapping theory is the method of moduli, see, e.g., the monographs [16, 49, 55, 64, 88, 89] and [91].

One of the main foundations in the present paper is the theory of the boundary behavior by prime ends for the so-called lower Q -homeomorphisms developed here in the first part. The second one is Theorem 3.1 in [24] on solutions of the Beltrami equations that can be reformulated in the following way.

Theorem 1.1. *Let f be a homeomorphism with finite distortion in a domain $D \subseteq \mathbb{C}$. Then f is a lower Q -homeomorphism at each point $z_0 \in \overline{D}$ with $Q = K_f$.*

As usual, $K_f(z)$ denotes the **dilatation** of the mapping f at z , namely,

$$K_f(z) = \frac{|f_z| + |f_{\bar{z}}|}{|f_z| - |f_{\bar{z}}|} \quad (1)$$

if $f_z \neq 0$, 1 if $f_z = 0$ and ∞ otherwise, where $f_{\bar{z}} = (f_x + if_y)/2$, $f_z = (f_x - if_y)/2$, $z = x + iy$, and f_x and f_y are partial derivatives of f in x and y , correspondingly. For the definition of homeomorphisms with finite distortion by Iwaniec, see Section 8.

2010 *Mathematics Subject Classification.* Primary 30C62, 30C65, 30D40, 37E30; Secondary 35A16, 35A23, 35J67, 35J70, 35J75

Keywords. Boundary behavior, prime ends, mappings of finite distortion, lower Q -homeomorphisms, finitely bi-Lipschitz mappings, isometric and quasiisometric mappings, Beltrami equations

Received: 10 February 2015; Accepted: 15 July 2015

Communicated by Miodrag Mateljević

Email addresses: denis_kovtonyuk@bk.ru (Denis Kovtonyuk), igorpetkov@list.ru (Igor Petkov), vl.ryazanov1@gmail.com (Vladimir Ryazanov)

Here we follow Caratheodory [5] in the definition of the prime ends for bounded finitely connected domains in \mathbb{C} and refer readers to Chapter 9 in [7], see also [1] and [54], for the history of the question. Recall relevant notions. Given a domain D in \mathbb{C} , a Jordan arc $\sigma : (0, 1) \rightarrow D$ with its end points in ∂D is called a **cross-cut** of D . Later on, we use the designation σ also for the whole image of $(0, 1)$ under the map σ and the designation $\bar{\sigma}$ for its closure.

A sequence $\sigma_1, \dots, \sigma_m, \dots$ of cross-cuts of D is called a **chain** if:

- (i) $\bar{\sigma}_i \cap \bar{\sigma}_j = \emptyset$ for every $i \neq j, i, j = 1, 2, \dots$;
- (ii) σ_{m-1} and σ_{m+1} are contained in different components of $D \setminus \sigma_m$ for every $m > 1$;
- (iii) $\delta(\sigma_m) \rightarrow 0$ as $m \rightarrow \infty$.

Here $\delta(E)$ denotes the Euclidean diameter of a set E in \mathbb{C} .

Correspondingly to the definition, a chain of cross-cuts $\{\sigma_m\}$ determines a chain of domains $d_m \subset D$ such that $\partial d_m \cap D \subseteq \sigma_m$ and $d_1 \supset d_2 \supset \dots \supset d_m \supset \dots$. Two chains of cross-cuts $\{\sigma_m\}$ and $\{\sigma'_k\}$ are called **equivalent** if, for every $m = 1, 2, \dots$, the domain d_m contains all domains d'_k except a finite number and, for every $k = 1, 2, \dots$, the domain d'_k contains all domains d_m except a finite number, too. A **prime end** K of the domain D is an equivalence class of chains of cross-cuts of D .

Reducing the case of bounded finitely connected domains to Theorem 9.3 in [7] for simple connected domains, we obtain one of the basic facts for applications of the theory of lower Q -homeomorphisms in what follows:

Lemma 1.1. *Each prime end P of a bounded finitely connected domain D in \mathbb{C} contains a chain of cross-cuts σ_m lying on circles $S(z_0, r_m) = \{z \in \mathbb{C} : |z - z_0| = r_m\}$ with $z_0 \in \partial D$ and $r_m \rightarrow 0$ as $m \rightarrow \infty$.*

Remark 1.1. *As known, every bounded finitely connected domain D in \mathbb{C} can be mapped by a conformal mapping g_0 onto the so-called circular domain D_0 whose boundary consists of a finite collection of mutually disjoint circles and isolated points, see, e.g., Theorem V.6.2 in [15]. Moreover, isolated singular points of bounded conformal mappings are removable by Theorem 1.2 in [7]. Hence isolated points of ∂D correspond to isolated points of ∂D_0 and inversely.*

Reducing this case to the Caratheodory theorem, see, e.g., Theorem 9.4 in [7] for simple connected domains, we have a natural one-to-one correspondence between points of ∂D_0 and prime ends of the domain D , see also Theorem 4.1 in [54]. Denote by \bar{D}_P the completion of D with its prime ends and determine in \bar{D}_P the metric $\rho_0(p_1, p_2) = |\bar{g}_0(p_1) - \bar{g}_0(p_2)|$ where \bar{g}_0 is the extension of g_0 .

If g_ is another conformal mapping of the domain D on a circular domain D_* , then the corresponding metric $\rho_*(p_1, p_2) = |\bar{g}_*(p_1) - \bar{g}_*(p_2)|$ generates the same convergence in \bar{D}_P as the metric ρ_0 because $g_0 \circ g_*^{-1}$ is a conformal mapping between the domains D_* and D_0 that is extended to a homeomorphism between \bar{D}_* and \bar{D}_0 . Consequently, the given metrics induce the same topology in the space \bar{D}_P that we will call the **topology of prime ends**.*

This topology can be also described in inner terms of the domain D , see, e.g., Section 9.5 in [7], however, we prefer the definition through the metrics because it is more clear, more convenient and it is important for us just metrizable of \bar{D}_P . Note also that the space \bar{D}_P for every bounded finitely connected domain D in \mathbb{C} with the given topology is compact because the closure of the circular domain D_0 is a compact space and by the construction $\bar{g}_0 : \bar{D}_P \rightarrow \bar{D}_0$ is a homeomorphism.

Later on, we mean the continuity of mappings $f : \bar{D}_P \rightarrow \bar{D}'_P$ with respect to the topology of prime ends.

2. On Lower Q -Homeomorphisms

A continuous mapping γ of an open subset Δ of the real axis \mathbb{R} or a circle into D is called a **dashed line**, see, e.g., Section 6.3 in [49]. Recall that every open set Δ in \mathbb{R} consists of a countable collection of mutually disjoint intervals. This is the motivation for the term.

Given a family Γ of dashed lines γ in complex plane \mathbb{C} , a Borel function $\varrho : \mathbb{C} \rightarrow [0, \infty]$ is called **admissible** for Γ , write $\varrho \in \text{adm } \Gamma$, if

$$\int_{\gamma} \varrho \, ds \geq 1 \quad \forall \gamma \in \Gamma \tag{2}$$

The **(conformal) modulus** of Γ is the quantity

$$M(\Gamma) = \inf_{\varrho \in \text{adm } \Gamma} \int_{\mathbb{C}} \varrho^2(z) \, dm(z) \tag{3}$$

where $dm(z)$ corresponds to the Lebesgue measure in \mathbb{C} . We say that a property P holds for **a.e.** (almost every) $\gamma \in \Gamma$ if a subfamily of all lines in Γ for which P fails has the modulus zero. Later on, we also say that a Lebesgue measurable function $\varrho : \mathbb{C} \rightarrow [0, \infty]$ is **extensively admissible** for Γ , write $\varrho \in \text{ext adm } \Gamma$, if (2) holds for a.e. $\gamma \in \Gamma$, see, e.g., Section 9.2 in [49].

The following concept was motivated by Gehring’s ring definition of quasiconformality in [10] and first introduced in the paper [26]. Given bounded domains D and D' in \mathbb{C} , $z_0 \in \overline{D}$, and a measurable function $Q : \mathbb{C} \rightarrow (0, \infty)$, we say that a homeomorphism $f : D \rightarrow D'$ is a **lower Q -homeomorphism at the point z_0** if

$$M(f(\Sigma_{\varepsilon})) \geq \inf_{\varrho \in \text{ext adm } \Sigma_{\varepsilon}} \int_{D \cap R_{\varepsilon}} \frac{\varrho^2(z)}{Q(z)} \, dm(z) \tag{4}$$

for every ring $R_{\varepsilon} = \{z \in \overline{\mathbb{C}} : \varepsilon < |z - z_0| < \varepsilon_0\}$, $\varepsilon \in (0, \varepsilon_0)$, $\varepsilon_0 \in (0, d_0)$, where $d_0 = \sup_{z \in D} |z - z_0|$, and Σ_{ε} denotes the family of all intersections of the circles $S(z_0, r) = \{z \in \mathbb{C} : |z - z_0| = r\}$, $r \in (\varepsilon, \varepsilon_0)$, with the domain D . We also say that a homeomorphism $f : D \rightarrow D'$ is a **lower Q -homeomorphism** if f is a lower Q -homeomorphism at every point $x_0 \in \overline{D}$.

Recall the criterion for homeomorphisms in \mathbb{C} to be lower Q -homeomorphisms, see Theorem 2.1 in [27], correspondingly, Theorem 9.2 in [49].

Proposition 2.1. *Let D and D' be bounded domains in \mathbb{C} , $z_0 \in \overline{D}$, and $Q : \mathbb{C} \rightarrow (0, \infty)$ be a measurable function. A homeomorphism $f : D \rightarrow D'$ is a lower Q -homeomorphism at z_0 if and only if*

$$M(f(\Sigma_{\varepsilon})) \geq \int_{\varepsilon}^{\varepsilon_0} \frac{dr}{\|Q\|(z_0, r)} \quad \forall \varepsilon \in (0, \varepsilon_0), \varepsilon_0 \in (0, d_0), \tag{5}$$

where $d_0 = \sup_{z \in D} |z - z_0|$ and $\|Q\|(z_0, r)$ is the L_1 -norm of Q over $D \cap S(z_0, r)$.

3. On Regular Domains

Recall first of all the following topological notion. A domain $D \subset \mathbb{C}$ is said to be **locally connected at a point** $z_0 \in \partial D$ if, for every neighborhood U of the point z_0 , there is a neighborhood $V \subseteq U$ of z_0 such that $V \cap D$ is connected. Note that every Jordan domain D in \mathbb{C} is locally connected at each point of ∂D , see e.g. [93], p. 66.

We say that ∂D is **weakly flat at a point** $z_0 \in \partial D$ if, for every neighborhood U of the point z_0 and every number $P > 0$, there is a neighborhood $V \subset U$ of z_0 such that

$$M(\Delta(E, F; D)) \geq P \tag{6}$$

for all continua E and F in D intersecting ∂U and ∂V . Here and later on, $\Delta(E, F; D)$ denotes the family of all paths $\gamma : [a, b] \rightarrow \overline{C}$ connecting E and F in D , i.e. $\gamma(a) \in E, \gamma(b) \in F$ and $\gamma(t) \in D$ for all $t \in (a, b)$. We say that the boundary ∂D is **weakly flat** if it is weakly flat at every point in ∂D .

We also say that a point $z_0 \in \partial D$ is **strongly accessible** if, for every neighborhood U of the point z_0 , there exist a compactum E in D , a neighborhood $V \subset U$ of z_0 and a number $\delta > 0$ such that

$$M(\Delta(E, F; D)) \geq \delta \tag{7}$$

for all continua F in D intersecting ∂U and ∂V . We say that the boundary ∂D is **strongly accessible** if every point $z_0 \in \partial D$ is strongly accessible.

It is easy to see that if a domain D in C is weakly flat at a point $z_0 \in \partial D$, then the point z_0 is strongly accessible from D . Moreover, it was proved by us that if a domain D in C is weakly flat at a point $z_0 \in \partial D$, then D is locally connected at z_0 , see, e.g., Lemma 5.1 in [27] or Lemma 3.15 in [49].

The notions of strong accessibility and weak flatness at boundary points of a domain in C defined in [26] are localizations and generalizations of the corresponding notions introduced in [47]–[48], cf. with the properties P_1 and P_2 by Väisälä in [89] and also with the quasiconformal accessibility and the quasiconformal flatness by Näkki in [53].

Recently we have proved the following significant statements, see either Theorem 10.1 (Lemma 6.1) in [27] or Theorem 9.8 (Lemma 9.4) in [49].

Proposition 3.1. *Let D and D' be bounded domains in C , $Q : D \rightarrow (0, \infty)$ a measurable function and $f : D \rightarrow D'$ a lower Q -homeomorphism in ∂D . Suppose that the domain D is locally connected on ∂D and that the domain D' has a (strongly accessible) weakly flat boundary. If*

$$\int_0^{\delta(z_0)} \frac{dr}{\|Q\|(z_0, r)} = \infty \quad \forall z_0 \in \partial D \tag{8}$$

for some $\delta(z_0) \in (0, d(z_0))$ where $d(z_0) = \sup_{z \in D} |z - z_0|$ and

$$\|Q\|(z_0, r) = \int_{S(z_0, r)} Q(z) ds,$$

then f has a (continuous) homeomorphic extension $\bar{f} : \overline{D} \rightarrow \overline{D}'$.

Bounded convex domains and bounded domains with smooth boundaries are simple examples of weakly flat boundaries, see a more detailed discussion, e.g., in [24].

4. On Continuous Extension of Lower Q -Homeomorphisms

Lemma 4.1. *Let D and D' be bounded finitely connected domains in C and let $f : D \rightarrow D'$ be a lower Q -homeomorphism. If*

$$\int_0^{\varepsilon_0} \frac{dr}{\|Q\|(z_0, r)} = \infty \quad \forall z_0 \in \partial D \tag{9}$$

where $0 < \varepsilon_0 < d_0 = \sup_{z \in D} |z - z_0|$ and

$$\|Q\|(z_0, r) = \int_{S(z_0, r)} Q(z) ds, \tag{10}$$

then f can be extended to a continuous mapping of \overline{D}_p onto \overline{D}'_p .

Proof. With no loss of generality we may assume that D' is a circular domain and, thus, $\overline{D'}_P = \overline{D'}$. By metrizable and compactness of $\overline{D'}$, it suffices to prove that, for each prime end P of the domain D , the cluster set

$$L = C(P, f) := \left\{ \zeta \in \mathbb{C} : \zeta = \lim_{n \rightarrow \infty} f(z_n), z_n \rightarrow P, z_n \in D, n = 1, 2, \dots \right\}$$

consists of a single point $\zeta_0 \in \partial D'$.

Note that $L \neq \emptyset$ by compactness of the set $\overline{D'}$, and L is a subset of $\partial D'$, see, e.g., Proposition 2.5 in [68] or Proposition 13.5 in [49]. Let us assume that there is at least two points ζ_0 and $\zeta_* \in L$. Set $U = B(\zeta_0, r_0) = \{\zeta \in \mathbb{C} : |\zeta - \zeta_0| < r_0\}$ where $0 < r_0 < |\zeta_* - \zeta_0|$.

Let $\sigma_k, k = 1, 2, \dots$, be a chain of cross-cuts of D in the prime end P lying on circles $S_k = S(z_0, r_k)$ from Lemma 1.1 where $z_0 \in \partial D$. Let $D_k, k = 1, 2, \dots$ be the domains associated with σ_k . Then there exist points ζ_k and ζ_k^* in the domains $D'_k = f(D_k)$ such that $|\zeta_0 - \zeta_k| < r_0$ and $|\zeta_0 - \zeta_k^*| > r_0$ and, moreover, $\zeta_k \rightarrow \zeta_0$ and $\zeta_k^* \rightarrow \zeta_*$ as $k \rightarrow \infty$. Let C_k be continuous curves joining ζ_k and ζ_k^* in D'_k . Note that by the construction $\partial U \cap C_k \neq \emptyset$.

By the condition of strong accessibility of the point ζ_0 , there is a continuum $E \subset D'$ and a number $\delta > 0$ such that

$$M(\Delta(E, C_k; D')) \geq \delta$$

for all large enough k .

Without loss of generality, we may assume that the latter condition holds for all $k = 1, 2, \dots$. Note that $C = f^{-1}(E)$ is a compact subset of D and hence $\varepsilon_0 = \text{dist}(z_0, C) > 0$. Again, with no loss of generality, we may assume that $r_k < \varepsilon_0$ for all $k = 1, 2, \dots$

Let Γ_m be a family of all continuous curves in $D \setminus D_m$ joining the circle $S_0 = S(z_0, \varepsilon_0)$ and $\overline{\sigma_m}, m = 1, 2, \dots$. Note that by the construction $C_k \subset D'_k \subset D'_m$ for all $m \leq k$ and, thus, by the principle of minorization $M(f(\Gamma_m)) \geq \delta$ for all $m = 1, 2, \dots$

On the other hand, the quantity $M(f(\Gamma_m))$ is equal to the capacity of the condenser in D' with facings $\overline{D'_m}$ and $f(D \setminus B_0)$ where $B_0 = B(z_0, \varepsilon_0)$, see, e.g., [80]. Thus, by the principle of minorization and Theorem 3.13 in [94]

$$M(f(\Gamma_m)) \leq \frac{1}{M(f(\Sigma_m))}$$

where Σ_m is the collection of all intersections of the domain D and the spheres $S(z_0, \rho), \rho \in (r_m, \varepsilon_0)$, because $f(\Sigma_m) \subset \Sigma(f(S_m), f(S_0))$ where $\Sigma(f(S_m), f(S_0))$ consists of all closed subsets of D' separating $f(S_m)$ and $f(S_0)$. Finally, by the condition (9) we obtain that $M(f(\Gamma_m)) \rightarrow 0$ as $m \rightarrow \infty$.

The obtained contradiction disproves the assumption that the cluster set $C(P, f)$ consists of more than one point. \square

5. Extension of Inverse Maps of Lower Q -Homeomorphisms

The base for the proof on extending the inverse mappings of lower Q -homeomorphism by prime ends in the plane is the following fact on the cluster sets.

Lemma 5.1. *Let D and D' be bounded finitely connected domains in \mathbb{C} , and let $f : D \rightarrow D'$ be a lower Q -homeomorphism. Denote by P_1 and P_2 different prime ends of the domain D and by $\sigma_m, m = 1, 2, \dots$, a chain of cross-cuts in the prime end P_1 from Lemma 1.1, lying on circles $S(z_1, r_m), z_1 \in \partial D$, with associated domains d_m . Suppose that Q is integrable over $D \cap S(z_1, r)$ for a set E of numbers $r \in (0, \delta)$ of a positive linear measure where $\delta = r_{m_0}$ and m_0 is a minimal number such that the domain d_{m_0} does not contain sequences of points converging to P_2 . If $\partial D'$ is weakly flat, then*

$$C(P_1, f) \cap C(P_2, f) = \emptyset. \tag{11}$$

Note that in view of metrizable of the completion \overline{D}_P of the domain D with prime ends, see Remark 1.1, the number m_0 in Lemma 5.1 always exists.

Proof. Let us choose $\varepsilon \in (0, \delta)$ such that $E_0 := \{r \in E : r \in (\varepsilon, \delta)\}$ has a positive linear measure. Such a choice is possible in view of subadditivity of the linear measure and the exhaustion $E = \cup E_m$ where $E_m = \{r \in E : r \in (1/m, \delta)\}$, $m = 1, 2, \dots$. Note that by Proposition 2.1

$$M(f(\Sigma_\varepsilon)) > 0 \tag{12}$$

where Σ_ε is the family of all surfaces $D(r) = D \cap S(z_1, r)$, $r \in (\varepsilon, \delta)$.

Let us assume that $C_1 \cap C_2 \neq \emptyset$ where $C_i = C(P_i, f)$, $i = 1, 2$. By the construction there is $m_1 > m_0$ such that σ_{m_1} lies on the circle $S(z_1, r_{m_1})$ with $r_{m_1} < \varepsilon$. Let $d_0 = d_{m_1}$ and $d_* \subseteq D \setminus d_{m_0}$ be a domain associated with a chain of cross-cuts in the prime end P_2 . Let $\zeta_0 \in C_1 \cap C_2$. Choose $r_0 > 0$ such that $S(\zeta_0, r_0) \cap f(d_0) \neq \emptyset$ and $S(\zeta_0, r_0) \cap f(d_*) \neq \emptyset$.

Set $\Gamma = \Delta(\overline{d_0}, \overline{d_*}; D)$. Correspondingly (12), by the principle of minorization and Theorem 3.13 in [94],

$$M(f(\Gamma)) \leq \frac{1}{M(f(\Sigma_\varepsilon))} < \infty. \tag{13}$$

Let $M_0 > M(f(\Gamma))$ be a finite number. By the condition of the lemma, $\partial D'$ is weakly flat and hence there is $r_* \in (0, r_0)$ such that

$$M(\Delta(E, F; D')) \geq M_0$$

for all continua E and F in D' intersecting the circles $S(\zeta_0, r_0)$ and $S(\zeta_0, r_*)$. However, these circles can be joined by continuous curves c_1 and c_2 in the domains $f(d_0)$ and $f(d_*)$, correspondingly, and, in particular, for these curves

$$M_0 \leq M(\Delta(c_1, c_2; D')) \leq M(f(\Gamma)). \tag{14}$$

The obtained contradiction disproves the assumption that $C_1 \cap C_2 \neq \emptyset$. \square

Theorem 5.1. *Let D and D' be bounded finitely connected domains in \mathbb{C} and $f : D \rightarrow D'$ be a lower Q -homeomorphism with $Q \in L^1(D)$. Then f^{-1} can be extended to a continuous mapping of \overline{D}'_P onto \overline{D}_P .*

Proof. By Remark 1.1, we may assume with no loss of generality that D' is a circular domain, $\overline{D}'_P = \overline{D}'$; $C(\zeta_0, f^{-1}) \neq \emptyset$ for every $\zeta_0 \in \partial D'$ because \overline{D}_P is metrizable and compact. Moreover, $C(\zeta_0, f^{-1}) \cap D = \emptyset$, see, e.g., Proposition 2.5 in [68] or Proposition 13.5 in [49].

Let us assume that there is at least two different prime ends P_1 and P_2 in $C(\zeta_0, f^{-1})$. Then $\zeta_0 \in C(P_1, f) \cap C(P_2, f)$ and, thus, (11) does not hold. Let $z_1 \in \partial D$ be a point corresponding to P_1 from Lemma 1.1. Note that

$$E = \{r \in (0, \delta) : Q|_{D \cap S(z_1, r)} \in L^1(D \cap S(z_1, r))\} \tag{15}$$

has a positive linear measure for every $\delta > 0$ by the Fubini theorem, see, e.g., [77], because $Q \in L^1(D)$. The obtained contradiction with Lemma 5.1 shows that $C(\zeta_0, f^{-1})$ contains only one prime end of D .

Thus, we have the extension g of f^{-1} to \overline{D}' such that $C(\partial D', f^{-1}) \subseteq \overline{D}_P \setminus D$. Really $C(\partial D', f^{-1}) = \overline{D}_P \setminus D$. Indeed, if P_0 is a prime end of D , then there is a sequence z_n in D being convergent to P_0 . We may assume without loss of generality that $z_n \rightarrow z_0 \in \partial D$ and $f(z_n) \rightarrow \zeta_0 \in \partial D'$ because \overline{D} and \overline{D}' are compact. Hence $P_0 \in C(\zeta_0, f^{-1})$.

Finally, let us show that the extended mapping $g : \overline{D}' \rightarrow \overline{D}_P$ is continuous. Indeed, let $\zeta_n \rightarrow \zeta_0$ in \overline{D}' . If $\zeta_0 \in D'$, then the statement is obvious. If $\zeta_0 \in \partial D'$, then take $\zeta_n^* \in D'$ such that $|\zeta_n - \zeta_n^*| < 1/n$ and $\rho(g(\zeta_n), g(\zeta_n^*)) < 1/n$ where ρ is one of the metrics in Remark 1.1. Note that by the construction $g(\zeta_n^*) \rightarrow g(\zeta_0)$ because $\zeta_n^* \rightarrow \zeta_0$. Consequently, $g(\zeta_n) \rightarrow g(\zeta_0)$, too. \square

Theorem 5.2. *Let D and D' be bounded finitely connected domains in \mathbb{C} . If $f : D \rightarrow D'$ is a lower Q -homeomorphism with condition (9), then f^{-1} can be extended to a continuous mapping of \overline{D}'_P onto \overline{D}_P .*

Proof. Indeed, by Lemma 9.2 in [27] or Lemma 9.6 in [49], condition (9) implies that

$$\int_0^\delta \frac{dr}{\|Q\|(z_0, r)} = \infty \quad \forall z_0 \in \partial D \quad \forall \delta \in (0, \varepsilon_0) \tag{16}$$

and, thus, the set

$$E = \{r \in (0, \delta) : Q|_{D \cap S(z_0, r)} \in L^1(D \cap S(z_0, r))\} \tag{17}$$

has a positive linear measure for all $z_0 \in \partial D$ and all $\delta \in (0, \varepsilon_0)$. The rest of arguments based on Lemma 5.1 is perfectly similar to one in the proof of Theorem 5.1. \square

6. On Functions of Finite Mean Oscillation

Recall that a real-valued function u in a domain D in \mathbb{C} is said to be of **bounded mean oscillation** in D , abbr. $u \in \text{BMO}(D)$, or simply $u \in \mathbf{BMO}$ if $u \in L^1_{\text{loc}}(D)$ and

$$\|u\|_* := \sup_B \frac{1}{|B|} \int_B |u(z) - u_B| dm(z) < \infty, \tag{18}$$

where the supremum is taken over all discs B in D , $dm(z)$ corresponds to the Lebesgue measure in \mathbb{C} and u_B is the average of u over B . We write $u \in \text{BMO}_{\text{loc}}(D)$ if $u \in \text{BMO}(U)$ for every relatively compact subdomain U of D (we also write BMO or BMO_{loc} if it is clear from the context what D is).

The class BMO was introduced by John and Nirenberg in the paper [23] and soon became an important concept in harmonic analysis, partial differential equations and related areas, see, e.g., [17] and [62].

A function φ in BMO is said to have **vanishing mean oscillation** (abbreviated as $\varphi \in \mathbf{VMO}$), if the supremum in (18) taken over all disks B in D with $|B| < \varepsilon$ converges to 0 as $\varepsilon \rightarrow 0$. \mathbf{VMO} was introduced by Sarason in [78]. Note that a large number of papers are devoted to the existence, uniqueness and properties of solutions for various kinds of differential equations and, in particular, of elliptic type with coefficients of the class \mathbf{VMO} , see, e.g., [6, 21, 51, 56, 61].

Following the paper [19], see also the monographs [16] and [49], we say that a function $\varphi : D \rightarrow \mathbb{R}$ has **finite mean oscillation** at a point $z_0 \in D$ if

$$\overline{\lim}_{\varepsilon \rightarrow 0} \int_{B(z_0, \varepsilon)} |\varphi(z) - \tilde{\varphi}_\varepsilon(z_0)| dm(z) < \infty, \tag{19}$$

where

$$\tilde{\varphi}_\varepsilon(z_0) = \int_{B(z_0, \varepsilon)} \varphi(z) dm(z) \tag{20}$$

is the mean value of the function $\varphi(z)$ over the disk $B(z_0, \varepsilon)$. Note that the condition (19) includes the assumption that φ is integrable in some neighborhood of the point z_0 . We say also that a function $\varphi : D \rightarrow \mathbb{R}$ is of **finite mean oscillation in D** , abbr. $\varphi \in \text{FMO}(D)$ or simply $\varphi \in \text{FMO}$, if $\varphi \in \text{FMO}(z_0)$ for all points $z_0 \in D$. We write $\varphi \in \text{FMO}(\overline{D})$ if φ is given in a domain G in \mathbb{C} such that $\overline{D} \subset G$ and $\varphi \in \text{FMO}(z_0)$ for all $z_0 \in \overline{D}$.

The following statement is obvious by the triangle inequality.

Proposition 6.1. *If, for a collection of numbers $\varphi_\varepsilon \in \mathbb{R}$, $\varepsilon \in (0, \varepsilon_0]$,*

$$\overline{\lim}_{\varepsilon \rightarrow 0} \int_{B(z_0, \varepsilon)} |\varphi(z) - \varphi_\varepsilon| dm(z) < \infty, \quad (21)$$

then φ is of finite mean oscillation at z_0 .

In particular choosing in Proposition 6.1, $\varphi_\varepsilon \equiv 0$, $\varepsilon \in (0, \varepsilon_0]$, we obtain the following conclusion.

Corollary 6.1. *If, for a point $z_0 \in D$,*

$$\overline{\lim}_{\varepsilon \rightarrow 0} \int_{B(z_0, \varepsilon)} |\varphi(z)| dm(z) < \infty, \quad (22)$$

then φ has finite mean oscillation at z_0 .

Recall that a point $z_0 \in D$ is called a **Lebesgue point** of a function $\varphi : D \rightarrow \mathbb{R}$ if φ is integrable in a neighborhood of z_0 and

$$\lim_{\varepsilon \rightarrow 0} \int_{B(z_0, \varepsilon)} |\varphi(z) - \varphi(z_0)| dm(z) = 0. \quad (23)$$

It is known that almost every point in D is a Lebesgue point for every function $\varphi \in L^1(D)$. Thus, we have by Proposition 6.1 the following corollary showing that the FMO condition is natural.

Corollary 6.2. *Every locally integrable function $\varphi : D \rightarrow \mathbb{R}$ has a finite mean oscillation at almost every point in D .*

Remark 6.1. *Note that the function $\varphi(z) = \log(1/|z|)$ belongs to BMO in the unit disk \mathbb{D} in \mathbb{C} , see, e.g., [60], p. 5, and hence also to FMO. However, $\widehat{\varphi}_\varepsilon(0) \rightarrow \infty$ as $\varepsilon \rightarrow 0$, showing that condition (22) is only sufficient but not necessary for a function φ to be of finite mean oscillation at z_0 . Clearly, $\text{BMO}(D) \subset \text{BMO}_{\text{loc}}(D) \subset \text{FMO}(D)$ and as well-known $\text{BMO}_{\text{loc}} \subset L^p_{\text{loc}}$ for all $p \in [1, \infty)$, see, e.g., [23]. However, FMO is not a subclass of L^p_{loc} for any $p > 1$ but only L^1_{loc} , see examples in [49], p. 211. Thus, the class FMO is essentially wider than BMO_{loc} .*

Versions of the next lemma has been first proved for the class BMO in the planar case in [72] and then in the space case in [48]. For the FMO case, see the papers [19] and [70] and also the monographs [16] and [49].

Lemma 6.1. *Let D be a domain in \mathbb{C} and let $\varphi : D \rightarrow \mathbb{R}$ be a non-negative function of the class $\text{FMO}(z_0)$ for some $z_0 \in D$. Then*

$$\int_{\varepsilon < |z-z_0| < \varepsilon_0} \frac{\varphi(z) dm(z)}{\left(|z-z_0| \log \frac{1}{|z-z_0|}\right)^2} = O\left(\log \log \frac{1}{\varepsilon}\right) \quad \text{as } \varepsilon \rightarrow 0 \quad (24)$$

for some $\varepsilon_0 \in (0, \delta_0)$ where $\delta_0 = \min(e^{-e}, d_0)$, $d_0 = \sup_{z \in D} |z - z_0|$.

7. Homeomorphic Extension of Lower Q -Homeomorphisms

Combining Lemma 4.1 and Theorem 5.2, we come to the following result.

Theorem 7.1. *Let D and D' be bounded finitely connected domains in \mathbb{C} and let $f : D \rightarrow D'$ be a lower Q -homeomorphism with*

$$\int_0^{\varepsilon_0} \frac{dr}{\|Q\|(z_0, r)} = \infty \quad \forall z_0 \in \partial D \quad (25)$$

where $0 < \varepsilon_0 < d_0 = \sup_{z \in D} |z - z_0|$ and

$$\|Q\|(z_0, r) = \int_{S(z_0, r)} Q(z) ds.$$

Then f can be extended to a homeomorphism of \overline{D}_P onto \overline{D}'_P .

Corollary 7.1. *In particular, the conclusion of Theorem 7.1 holds if*

$$q_{z_0}(r) = O\left(\log \frac{1}{r}\right) \quad \forall z_0 \in \partial D \tag{26}$$

as $r \rightarrow 0$ where $q_{z_0}(r)$ is the average of Q over the circle $|z - z_0| = r$.

Using Lemma 2.2 in [70], see also Lemma 7.4 in [49], by Theorem 7.1 we obtain the following general lemma that, in turn, makes possible to obtain new criteria in a great number.

Lemma 7.1. *Let D and D' be bounded finitely connected domains in \mathbb{C} and let $f : D \rightarrow D'$ be a lower Q -homeomorphism. Suppose that*

$$\int_{D(z_0, \varepsilon)} Q(z) \cdot \psi_{z_0, \varepsilon}^2(|z - z_0|) dm(z) = o\left(I_{z_0}^2(\varepsilon)\right) \quad \forall z_0 \in \partial D \tag{27}$$

as $\varepsilon \rightarrow 0$ where $D(z_0, \varepsilon) = \{z \in D : \varepsilon < |z - z_0| < \varepsilon_0\}$ for $0 < \varepsilon_0 < d(z_0) = \sup_{z \in D} |z - z_0|$ and where $\psi_{z_0, \varepsilon}(t) : (0, \infty) \rightarrow [0, \infty]$, $\varepsilon \in (0, \varepsilon_0)$, is a two-parameter family of measurable functions such that

$$0 < I_{z_0}(\varepsilon) := \int_{\varepsilon}^{\varepsilon_0} \psi_{z_0, \varepsilon}(t) dt < \infty \quad \forall \varepsilon \in (0, \varepsilon_0).$$

Then f can be extended to a homeomorphism of \overline{D}_P onto \overline{D}'_P .

Remark 7.1. *Note that (27) holds, in particular, if*

$$\int_{B(z_0, \varepsilon_0)} Q(z) \cdot \psi^2(|z - z_0|) dm(z) < \infty \quad \forall z_0 \in \partial D \tag{28}$$

where $B(z_0, \varepsilon_0) = \{z \in D : |z - z_0| < \varepsilon_0\}$ and where $\psi(t) : (0, \infty) \rightarrow [0, \infty]$ is a measurable function such that $I_{z_0}(\varepsilon) \rightarrow \infty$ as $\varepsilon \rightarrow 0$. In other words, for the extendability of f to a homeomorphism of \overline{D}_P onto \overline{D}'_P , it suffices the integrals in (28) to be convergent for some nonnegative function $\psi(t)$ that is locally integrable on $(0, \infty)$ but it has a non-integrable singularity at zero.

Note also that it is not only Lemma 7.1 follows from Theorem 7.1 but, inversely, Theorem 7.1 follows from Lemma 7.1, too. Indeed, for the function

$$\psi_{z_0}(t) = \begin{cases} 1/\|Q\|(z_0, t), & t \in (0, \varepsilon_0), \\ 0, & t \in [\varepsilon_0, \infty), \end{cases} \tag{29}$$

we have by the Fubini theorem that

$$\int_{S(z_0, \varepsilon)} Q(z) \cdot \psi_{z_0}^2(|z - z_0|) dm(z) = \int_{\varepsilon}^{\varepsilon_0} \frac{dr}{\|Q\|(z_0, r)}. \tag{30}$$

Thus, Theorem 7.1 is equivalent to Lemma 7.1 but each of them is sometimes more convenient for applications than another one.

Choosing in Lemma 7.1 $\psi(t) := \frac{1}{t \log 1/t}$ and applying Lemma 6.1, we obtain the next result.

Theorem 7.2. *Let D and D' be bounded finitely connected domains in \mathbb{C} and let $f : D \rightarrow D'$ be a lower Q -homeomorphism. If $Q(z)$ has finite mean oscillation at every point $z_0 \in \partial D$, then f can be extended to a homeomorphism of \overline{D}_P onto \overline{D}'_P .*

Corollary 7.2. *In particular, the conclusion of Theorem 7.2 holds if*

$$\overline{\lim}_{\varepsilon \rightarrow 0} \int_{B(z_0, \varepsilon)} Q(z) dm(z) < \infty \quad \forall z_0 \in \partial D \tag{31}$$

Corollary 7.3. *The conclusion of Theorem 7.2 holds if every point $z_0 \in \partial D$ is a Lebesgue point of the function $Q : \mathbb{C} \rightarrow (0, \infty)$.*

The next statement also follows from Lemma 7.1 under the choice $\psi(t) = 1/t$.

Theorem 7.3. *Let D and D' be bounded finitely connected domains in \mathbb{C} and $f : D \rightarrow D'$ be a lower Q -homeomorphism. If, for some $\varepsilon_0 = \varepsilon(z_0) > 0$,*

$$\int_{\varepsilon < |z-z_0| < \varepsilon_0} Q(z) \frac{dm(z)}{|z-z_0|^2} = o\left(\left[\log \frac{1}{\varepsilon}\right]^2\right) \quad \forall z_0 \in \partial D \tag{32}$$

as $\varepsilon \rightarrow 0$, then f can be extended to a homeomorphism of \overline{D}_P onto \overline{D}'_P .

Remark 7.2. *Choosing in Lemma 7.1 the function $\psi(t) = 1/(t \log 1/t)$ instead of $\psi(t) = 1/t$, (32) can be replaced by the weaker condition*

$$\int_{\varepsilon < |z-z_0| < \varepsilon_0} \frac{Q(z) dm(z)}{|z-z_0| \log \frac{1}{|z-z_0|}} = o\left(\left[\log \log \frac{1}{\varepsilon}\right]^2\right). \tag{33}$$

Similarly, (26) can be replaced by the condition

$$q_{z_0}(r) = O\left(\log \frac{1}{r} \log \log \frac{1}{r}\right). \tag{34}$$

Of course, we could give here the whole scale of the corresponding conditions of the logarithmic type using suitable functions $\psi(t)$.

Theorem 7.1 has a magnitude of other fine consequences, for instance:

Theorem 7.4. *Let D and D' be bounded finitely connected domains in \mathbb{C} and let $f : D \rightarrow D'$ be a lower Q -homeomorphism with*

$$\int_D \Phi(Q(z)) dm(z) < \infty \tag{35}$$

for a nondecreasing convex function $\Phi : [0, \infty) \rightarrow [0, \infty)$ such that

$$\int_{\delta_*}^{\infty} \frac{d\tau}{\tau \Phi^{-1}(\tau)} = \infty \tag{36}$$

for $\delta_* > \Phi(0)$. Then f is extended to a homeomorphism of \overline{D}_P onto \overline{D}'_P .

Indeed, by Theorem 3.1 and Corollary 3.2 in [76], (35) and (36) imply (25) and, thus, Theorem 7.4 is a direct consequence of Theorem 7.1.

Corollary 7.4. *In particular, the conclusion of Theorem 7.2 holds if, for some $\alpha > 0$,*

$$\int_D e^{\alpha Q(z)} dm(z) < \infty. \tag{37}$$

Remark 7.3. *By Theorem 2.1 in [76], see also Proposition 2.3 in [71], (36) is equivalent to every of the conditions from the following series:*

$$\int_{\delta}^{\infty} H'(t) \frac{dt}{t} = \infty, \quad \delta > 0, \tag{38}$$

$$\int_{\delta}^{\infty} \frac{dH(t)}{t} = \infty, \quad \delta > 0, \tag{39}$$

$$\int_{\delta}^{\infty} H(t) \frac{dt}{t^2} = \infty, \quad \delta > 0, \tag{40}$$

$$\int_0^{\Delta} H\left(\frac{1}{t}\right) dt = \infty, \quad \Delta > 0, \tag{41}$$

$$\int_{\delta_0}^{\infty} \frac{d\eta}{H^{-1}(\eta)} = \infty, \quad \delta_0 > H(0), \tag{42}$$

where

$$H(t) = \log \Phi(t). \tag{43}$$

Here the integral in (39) is understood as the Lebesgue–Stieltjes integral and the integrals in (38) and (40)–(42) as the ordinary Lebesgue integrals.

It is necessary to give one more explanation. From the right hand sides in the conditions (38)–(42) we have in mind $+\infty$. If $\Phi(t) = 0$ for $t \in [0, t_*]$, then $H(t) = -\infty$ for $t \in [0, t_*]$ and we complete the definition $H'(t) = 0$ for $t \in [0, t_*]$. Note, the conditions (39) and (40) exclude that t_* belongs to the interval of integrability because in the contrary case the left hand sides in (39) and (40) are either equal to $-\infty$ or indeterminate. Hence we may assume in (38)–(41) that $\delta > t_0$, correspondingly, $\Delta < 1/t_0$ where $t_0 := \sup_{\Phi(t)=0} t$, set $t_0 = 0$ if $\Phi(0) > 0$.

The most interesting of the above conditions is (40) that can be rewritten in the form:

$$\int_{\delta}^{\infty} \log \Phi(t) \frac{dt}{t^2} = \infty. \tag{44}$$

Finally, note that if a domain D in \mathbb{C} is locally connected on its boundary, then there is a natural one-to-one correspondence between prime ends of D and boundary points of D . Thus, if D and D' are in addition locally connected on their boundaries in theorems of Sections 4 and 5, then f is extended to a homeomorphism of \overline{D} onto \overline{D}' . We obtained before it similar results when $\partial D'$ was weakly flat which is a more strong condition than local connectivity of D' on its boundary, see, e.g., [24] and [25].

As known, every Jordan domain D in \mathbb{C} is locally connected on its boundary, see, e.g., [93], p. 66. It is easy to see, the latter implies that every bounded finitely connected domain D in \mathbb{C} whose boundary consists of mutually disjoint Jordan curves and isolated points is also locally connected on its boundary.

Inversely, every bounded finitely connected domain D in \mathbb{C} which is locally connected on its boundary has a boundary consisting of mutually disjoint Jordan curves and isolated points. Indeed, every such a domain D can be mapped by a conformal mapping f onto the so-called circular domain D_* bounded by a finite collection of mutually disjoint circles and isolated points, see, e.g., Theorem V.6.2 in [15], that is extended to a homeomorphism of \overline{D} onto \overline{D}_* .

Note also that, under every homeomorphism f between domains D and D' in $\overline{\mathbb{C}}$, there is a natural one-to-one correspondence between components of their boundaries ∂D and $\partial D'$, see, e.g., Lemma 5.3 in [19] or Lemma 6.5 in [49]. Thus, if a bounded domain D in \mathbb{C} is finitely connected and D' is bounded, then D' is finitely connected, too.

8. Boundary Behavior of Mappings with Finite Distortion

Recall that a homeomorphism f between domains D and D' in \mathbb{R}^n , $n \geq 2$, is called of **finite distortion** if $f \in W_{\text{loc}}^{1,1}$ and

$$\|f'(x)\|^n \leq K(x) \cdot J_f(x) \quad (45)$$

with some a.e. finite function K where $f'(x)$ denotes the Jacobian matrix of f at $x \in D$ if it exists, $J_f(x) = \det f'(x)$ is the Jacobian of f at x , and $\|f'(x)\|$ is the operator norm of $f'(x)$, i.e.,

$$\|f'(x)\| = \max\{|f'(x)h| : h \in \mathbb{R}^n, |h| = 1\}. \quad (46)$$

In the complex plane $\|f'\| = |f_z| + |f_{\bar{z}}|$ and $J_f = |f_z|^2 - |f_{\bar{z}}|^2$, i.e., (45) is equivalent to the condition $K_f(z) < \infty$ a.e., see (1).

First this notion was introduced on the plane for $f \in W_{\text{loc}}^{1,2}$ in the work [22]. Later on, this condition was replaced by $f \in W_{\text{loc}}^{1,1}$ but with the additional condition $J_f \in L_{\text{loc}}^1$ in the monograph [20]. The theory of the mappings with finite distortion had many successors, see many relevant references in the monographs [16, 18, 20] and [49]. They had as predecessors of the mappings with bounded distortion, see [63], and also [90], in other words, the quasiregular mappings, see, e.g., [17, 50, 64] and [91]. They are also closely connected to the earlier mappings with the bounded Dirichlet integral, see, e.g., the monographs [45, 84] and [85], and the mappings quasiconformal in the mean which had a rich history, see, e.g., [2]–[4], [11]–[14], [31]–[44], [57]–[59], [65]–[67], [82], [83], [86], [87], [95] and [96].

Note that the above additional condition $J_f \in L_{\text{loc}}^1$ in the definition of the mappings with finite distortion can be omitted for homeomorphisms. Indeed, for each homeomorphism f between domains D and D' in \mathbb{R}^n with the first partial derivatives a.e. in D , there is a set E of the Lebesgue measure zero such that f satisfies (N)-property by Lusin on $D \setminus E$ and

$$\int_A J_f(x) \, dm(x) = |f(A)| \quad (47)$$

for every Borel set $A \subset D \setminus E$, see, e.g., 3.1.4, 3.1.8 and 3.2.5 in [9].

On the basis of Theorem 1.1 and the corresponding results on lower Q -homeomorphisms in Sections 5 and 7, we obtain the following conclusions on the boundary behavior of mappings with finite distortion.

Theorem 8.1. *Let D and D' be bounded finitely connected domains in \mathbb{C} and $f : D \rightarrow \overline{D}'$ be a homeomorphism of finite distortion with $K_f \in L^1(D)$. Then f^{-1} can be extended to a continuous mapping of \overline{D}'_p onto \overline{D}_p .*

In fact, it is sufficient to assume in Theorem 8.1 that K_f is integrable only in a neighborhood of ∂D and even more weak conditions on K_f due to Lemma 5.1.

However, any degree of integrability of K_f cannot guarantee a continuous extension of the direct mappings f to the boundary, see an example in the proof of Proposition 6.3 in [49]. Conditions for it have a perfectly different nature. The principal related result is the following.

Theorem 8.2. *Let D and D' be bounded finitely connected domains in \mathbb{C} , $f : D \rightarrow D'$ be a homeomorphism of finite distortion with condition*

$$\int_0^{\varepsilon_0} \frac{dr}{\|K_f\|(z_0, r)} = \infty \quad \forall z_0 \in \partial D \tag{48}$$

where $0 < \varepsilon_0 < d_0 = \sup_{z \in D} |z - z_0|$ and

$$\|K_f\|(z_0, r) = \int_{S(z_0, r)} K_f(z) ds. \tag{49}$$

Then f can be extended to a homeomorphism of \overline{D}_p onto \overline{D}'_p .

Here we assume that K_f is extended by zero outside of the domain D .

Corollary 8.1. *In particular, the conclusion of Theorem 8.2 holds if*

$$k_{z_0}(r) = O\left(\log \frac{1}{r}\right) \quad \forall z_0 \in \partial D \tag{50}$$

as $r \rightarrow 0$ where $k_{z_0}(r)$ is the average of K_f over the circle $|z - z_0| = r$.

Lemma 8.1. *Let D and D' be bounded finitely connected domains in \mathbb{C} and let $f : D \rightarrow D'$ be a homeomorphism with finite distortion. Suppose that*

$$\int_{\varepsilon < |z - z_0| < \varepsilon_0} K_f(z) \cdot \psi_{z_0, \varepsilon}^2(|z - z_0|) dm(z) = o\left(I_{z_0}^2(\varepsilon)\right) \quad \forall z_0 \in \partial D \tag{51}$$

as $\varepsilon \rightarrow 0$ where $0 < \varepsilon_0 < \sup_{z \in D} |z - z_0|$ and where $\psi_{z_0, \varepsilon}(t) : (0, \infty) \rightarrow [0, \infty]$, $\varepsilon \in (0, \varepsilon_0)$, is a family of measurable functions such that

$$0 < I_{z_0}(\varepsilon) := \int_{\varepsilon}^{\varepsilon_0} \psi_{z_0, \varepsilon}(t) dt < \infty \quad \forall \varepsilon \in (0, \varepsilon_0).$$

Then f can be extended to a homeomorphism of \overline{D}_p onto \overline{D}'_p .

Theorem 8.3. *Let D and D' be bounded finitely connected domains in \mathbb{C} and let $f : D \rightarrow D'$ be a homeomorphism with finite distortion. If $K_f(z)$ has finite mean oscillation at every point $z_0 \in \partial D$, then f can be extended to a homeomorphism of \overline{D}_p onto \overline{D}'_p .*

In fact, here it is sufficient $K_f(z)$ to have a dominant of finite mean oscillation in a neighborhood of every point $z_0 \in \partial D$.

Corollary 8.2. *In particular, the conclusion of Theorem 8.3 holds if*

$$\overline{\lim}_{\varepsilon \rightarrow 0} \int_{B(z_0, \varepsilon)} K_f(z) dm(z) < \infty \quad \forall z_0 \in \partial D \tag{52}$$

Theorem 8.4. Let D and D' be bounded finitely connected domains in \mathbb{C} and let $f : D \rightarrow D'$ be a homeomorphism with finite distortion such that

$$\int_{\varepsilon < |z-z_0| < \varepsilon_0} K_f(z) \frac{dm(z)}{|z-z_0|^2} = o\left(\left[\log \frac{1}{\varepsilon}\right]^2\right) \quad \forall z_0 \in \partial D \tag{53}$$

Then f can be extended to a homeomorphism of \overline{D}_p onto \overline{D}'_p .

Remark 8.1. Choosing in Lemma 7.1 the function $\psi(t) = 1/(t \log 1/t)$ instead of $\psi(t) = 1/t$, (53) can be replaced by the weaker condition

$$\int_{\varepsilon < |z-z_0| < \varepsilon_0} \frac{K_f(z) dm(z)}{\left(|z-z_0| \log \frac{1}{|z-z_0|}\right)^2} = o\left(\left[\log \log \frac{1}{\varepsilon}\right]^2\right). \tag{54}$$

Of course, we could give here the whole scale of the corresponding conditions of the logarithmic type using suitable functions $\psi(t)$. In particular, condition (50) can be weakened with the help of Theorem 8.2 by condition

$$k_{z_0}(r) = O\left(\log \frac{1}{r} \log \log \frac{1}{r}\right). \tag{55}$$

Theorem 8.2 has a magnitude of other fine consequences, for instance:

Theorem 8.5. Let D and D' be bounded finitely connected domains in \mathbb{C} and let $f : D \rightarrow D'$ be a homeomorphism with finite distortion such that

$$\int_D \Phi(K_f(z)) dm(z) < \infty \tag{56}$$

for a nondecreasing convex function $\Phi : [0, \infty) \rightarrow [0, \infty)$ such that

$$\int_{\delta_*}^{\infty} \frac{d\tau}{\tau \Phi^{-1}(\tau)} = \infty \tag{57}$$

for $\delta_* > \Phi(0)$. Then f is extended to a homeomorphism of \overline{D}_p onto \overline{D}'_p .

Corollary 8.3. In particular, the conclusion of Theorem 8.5 holds if, for some $\alpha > 0$,

$$\int_D e^{\alpha K_f(z)} dm(z) < \infty. \tag{58}$$

Remark 8.2. Note that the condition (57) is not only sufficient but also necessary for a continuous extension to the boundary of the mappings f with integral restrictions of the form (56), see, e.g., Theorem 5.1 and Remark 5.1 in [29].

The boundary behavior of quasiconformal mappings in prime ends was well studied long ago, see, e.g., [53] and [54] and further references therein. However, this problem in prime ends remained open for mappings with finite distortion. Note that the case of regular boundaries was studied in [24] and [25] for the plane and in [30] for the space, cf. also [8] and [79]. Thus, all results in Section 8 are new.

9. Boundary Behavior of Finitely Bi-Lipschitz Mappings

Given a domain $D \subseteq \mathbb{C}$, following Section 5 in [28], see also Section 10.6 in [49], we say that a mapping $f : D \rightarrow \mathbb{C}$ is **finitely bi-Lipschitz** if

$$0 < l(z, f) \leq L(z, f) < \infty \quad \forall z \in D \tag{59}$$

where

$$L(z, f) = \limsup_{\zeta \rightarrow z} \frac{|f(\zeta) - f(z)|}{|\zeta - z|} \tag{60}$$

and

$$l(z, f) = \liminf_{\zeta \rightarrow z} \frac{|f(\zeta) - f(z)|}{|\zeta - z|}. \tag{61}$$

The class of finitely bi-Lipschitz homeomorphisms is a natural generalization of the well-known classes of isometries and quasi-isometries. However, they are generally speaking are not of finite distortion by Iwaniec.

To show the latter, let us give the corresponding example. Namely, let us consider the mappings $f : \mathbb{C} \rightarrow \mathbb{C}$ of the form $f(z) = ze^{i\varphi(|z|)}$, $f(0) = 0$, with real-valued decreasing functions $\varphi(t)$, $t > 0$, of the class C^1 such that $\varphi(t) \rightarrow +\infty$ as $t \rightarrow 0$ and demonstrate that, for instance, the function $\varphi(t) = 1/t^2$ is suitable for our goals.

Indeed, first of all it is easy to see that f is a finitely bi-Lipschitz homeomorphism of \mathbb{C} onto itself such that $|f(z)| = |z|$ transforming the radial lines $\arg z = \text{const}$ into the corresponding spirals. Moreover, for $z = re^{i\vartheta}$,

$$\begin{aligned} \frac{\partial f}{\partial r} &= \frac{\partial f}{\partial z} \cdot \frac{\partial z}{\partial r} + \frac{\partial f}{\partial \bar{z}} \cdot \frac{\partial \bar{z}}{\partial r} = \frac{z}{|z|} \cdot f_z + \frac{\bar{z}}{|z|} \cdot f_{\bar{z}}, \\ \frac{\partial f}{\partial \vartheta} &= \frac{\partial f}{\partial z} \cdot \frac{\partial z}{\partial \vartheta} + \frac{\partial f}{\partial \bar{z}} \cdot \frac{\partial \bar{z}}{\partial \vartheta} = i(zf_z - \bar{z}f_{\bar{z}}), \end{aligned}$$

and, thus,

$$\begin{aligned} zf_z &= \frac{1}{2} \left(r \frac{\partial f}{\partial r} - i \frac{\partial f}{\partial \vartheta} \right) = f(z) \cdot \left(1 + i \cdot \frac{r\varphi'(r)}{2} \right), \\ \bar{z}f_{\bar{z}} &= \frac{1}{2} \left(r \frac{\partial f}{\partial r} + i \frac{\partial f}{\partial \vartheta} \right) = f(z) \cdot i \cdot \frac{r\varphi'(r)}{2}, \end{aligned}$$

i.e.

$$|f_z| = \left| \frac{r\varphi'(r)}{2} \right|, \quad |f_{\bar{z}}| = \left| 1 + i \cdot \frac{r\varphi'(r)}{2} \right| = \sqrt{1 + \left(\frac{r\varphi'(r)}{2} \right)^2}.$$

By the way, it follows from here that its Jacobian $J_f(z) = |f_z|^2 - |f_{\bar{z}}|^2 = 1 > 0$ and, consequently, that f preserves the orientation as well as the area. However, it is more important that $|f_z| > |f_{\bar{z}}| = 1/r^2$ and, thus, the first derivatives of f is not integrable in a neighborhood of the origin under the choice of $\varphi(t) = 1/t^2$.

By the classic Stepanov theorem, see [81], see also [46], we obtain from the right hand inequality in (59) that finitely bi-Lipschitz mappings are differentiable a.e. and from the left hand inequality in (59) that $J_f(x) \neq 0$ a.e. Moreover, such mappings have (N)-property with respect to each Hausdorff measure, see, e.g., either Lemma 5.3 in [28] or Lemma 10.6 [49]. In particular, they are ACL but generally speaking not in the class $W_{loc}^{1,1}$ as in the above example.

However, by Corollary 5.15 in [28] and Corollary 10.10 in [49] we have the following statement:

Lemma 9.1. *Every finitely bi-Lipschitz homeomorphism $f : \Omega \rightarrow \mathbb{C}$ is a lower Q -homeomorphism with $Q = K_f$.*

Corollary 9.1. *All results on homeomorphisms with finite distortion in Section 8 are valid for finitely bi-Lipschitz homeomorphisms under the same conditions on K_f .*

All these results for finitely bi-Lipschitz homeomorphisms are perfectly similar to the corresponding results for homeomorphisms with finite distortion in Section 8. Hence we will not formulate them in the explicit form here.

The results of this paper can be applied to the theory of boundary value problems for the Beltrami equations including equations of the second kind that take an important part in many problems of mathematical physics.

Acknowledgements

We would like to thank our referee whose remarks have made possible us essentially to improve the text.

References

- [1] T. Adamowicz, A. Björna, J. Björna, N. Shanmugalingamb, Prime ends for domains in metric spaces, *Advances in Mathematics* 238 (2013) 459–505.
- [2] L. Ahlfors, On quasiconformal mappings, *J. Anal. Math.* 3 (1953/1954) 1–58.
- [3] P.P. Belinskii, General properties of quasiconformal mappings, Novosibirsk, Izdat. “Nauka” Sibirsk. Otdel., 1974 [in Russian].
- [4] P.A. Biluta, Extremal problems for mappings quasiconformal in the mean, *Sib. Mat. Zh.* 6 (1965) 717–726 [in Russian].
- [5] C. Caratheodory, Über die Begrenzung der einfach zusammenhängender Gebiete, *Math. Ann.* 73 (1913) 323–370.
- [6] F. Chiarenza, M. Frasca, P. Longo, $W^{2,p}$ -solvability of the Dirichlet problem for nondivergence elliptic equations with VMO coefficients, *Trans. Amer. Math. Soc.* 336, no. 2 (1993) 841–853.
- [7] E. F. Collingwood, A. J. Lohwater, *The Theory of Cluster Sets*, Cambridge Tracts in Math. and Math. Physics 56 (1966), Cambridge: Cambridge Univ. Press.
- [8] M. Cristea, Mappings of finite distortion: boundary extension, *Rev. Roumaine Math. Pures Appl.* 51, no. 5-6 (2006) 607–631.
- [9] H. Federer, *Geometric Measure Theory*, Berlin: Springer-Verlag, 1969.
- [10] F.W. Gehring, Rings and quasiconformal mappings in space, *Trans. Amer. Math. Soc.* 103, no. 3 (1962) 353–393.
- [11] A. Golberg, Distortions of p -moduli of separating sets under mappings with finite integral dilatations, *Bull. Soc. Sci. Lettres Łódź Sér. Rech. Déform.* 40 (2003) 41–51.
- [12] A. Golberg, Homeomorphisms with finite mean dilatations, *Contemporary Math.*, 382 (2005) 177–186.
- [13] A. Golberg, Coefficients of quasiconformality of ring domains, *Proc. Inst. Math. NAS Ukraine* 1(3) (2004) , 77–86.
- [14] A. Golberg, V.S. Kud’yavin, Mean coefficients of quasiconformality of pair of domains, *Ukrain. Mat. Zh.* 43(12) (1991), 1709–1712 [in Russian]; translation in *Ukrain. Math. J.* 43 (1991) 1594–1597.
- [15] G. M. Goluzin, *Geometric Theory of Functions of a Complex Variable*, Transl. of Math. Monographs, 26, Providence, AMS, 1969.
- [16] V. Gutlyanskii, V. Ryazanov, U. Srebro, E. Yakubov, *The Beltrami Equations: A Geometric Approach*, *Developments in Math.* 26, New York etc., Springer, 2012.
- [17] J. Heinonen, T. Kilpelainen, O. Martio, *Nonlinear Potential Theory of Degenerate Elliptic Equations*, Oxford-New York-Tokio, Clarendon Press, 1993.
- [18] S. Hencl, P. Koskela, *Lectures on mappings of finite distortion*, *Lecture Notes in Math.* 2096, Cham, Springer, 2014.
- [19] A. Ignat’ev, V. Ryazanov, Finite mean oscillation in the mapping theory, *Ukr. Mat. Vis.* 2, no. 3 (2005), 395–417, 443 [in Russian]; transl. in *Ukr. Math. Bull.* 2, no. 3 (2005), 403–424, 443.
- [20] T. Iwaniec, G. Martin, *Geometric function theory and non-linear analysis*, Oxford, Oxford Univ. Press, 2001.
- [21] T. Iwaniec, C. Sbordone, Riesz transforms and elliptic PDEs with VMO coefficients, *J. Anal. Math.* 74 (1998), 183–212.
- [22] T. Iwaniec, V. Sverak, On mappings with integrable dilatation, *Proc. Amer. Math. Soc.* 118 (1993), 181–188.
- [23] F. John, L. Nirenberg, On functions of bounded mean oscillation. *Comm. Pure Appl. Math.* 14 (1961), 415–426.
- [24] D. Kovtonyuk, I. Petkov, V. Ryazanov, On the boundary behaviour of solutions to the Beltrami equations, *Complex Variables and Elliptic Equations* 58, no. 5 (2013), 647–663.
- [25] D. Kovtonyuk, I. Petkov, V. Ryazanov, R.R. Salimov, Boundary behavior and Dirichlet problem for Beltrami equations, *Algebra and Analysis* 25, no. 4 (2013), 101–124 [in Russian]; transl in *St. Petersburg Math. J.* 25 (2014), 587–603.
- [26] D. Kovtonyuk, V. Ryazanov, On boundaries of space domains, *Proc. Inst. Appl. Math. & Mech. NAS of Ukraine.* 13 (2006), 110–120.
- [27] D. Kovtonyuk, V. Ryazanov, Toward the theory of lower Q -homeomorphisms, *Ukr. Mat. Visn.* 5, no. 2 (2008), 159–184 [in Russian]; transl. in *Ukrainian Math. Bull.* 5, no. 2 (2008), 157–181.
- [28] D. Kovtonyuk, V. Ryazanov, On the theory of mappings with finite area distortion, *J. Anal. Math.* 104 (2008), 291–306.
- [29] D. Kovtonyuk, V. Ryazanov, On the boundary behavior of generalized quasi-isometries, *J. Anal. Math.* 115 (2011), 103–119.

- [30] D.A. Kovtonyuk, V.I. Ryazanov, R.R. Salimov, E.A. Sevost'yanov, Toward the theory of classes of Orlicz–Sobolev, *Algebra and Analysis*. 25, no. 6 (2013), 49–101 [in Russian]; transl in *St. Petersburg Math. J.* 25, no. 6 (2014), 929–963.
- [31] V.I. Kruglikov, The existence and uniqueness of mappings that are quasiconformal in the mean, In: *Metric Questions of the Theory of Functions and Mappings*, Kiev, Naukova Dumka, 1973, 123–147 [in Russian].
- [32] V.I. Kruglikov, Capacities of condensers and quasiconformal in the mean mappings in space, *Mat. Sb.* 130(2) (1986), 185–206 [in Russian].
- [33] V.I. Kruglikov, V.M. Miklyukov, On classes of plane topological mappings with generalized derivatives, In: *Metric Questions of the Theory of Functions and Mappings*, Kiev, Naukova dumka, 1973, 102–122 [in Russian].
- [34] V.I. Kruglikov, V.I. Paikov, Capacities and prime ends of an n -dimensional domain, *Dokl. Akad. Nauk Ukrain. SSR Ser. A.* 5 (1987), 10–13 [in Russian].
- [35] S.L. Krushkal', On mappings quasiconformal in the mean, *Dokl. Akad. Nauk SSSR.* 157(3) (1964), 517–519 [in Russian].
- [36] S.L. Krushkal', On the absolute integrability and differentiability of some classes of mappings of many-dimensional domains, *Sib. Mat. Zh.* 6(3) (1965), 692–696 [in Russian].
- [37] S.L. Krushkal', On mappings ε -quasiconformal in the mean, *Sib. Mat. Zh.* 8(4) (1967), 798–806 [in Russian].
- [38] S.L. Krushkal', R. Kühnau, Quasiconformal mappings, new methods and applications, Novosibirsk, Nauka, 1984 [in Russian].
- [39] V.S. Kud'yavin, A characteristic property of a class of n -dimensional homeomorphisms, *Dokl. Akad. Nauk Ukrain. SSR ser. A.* 3 (1990), 7–9 [in Russian].
- [40] V.S. Kud'yavin, Quasiconformal mappings and α -moduli of families of curves, *Dokl. Akad. Nauk Ukrain.* 7 (1992), 11–13 [in Russian].
- [41] V.S. Kud'yavin, Estimation of the distortion of distances under mappings quasiconformal in the mean, *Dinam. Sploshn. Sred.* 52 (1981), 168–171 [in Russian].
- [42] V.S. Kud'yavin, Local boundary properties of mappings quasiconformal in the mean, In: *Collection of Scientific Works, Institute of Mathematics, Novosibirsk, Siberian Branch of the Academy of Sciences of the USSR*, 1981, 168–171 [in Russian].
- [43] V.S. Kud'yavin, Behavior of a class of mappings quasiconformal in the mean at an isolated singular point, *Dokl. Akad. Nauk SSSR.* 277(5) (1984), 1056–1058 [in Russian].
- [44] R. Kühnau, Über Extremalprobleme bei im Mittel quasikonformen Abbildungen, *Lecture Notes in Math.* 1013 (1983), 113–124 [in German].
- [45] J. Lelong-Ferrand, *Representation conforme et transformations à integrale de Dirichlet bornée*, Paris, Gauthier-Villars, 1955.
- [46] J. Maly, A simple proof of the Stepanov theorem on differentiability almost everywhere, *Exposition Math.* 17 (1999), 59–61.
- [47] O. Martio, V. Ryazanov, U. Srebro, E. Yakubov, Q -homeomorphisms. *Contemporary Math.* 364 (2004), 193–203.
- [48] O. Martio, V. Ryazanov, U. Srebro, E. Yakubov, On Q -homeomorphisms. *Ann. Acad. Sci. Fenn.* 30 (2005), 49–69.
- [49] O. Martio, V. Ryazanov, U. Srebro, E. Yakubov, *Moduli in Modern Mapping Theory*, Springer Monographs in Mathematics, New York etc., Springer, 2009.
- [50] O. Martio, S. Rickman, J. Vaisala, Definitions for quasiregular mappings, *Ann. Acad. Sci. Fenn. Ser. A1. Math.* 448 (1969), 1–40.
- [51] O. Martio, V. Ryazanov, M. Vuorinen, BMO and Injectivity of Space Quasiregular Mappings, *Math. Nachr.* 205 (1999), 149–161.
- [52] V.M. Miklyukov, G.D. Suvorov, On existence and uniqueness of quasiconformal mappings with unbounded characteristics, In: *Investigations in the Theory of Functions of Complex Variables and its Applications*, Kiev, Inst. Math., 1972 [in Russian].
- [53] R. Näkki, Boundary behavior of quasiconformal mappings in n -space, *Ann. Acad. Sci. Fenn. Ser. A1. Math.* 484 (1970), 1–50.
- [54] R. Näkki, Prime ends and quasiconformal mappings, *J. Anal. Math.* 35 (1979), 13–40.
- [55] M. Ohtsuka, *Extremal Length and Precise Functions*, Tokyo, Gakkotosho Co., 2003.
- [56] D.K. Palagachev, Quasilinear elliptic equations with VMO coefficients, *Trans. Amer. Math. Soc.* 347, no. 7 (1995), 2481–2493.
- [57] M. Perovich, Isolated singularity of the mean quasiconformal mappings, *Lect. Notes in Math.* 743 (1979), 212–214.
- [58] M. Perovich, Global homeomorphism of mappings quasiconformal in the mean, *Dokl. Akad. Nauk SSSR.* 230(4) (1976), 781–784 [in Russian].
- [59] I.N. Pesin, Mappings quasiconformal in the mean, *Dokl. Akad. Nauk SSSR.* 187(4) (1969), 740–742 [in Russian].
- [60] T. Rado, P.V. Reichelderfer, *Continuous Transformations in Analysis*, Berlin etc., Springer, 1955.
- [61] M.A. Ragusa, Elliptic boundary value problem in vanishing mean oscillation hypothesis, *Comment. Math. Univ. Carolin.* 40, no. 4 (1999), 651–663.
- [62] H.M. Reimann, T. Rychener, *Funktionen Beschränkter Mittlerer Oscillation*, Lecture Notes in Math. 487, Berlin etc., Springer-Verlag, 1975.
- [63] Yu.G. Reshetnyak, *Space mappings with bounded distortion*, Novosibirsk, Nauka, 1982; transl. in *Translations of Mathematical Monographs*, 73, Providence, RI, Amer. Math. Soc., 1988.
- [64] S. Rickman, *Quasiregular Mappings*, Berlin etc., Springer-Verlag, 1993.
- [65] V.I. Ryazanov, On compactification of classes with integral restrictions on the Lavrent'ev characteristics, *Sibirsk. Mat. Zh.* 33(1) (1992), 87–104 [in Russian]; transl. in *Siberian Math. J.* 33(1) (1992), 70–86.
- [66] V.I. Ryazanov, On quasiconformal mappings with measure restrictions, *Ukrain. Mat. Zh.* 45(7) (1993), 1009–1019 [in Russian]; transl. in *Ukrainian Math. J.* 45(7) (1993), 1121–1133.
- [67] V.I. Ryazanov, On mappings that are quasiconformal in the mean, *Sibirsk. Mat. Zh.* 37(2) (1996), 378–388 [in Russian]; transl. in *Siberian Math. J.* 37(2) (1996), 325–334.
- [68] V.I. Ryazanov, R.R. Salimov, Weakly flat spaces and boundaries in the mapping theory, *Ukr. Mat. Vis.* 4, 2 (2007), 199–234 [in Russian]; transl. in *Ukrainian Math. Bull.* 4, no. 2 (2007), 199–233.
- [69] V. Ryazanov, R. Salimov, U. Srebro, E. Yakubov, On Boundary Value Problems for the Beltrami Equations, *Contemporary Math.* 591 (2013), 211–242.
- [70] V. Ryazanov, E. Sevost'yanov, Equicontinuous classes of ring Q -homeomorphisms, *Sibirsk. Math. Zh.* 48, 6 (2007), 1361–1376 [in Russian]; transl. in *Siberian Math. J.* 48, no. 6 (2007), 1093–1105.

- [71] V. Ryazanov, E. Sevost'yanov, Equicontinuity of mappings quasiconformal in the mean, *Ann. Acad. Sci. Fenn.* 36 (2011), 231–244.
- [72] V. Ryazanov, U. Srebro, E. Yakubov, On ring solutions of Beltrami equation, *J. Anal. Math.* 96 (2005), 117–150.
- [73] V. Ryazanov, U. Srebro, E. Yakubov, To strong ring solutions of the Beltrami equations, *Uzbek. Math. J.* 1 (2009), 127–137.
- [74] V. Ryazanov, U. Srebro, E. Yakubov, On strong solutions of the Beltrami equations, *Complex Var. Elliptic Equ.* 55, no. 1–3 (2010), 219–236.
- [75] V. Ryazanov, U. Srebro, E. Yakubov, Integral conditions in the theory of the Beltrami equations, *Complex Var. Elliptic Equ.* 57, no. 12 (2012), 1247–1270.
- [76] V. Ryazanov, U. Srebro, E. Yakubov, On integral conditions in the mapping theory, *Ukrainian Math. Bull.* 7, no. 1 (2010), 73–87.
- [77] S. Saks, *Theory of the Integral*, New York, Dover Publications Inc., 1964.
- [78] D. Sarason, Functions of vanishing mean oscillation, *Trans. Amer. Math. Soc.* 207 (1975), 391–405.
- [79] E.A. Sevost'yanov, On the boundary behavior of open discrete mappings with unbounded characteristic, *Ukrainian Math. J.* 64, no. 6 (2012), 979–984.
- [80] V.A. Shlyk, On the equality between p -capacity and p -modulus, *Sibirsk. Mat. Zh.* 34, no. 6 (1993), 216–221 [in Russian]; transl. in *Siberian Math. J.* 34, no. 6 (1993), 1196–1200.
- [81] W. Stepanov, Sur les conditions de l'existence de la differentielle totale, *Mat. Sb.* 32 (1925), 511–526.
- [82] Y.F. Strugov, Compactness of the classes of mappings quasiconformal in the mean, *Dokl. Akad. Nauk SSSR.* 243(4) (1978), 859–861 [in Russian].
- [83] Y.F. Strugov, A.V. Sychov, On different classes of mappings quasiconformal in the mean, *Vest. PANI.* 7 (2002), 14–19 [in Russian].
- [84] G.D. Suvorov, Generalized principle of the length and area in the mapping theory, *Kiev, Naukova Dumka*, 1985 [in Russian].
- [85] G.D. Suvorov, The metric theory of prime ends and boundary properties of plane mappings with bounded Dirichlet integrals, *Kiev, Naukova Dumka*, 1981 [in Russian].
- [86] A. Uhlov, S. Vodopyanov, Mappings associated with weighted Sobolev spaces, *Complex Anal. Dynam. Sys. III. Contemp. Math.* 455 (2008), 363–382.
- [87] A. Uhlov, S. Vodopyanov, Sobolev spaces and (P, Q) -quasiconformal mappings of Carnot groups, *Siberian Math. J.* 39 (1998), 665–682.
- [88] A. Vasil'ev, *Moduli of Families of Curves for Conformal and Quasiconformal Mappings*, Lecture Notes in Math. 1788, Berlin etc., Springer–Verlag, 2002.
- [89] J. Väisälä, *Lectures on n -Dimensional Quasiconformal Mappings*, Lecture Notes in Math. 229, Berlin etc., Springer–Verlag, 1971.
- [90] S. Vodop'yanov, Mappings with bounded distortion and with finite distortion on Carnot groups, *Sibirsk. Mat. Zh.* 40, no. 4 (1999), 764–804; transl. in *Siberian Math. J.* 40, no. 4 (1999), 644–677.
- [91] M. Vuorinen, *Conformal Geometry and Quasiregular Mappings*, Lecture Notes in Math. 1319, Berlin etc., Springer–Verlag, 1988.
- [92] G.Th. Whyburn, *Analytic Topology*, Providence, AMS, 1942.
- [93] R.L. Wilder, *Topology of Manifolds*, New York, AMS, 1949.
- [94] W.P. Ziemer, Extremal length and conformal capacity, *Trans. Amer. Math. Soc.* 126, no. 3 (1967), 460–473.
- [95] V.A. Zorich, Admissible order of growth of the characteristic of quasiconformality in the Lavrent'ev theory, *Dokl. Akad. Nauk SSSR.* 181 (1968), 530–533 [in Russian].
- [96] V.A. Zorich, Isolated singularities of mappings with unbounded distortion, *Mat. Sb.* 81 (1970), 634–638 [in Russian].