



Two Iterative Algorithms for Solving the Split Common Fixed Point Problems

Dingping Wu^a, Mihai Postolache^{b,c,d,e}

^aCollege of Applied Mathematics, Chengdu University of Information Technology, Chengdu, China

^bCenter for General Education, China Medical University, Taichung 40402, Taiwan

^cDepartment of Interior Design, Asia University, Taichung 41354, Taiwan

^dRomanian Academy, Gh. Mihoc-C. Iacob Institute of Mathematical Statistics and Applied Mathematics, Bucharest 050711, Romania

^eUniversity "Politehnica" of Bucharest, Department of Mathematics and Informatics Bucharest 060042, Romania

Abstract. The purpose of this paper is to study the split common fixed point problems (SCFP) involved in nonexpansive mappings in real Hilbert space. We introduce two iterative algorithms for finding a solution of the SCFP involved in nonexpansive mappings, where one is a Mann-type iterative algorithm and another is a Halpern-type iterative algorithm.

1. Introduction

In 1994, the split feasibility problem (SFP) was first presented by Censor and Elfving [7] for modeling inverse problems, which arise from phase retrievals and in medical image reconstruction ([1–3]). Recently, the SFP had also been applied to study intensity modulated radiation therapy when one attempted to describe physical dose constraints and equivalent uniform dose constraints within a single model([6]).

To begin with, let us recall that the SFP is to find a point

$$x^* \in C \text{ such that } Ax^* \in Q, \quad (1)$$

where C is a close convex subset of a Hilbert space H_1 , Q is a close convex subset of a Hilbert space H_2 and $A: H_1 \rightarrow H_2$ is a bounded linear operator. To solve problem (1), Byrne [2] proposed his CQ algorithm, which generates a sequence $\{x_n\}$ by

$$x_{n+1} = P_C(I + \gamma A^*(P_Q - I)A)x_n, \quad n \geq 0, \quad (2)$$

where $\gamma \in (0, 2/\lambda)$ and λ is the spectral radius of the operator A^*A .

Many authors have also made a continuation of the study on the CQ algorithm and its variant form, refer to [11, 14, 17, 18, 22, 27, 30].

2010 *Mathematics Subject Classification.* 47J25; 47H09; 65J15; 90C25.

Keywords. Split common fixed point problems, Nonexpansive mappings, Halpern-type algorithm, Mann-type algorithm.

Received: 11 January 2020; Accepted: 06 August 2020

Communicated by Naseer Shahzad

Corresponding author: Mihai Postolache

Email addresses: wdp68@163.com (Dingping Wu), mihai@mathem.pub.ro (Mihai Postolache)

On the other hand, the split common fixed point problem (SCFP) is a generalization of the SFP. The purpose of this paper is to study the SCFP, that is, finding a point x^* with the property

$$x^* \in \text{Fix}(T) \text{ and } Ax^* \in \text{Fix}(S). \tag{3}$$

This problem was first introduced by Censor and Segal [8]. We use Γ_0 to denote the set of the solutions of (3), that is, $\Gamma_0 = \{x^* : x^* \in \text{Fix}(T), Ax^* \in \text{Fix}(S)\}$. For solving (3), they suggested an algorithm, which generates a sequence $\{x_n\}$ in the following manner:

$$x_{n+1} = T(x_n + \gamma A^*(S - I)Ax_n), \quad n \geq 0. \tag{4}$$

Note that (4) is more general than (1). Some further generation of this algorithm were studied by [4, 5, 12–29, 31–34] and others. One of them, Moudafi extended (4) to the following relaxed algorithm:

$$x_{n+1} = U_{\alpha_n}(x_n + \gamma A^*(T_\beta - I)Ax_n), \quad n \geq 0,$$

where $\beta \in (0, 1)$, $\alpha_n \in (0, 1)$ are relaxation parameters. Using the definition of relaxed operators, observe that this algorithm takes the following form:

$$\begin{cases} y_n = x_n + \gamma \beta A^*(T - I)Ax_n, \\ x_{n+1} = (1 - \alpha_n)y_n + \alpha_n U y_n, \quad n \geq 0, \end{cases}$$

Recently, Chen et. al. [10] presented the following iterative method to have a common fixed point of three nonexpansive mappings.

$$\begin{cases} x_0 \in C \text{ chosen arbitrarily,} \\ y_n = \beta_n x_n + (1 - \beta_n)Tx_n, \\ z_n = \gamma_n x_n + (1 - \gamma_n)Sx_n, \\ x_{n+1} = \alpha_n y_n + (1 - \alpha_n)Rz_n, \quad n \geq 0, \end{cases}$$

where $\beta_n \in (0, 1)$, $\gamma_n \in (0, 1)$, $\alpha_n \in (0, 1)$.

Motivated and inspired by the above works and the research going on in these direction, the purpose here is to find a solution of the SCFP involved in nonexpansive mappings by proposed the following algorithm:

$$\begin{cases} y_n = x_n + \gamma \delta A^*(S - I)Ax_n, \\ z_n = (1 - \beta_n)y_n + \beta_n Ty_n, \\ w_n = (1 - \gamma_n)y_n + \gamma_n Ry_n, \\ x_{n+1} = (1 - \alpha_n)z_n + \alpha_n U w_n, \quad n \geq 0, \end{cases} \tag{5}$$

The weak convergence result of the algorithm will be established. Our results improve and develop previously discussed feasibility problem and related algorithm.

On the other hand, we also suggest the following Halpern-type iterative algorithm to solve the SCFP (3). Define $\{x_n\}$ in the following iteration process:

$$\begin{cases} y_n = Tx_n + (1 - \beta_n)\delta A^*(S - I)ATx_n, \\ x_{n+1} = \alpha_n u + (1 - \alpha_n)y_n, \quad n \geq 0, \end{cases} \tag{6}$$

where u is an arbitrary fixed element in H_1 . We prove, under certain appropriate assumptions on the sequences $\{\alpha_n\}$ and $\{\beta_n\}$ that $\{x_n\}$ defined by (6) converges to a split common fixed point x^* of Γ_0 . Strong convergence theorem will be proven.

2. Preliminaries

In this section, we collect definitions and lemmas which will be used in the sequel.

Let H be a real Hilbert space with inner product $\langle \cdot, \cdot \rangle$ and norm $\| \cdot \|$, respectively. Assume that C be a nonempty and closed convex subset of H . Denote $F(T)$ the set of fixed points of a mapping $T : C \rightarrow C$, that is, $Fix(T) = \{x \in C : x = Tx\}$. We shall use the following notation: 1. $x_n \rightarrow x$ stands for the strong convergence of $\{x_n\}$ to x ; 2. $x_n \rightharpoonup x$ stands for the weak convergence of $\{x_n\}$ to x ; 3. $\omega_w(x_n) = \{x : \exists x_{n_i} \rightarrow x \text{ weakly}\}$ denote the weak w -limit set of $\{x_n\}$.

Definition 2.1. $T : C \rightarrow C$ is said to be nonexpansive mapping if

$$\|Tx - Ty\| \leq \|x - y\|, \quad \forall x, y \in C.$$

It is well known that in real Hilbert space H , the following statements hold

$$\|tx + (1 - t)y\|^2 = t\|x\|^2 + (1 - t)\|y\|^2 - t(1 - t)\|x - y\|^2, \tag{7}$$

for all $x, y \in H$ and $t \in [0, 1]$, and identical equation

$$\|x + y\|^2 = \|x\|^2 + 2\langle x, y \rangle + \|y\|^2, \tag{8}$$

for all $x, y \in H$. It follows that

$$\|x + y\|^2 \leq \|x\|^2 + 2\langle y, x + y \rangle. \tag{9}$$

for all $x, y \in H$.

Definition 2.2. A mapping T is said to be demiclosed if for any sequence $\{x_n\}$ which converges weakly to x , and if the sequence $\{Tx_n\}$ converges strongly to z , then $Tx = z$.

In what follows, only the particular case of demiclosedness at zero will be used, which is the particular case when $z = 0$. It is true that if T be a nonexpansive mapping, then $I - T$ is demiclosed.

Lemma 2.3 ([30]). Let H be a Hilbert space and $\{x_n\}$ be a sequence in H such that there exists a nonempty set $W \in H$ satisfying:

(1) for every $w \in W$, $\lim_{n \rightarrow \infty} \|x_n - w\|$ exists; (2) each weak-cluster point of the sequence $\{x_n\}$ is in W . Then there exists $w^* \in W$ such that $\{x_n\}$ weakly converges to w^* .

Lemma 2.4 ([31]). Let $\{a_n\}$ be a sequence of nonnegative real numbers satisfying the property

$$a_{n+1} \leq (1 - \gamma_n)a_n + \sigma_n, \quad n \geq 0,$$

where $\{\gamma_n\} \subset (0, 1)$ and $\{\sigma_n\}$ are such that

(1) $\sum_{n=0}^{\infty} \gamma_n = \infty$; (2) either $\limsup_{n \rightarrow \infty} \frac{\sigma_n}{\gamma_n} \leq 0$ or $\sum_{n=0}^{\infty} |\sigma_n| < \infty$.

Then $\{a_n\}$ converges to zero.

To attain strong convergence result, we need to use the following lemma.

Lemma 2.5 ([29]). Let $\{u_n\}$ be a sequence of real numbers. Assume $\{u_n\}$ does not decrease at infinity, that is, there exists at least a subsequence $\{u_{n_k}\}$ of $\{u_n\}$ such that $u_{n_k} \leq u_{n_k+1}$ for all $k \geq 0$. For every $n \geq N_0$, define an $\{\tau(n)\}$ as

$$\tau(n) = \max\{i \leq n : u_{n_i} < u_{n_i+1}\}.$$

Then $\tau(n) \rightarrow \infty$ as $n \rightarrow \infty$ and for all $n \geq N_0$,

$$\max\{u_{\tau(n)}, u_n\} \leq u_{\tau(n)+1}.$$

3. Weak convergence to split common fixed point of nonexpansive mappings

In this part, we will focus our attention on the following four-operator SCFP:

$$\text{Find } x^* \in \text{Fix}(T) \cap \text{Fix}(R) \cap \text{Fix}(U) \text{ such that } Ax^* \in \text{Fix}(S), \tag{10}$$

to solve (10), we suggested algorithm (5). We use Γ to denote the set of the solutions of (10).

Now, we are in a position to prove our convergence results.

Theorem 3.1. *Let H_1 and H_2 be two real Hilbert spaces. Let $A : H_1 \rightarrow H_2$ be a bounded linear operator. Let $S : H_2 \rightarrow H_2$ and $T, R, U : H_1 \rightarrow H_1$ are four nonexpansive mappings. Assume that $\Gamma \neq \emptyset$. Given $\{\alpha_n\}, \{\beta_n\}, \{\gamma_n\} \subset (0,1)$ and constants γ, δ satisfying the following conditions:*

- (i) $\liminf_{n \rightarrow \infty} \beta_n(1 - \alpha_n)(1 - \beta_n) > 0$; $\liminf_{n \rightarrow \infty} \alpha_n \gamma_n(1 - \gamma_n) > 0$; $\liminf_{n \rightarrow \infty} \alpha_n(1 - \alpha_n) > 0$;
- (ii) $0 < \gamma < 1$ and $0 < \delta < \frac{1}{\gamma\lambda}$, with λ being the spectral radius of the operator AA^* .

Define a sequence $\{x_n\}$ by the algorithm (5), then $\{x_n\}$ converges weakly to a $p \in \Gamma$.

Proof. First, we observe that $\{x_n\}$ is bounded, if we take an arbitrary fixed point x^* of Γ . Then we get $Ax^* \in \text{Fix}(S)$ and $x^* \in \text{Fix}(T) \cap \text{Fix}(R) \cap \text{Fix}(U)$, noting that S is nonexpansive mapping and using (8), we have

$$\begin{aligned} \|y_n - x^*\|^2 &= \|x_n - x^*\|^2 + 2\gamma\delta\langle Ax_n - Ax^*, (S - I)Ax_n \rangle \\ &\quad + \gamma^2\delta^2\langle AA^*(S - I)Ax_n, (S - I)Ax_n \rangle \\ &\leq \|x_n - x^*\|^2 + 2\gamma\delta\langle Ax_n - Ax^*, (S - I)Ax_n \rangle + \gamma^2\delta^2\lambda\|(S - I)Ax_n\|^2 \\ &= \|x_n - x^*\|^2 + \gamma^2\delta^2\lambda\|SAx_n - Ax_n\|^2 \\ &\quad + \gamma\delta(\|SAx_n - Ax^*\|^2 - \|Ax_n - Ax^*\|^2 - \|SAx_n - Ax_n\|^2) \\ &\leq \|x_n - x^*\|^2 - \gamma\delta(1 - \gamma\delta\lambda)\|SAx_n - Ax_n\|^2. \end{aligned} \tag{11}$$

From (5), (7) and (11), we obtain that

$$\begin{aligned} \|x_{n+1} - x^*\|^2 &= (1 - \alpha_n)\|z_n - x^*\|^2 + \alpha_n\|Uw_n - x^*\|^2 - \alpha_n(1 - \alpha_n)\|Uw_n - z_n\|^2 \\ &\leq (1 - \alpha_n)\|(1 - \beta_n)(y_n - x^*) + \beta_n(Ty_n - x^*)\|^2 \\ &\quad + \alpha_n\|(1 - \gamma_n)(y_n - x^*) + \gamma_n(Ry_n - x^*)\|^2 - \alpha_n(1 - \alpha_n)\|Uw_n - z_n\|^2 \\ &\leq \alpha_n(1 - \gamma_n)\|y_n - x^*\|^2 + \alpha_n\gamma_n\|y_n - x^*\|^2 - \alpha_n\gamma_n(1 - \gamma_n)\|Ry_n - y_n\|^2 \\ &\quad + (1 - \alpha_n)(1 - \beta_n)\|y_n - x^*\|^2 + (1 - \alpha_n)\beta_n\|y_n - x^*\|^2 \\ &\quad - \beta_n(1 - \alpha_n)(1 - \beta_n)\|Ty_n - y_n\|^2 - \alpha_n(1 - \alpha_n)\|Uw_n - z_n\|^2 \\ &= \|y_n - x^*\|^2 - \beta_n(1 - \alpha_n)(1 - \beta_n)\|Ty_n - y_n\|^2 - \alpha_n\gamma_n(1 - \gamma_n)\|Ry_n - y_n\|^2 \\ &\quad - \alpha_n(1 - \alpha_n)\|Uw_n - z_n\|^2 \\ &\leq \|x_n - x^*\|^2 - \gamma\delta(1 - \gamma\delta\lambda)\|SAx_n - Ax_n\|^2 - \beta_n(1 - \alpha_n)(1 - \beta_n)\|Ty_n - y_n\|^2 \\ &\quad - \alpha_n\gamma_n(1 - \gamma_n)\|Ry_n - y_n\|^2 - \alpha_n(1 - \alpha_n)\|Uw_n - z_n\|^2. \end{aligned} \tag{12}$$

We deduce immediately by conditions (i) and (ii) that

$$\|x_{n+1} - x^*\| \leq \|x_n - x^*\|,$$

for every $x^* \in \Gamma$, and for all $n \geq 0$. Thus, $\{x_n\}$ generated by algorithm (5) is the Féjer-monotone with respect to Γ . So, we obtain $\lim_{n \rightarrow \infty} \|x_n - x^*\|$ exists immediately, this implies that $\{x_n\}$ is bounded, the sequence $\{\|x_n - x^*\|\}$ is monotonically decreasing. From (12), we have

$$\lim_{n \rightarrow \infty} \|SAx_n - Ax_n\| = 0. \tag{13}$$

Since $\{x_n\}$ is bounded, $\omega_w(x_n) \neq \emptyset$. Denoting by p a weak-cluster point of $\{x_n\}$, we have $w - \lim_{i \rightarrow \infty} x_{n_i} = p$. Noting that demiclosedness of $I - S$ at 0, from (13), we obtain

$$SAp = Ap, \tag{14}$$

from which it follows that $Ap \in \text{Fix}(S)$. By setting $y_n = x_n + \gamma\delta A^*(S - I)Ax_n$, it follows that $w - \lim_{i \rightarrow \infty} y_{n_i} = p$. Again, from (12), we have

$$\lim_{n \rightarrow \infty} \|Ty_n - y_n\| = 0 \text{ and } \lim_{n \rightarrow \infty} \|Ry_n - y_n\| = 0, \tag{15}$$

by the demiclosedness of $I - T, I - R$ at 0 and the weak convergence of $\{y_{n_i}\}$ to p yields

$$Tp = p \text{ and } Rp = p. \tag{16}$$

Next, we prove that $\lim_{n \rightarrow \infty} \|Uw_n - w_n\| = 0$.

From (12), we deduce

$$\lim_{n \rightarrow \infty} \|Uw_n - z_n\| = 0. \tag{17}$$

By (15), we obtain that $\lim_{n \rightarrow \infty} \|z_n - y_n\| = 0$ and $\lim_{n \rightarrow \infty} \|w_n - y_n\| = 0$. Thus, we get

$$\lim_{n \rightarrow \infty} \|w_n - z_n\| = 0. \tag{18}$$

By (17) and (18), we get $\lim_{n \rightarrow \infty} \|Uw_n - w_n\| = 0$. The key limit above combined with $\lim_{n \rightarrow \infty} \|w_n - y_n\| = 0$, the demiclosedness of $I - U$ at 0 and the weak convergence of $\{y_{n_i}\}$ to p yield $Up = p$, combined with (14) and (16) we get $p \in \Gamma$. Since there is no more than one weak-cluster point, the weak convergence of the whole sequence $\{x_n\}$ follows by applying Lemma 2.3 with $W = \Gamma$. This completes the proof. \square

Chen et. al. [9], proposed a sequence $\{x_n\}$ generate by the following iterative method:

$$\begin{cases} y_n = \beta_n x_n + (1 - \beta_n)Tx_n, \\ z_n = \gamma_n x_n + (1 - \gamma_n)Sx_n, \\ x_{n+1} = \alpha_n y_n + (1 - \alpha_n)z_n, \quad n \geq 0, \end{cases}$$

where T, S are nonexpansive mappings, $\{\alpha_n\}, \{\beta_n\}$ are sequences in $(0,1)$.

Inspired by above algorithm, we suggest and analyze the corollary 3.2 as below. We denote $\Gamma_1 = \{x^* : x^* \in \text{Fix}(T) \cap \text{Fix}(R), Ax^* \in \text{Fix}(S)\}$.

By the careful analysis of the proof of Theorem 3.1, we can obtain the following result. Because its proof is much simpler than that of Theorem 3.1, we omit its proof.

Corollary 3.2. *Let H_1 and H_2 be two real Hilbert spaces. Let $A : H_1 \rightarrow H_2$ be a bounded linear operator. Let $S : H_2 \rightarrow H_2$ and $T, R : H_1 \rightarrow H_1$ are three nonexpansive mappings. Assume that $\Gamma_1 \neq \emptyset$. Given $\{\alpha_n\}, \{\beta_n\} \subset (0,1)$ and constants γ, δ satisfying the following conditions:*

- (1) $\liminf_{n \rightarrow \infty} \beta_n(1 - \alpha_n)(1 - \beta_n) > 0$; $\liminf_{n \rightarrow \infty} \alpha_n \gamma_n(1 - \gamma_n) > 0$;
- (2) $0 < \gamma < 1$ and $0 < \delta < \frac{1}{\gamma\lambda}$, with λ being the spectral radius of the operator AA^* .

Define a sequence $\{x_n\}$ by the following algorithm:

$$\begin{cases} y_n = x_n + \gamma\delta A^*(S - I)Ax_n, \\ z_n = (1 - \beta_n)y_n + \beta_nTy_n, \\ w_n = (1 - \gamma_n)y_n + \gamma_nRy_n, \\ x_{n+1} = (1 - \alpha_n)z_n + \alpha_nw_n, \quad n \geq 0, \end{cases}$$

then $\{x_n\}$ converges weakly to a split common fixed point $p \in \Gamma_1$.

Next, applying Theorem 3.1, we get the result as below.

Corollary 3.3. *Let H_1 and H_2 be two real Hilbert spaces. Let $A : H_1 \rightarrow H_2$ be a bounded linear operator. Let $S : H_2 \rightarrow H_2$ and $T, R : H_1 \rightarrow H_1$ are three nonexpansive mappings. Assume that $\Gamma_1 \neq \emptyset$. Given $\{\alpha_n\}, \{\beta_n\} \subset (0,1)$ and constants γ, δ satisfying the following conditions:*

- (1) $\liminf_{n \rightarrow \infty} \alpha_n \beta_n (1 - \beta_n) > 0; \liminf_{n \rightarrow \infty} \alpha_n (1 - \alpha_n) > 0;$
- (2) $0 < \gamma < 1$ and $0 < \delta < \frac{1}{\gamma \lambda}$, with λ being the spectral radius of the operator AA^* .

Define a sequence $\{x_n\}$ by the following algorithm:

$$\begin{cases} y_n = x_n + \gamma \delta A^*(S - I)Ax_n, \\ z_n = (1 - \beta_n)y_n + \beta_n T y_n, \\ x_{n+1} = (1 - \alpha_n)y_n + \alpha_n R z_n, \quad n \geq 0, \end{cases}$$

then $\{x_n\}$ converges weakly to a split common fixed point $p \in \Gamma_1$.

Now, we suggest a new split common fixed point problem, that is, finding a point x^* with the property

$$x^* \in \text{Fix}(R) \text{ and } Ax^* \in \text{Fix}(T) \cap \text{Fix}(S). \tag{19}$$

We use Γ_2 to denote the set of the solutions of (19).

Theorem 3.4. *Let H_1 and H_2 be two real Hilbert spaces. Let $A : H_1 \rightarrow H_2$ be a bounded linear operator. Let $S, T : H_2 \rightarrow H_2$ and $R : H_1 \rightarrow H_1$ are three nonexpansive mappings. Assume that $\Gamma_2 \neq \emptyset$. The following conditions hold for the sequence $\{\alpha_n\} \subset (0,1)$ and constants γ, δ :*

- (i) $\liminf_{n \rightarrow \infty} \alpha_n (1 - \alpha_n) > 0;$
- (ii) $0 < \gamma, \delta < \frac{1}{\lambda}$, with λ being the spectral radius of the operator AA^* .

Then a sequence $\{x_n\}$ by the following algorithm:

$$\begin{cases} y_n = x_n + \gamma A^*(T - I)Ax_n, \\ z_n = x_n + \delta A^*(S - I)Ax_n, \\ x_{n+1} = (1 - \alpha_n)y_n + \alpha_n R z_n, \quad n \geq 0, \end{cases} \tag{20}$$

converges weakly to a split common fixed point $p \in \Gamma_2$.

Proof. As proved in Theorem 3.1, only a sketch of the proof is given here.

First, we observe that $\{x_n\}$ is bounded, taking x^* of Γ_2 , i.e., $Ax^* \in \text{Fix}(T) \cap \text{Fix}(S)$ and $x^* \in \text{Fix}(R)$, noting that T is nonexpansive mapping and (8), using the same argument in Theorem 3.1, we have

$$\|y_n - x^*\|^2 \leq \|x_n - x^*\|^2 - \gamma(1 - \gamma\lambda)\|TAx_n - Ax_n\|^2. \tag{21}$$

Similarly, we get

$$\|z_n - x^*\|^2 \leq \|x_n - x^*\|^2 - \delta(1 - \delta\lambda)\|SAx_n - Ax_n\|^2. \tag{22}$$

From (20), (7), (21) and (22), we obtain that

$$\begin{aligned} \|x_{n+1} - x^*\|^2 &\leq (1 - \alpha_n)\|y_n - x^*\|^2 + \alpha_n\|z_n - x^*\|^2 - \alpha_n(1 - \alpha_n)\|Rz_n - y_n\|^2 \\ &\leq (1 - \alpha_n)(\|x_n - x^*\|^2 - \gamma(1 - \gamma\lambda)\|TAx_n - Ax_n\|^2) \\ &\quad + \alpha_n(\|x_n - x^*\|^2 - \delta(1 - \delta\lambda)\|SAx_n - Ax_n\|^2) - \alpha_n(1 - \alpha_n)\|Rz_n - y_n\|^2 \\ &\leq \|x_n - x^*\|^2 - (1 - \alpha_n)\gamma(1 - \gamma\lambda)\|TAx_n - Ax_n\|^2 - \alpha_n(1 - \alpha_n)\|Rz_n - y_n\|^2 \\ &\quad - \alpha_n\delta(1 - \delta\lambda)\|SAx_n - Ax_n\|^2. \end{aligned} \tag{23}$$

We deduce immediately by conditions (i) and (ii) that $\|x_{n+1} - x^*\| \leq \|x_n - x^*\|$, for every $x^* \in \Gamma_2$, and for all $n \in \mathbb{N}$. Thus, $\{x_n\}$ generated by algorithm (3.11) is the Féjer-monotone with respect to Γ_2 . So, we

obtain $\lim_{n \rightarrow \infty} \|x_n - x^*\|$ exists immediately, this implies that $\{x_n\}$ is bounded, the sequence $\{\|x_n - x^*\|\}$ is monotonically decreasing.

From (23), we have

$$\lim_{n \rightarrow \infty} \|SAx_n - Ax_n\| = 0 \text{ and } \lim_{n \rightarrow \infty} \|TAx_n - Ax_n\| = 0.$$

On the lines similar to Theorem 3.1, we get $p \in \Gamma$. Since there is no more than one weak-cluster point, the weak convergence of the whole sequence $\{x_n\}$ follows by applying Lemma 2.3 with $W = \Gamma_2$. This completes the proof. \square

We suggest an algorithm in the following Corollary to solve (3).

Corollary 3.5. *Let H_1 and H_2 be two real Hilbert spaces. Let $A : H_1 \rightarrow H_2$ be a bounded linear operator. Let $S : H_2 \rightarrow H_2$ and $T : H_1 \rightarrow H_1$ are two nonexpansive mappings. Assume that $\Gamma_0 \neq \emptyset$. The following conditions hold for the sequences $\{\alpha_n\} \subset (0,1)$ and constant γ :*

- (1) $\liminf_{n \rightarrow \infty} \alpha_n(1 - \alpha_n) > 0$;
- (2) $0 < \gamma < \frac{1}{\lambda}$, with λ being the spectral radius of the operator AA^* .

Then a sequence $\{x_n\}$ by the following algorithm:

$$\begin{cases} y_n = x_n + \gamma A^*(S - I)Ax_n, \\ x_{n+1} = (1 - \alpha_n)x_n + \alpha_n T y_n, \quad n \geq 0, \end{cases}$$

converges weakly to a split common fixed point $p \in \Gamma_0$.

Finally, we propose the following split common fixed point problem, that is, finding a point x^* with the property

$$x^* \in H_1 \text{ and } Ax^* \in \text{Fix}(T) \cap \text{Fix}(S). \tag{24}$$

We use Γ_3 to denote the set of the solutions of (24).

We introduce the following algorithm, which is take $R = I$ in the algorithm (20) in Theorem 3.4.

Corollary 3.6. *Let H_1 and H_2 be two real Hilbert spaces. Let $A : H_1 \rightarrow H_2$ be a bounded linear operator. Let $T, S : H_2 \rightarrow H_2$ are two nonexpansive mappings. Assume that $\Gamma_3 \neq \emptyset$. Given $\{\alpha_n\}$ be sequence in $(0,1)$ and constants γ, δ satisfying $0 < \gamma, \delta < \frac{1}{\lambda}$, with λ being the spectral radius of the operator AA^* .*

Then a sequence $\{x_n\}$ by the following algorithm:

$$\begin{cases} y_n = x_n + \gamma A^*(T - I)Ax_n, \\ z_n = x_n + \delta A^*(S - I)Ax_n, \\ x_{n+1} = (1 - \alpha_n)y_n + \alpha_n z_n, \quad n \geq 0, \end{cases}$$

converges weakly to a split common fixed point $p \in \Gamma_3$.

Proof. Take $x^* \in \Gamma_3$, i.e., $Ax^* \in \text{Fix}(T) \cap \text{Fix}(S)$, using the same argument in Theorem 3.4, we have the bounded of $\{x_n\}$, $\omega_w(x_n) \neq \emptyset$, such that $w - \lim_{i \rightarrow \infty} x_{n_i} = p$. By setting $\{y_n\}, \{z_n\}$ and combined with the demiclosedness of $I - T, I - S$ at 0, all the conditions in this corollary are satisfied, the conclusion of Corollary 3.6 can be obtained from Theorem 3.4 immediately. \square

We denote $\Gamma_4 = \{x^* : x^* \in H_1, Ax^* \in \text{Fix}(S)\}$.

Corollary 3.7. *Let H_1 and H_2 be two real Hilbert spaces. Let $A : H_1 \rightarrow H_2$ be a bounded linear operator. Let $S : H_2 \rightarrow H_2$ be a nonexpansive mapping. Assume that $\Gamma_4 \neq \emptyset$. Given $\{\alpha_n\}$ be sequence in $(0,1)$ and constants γ , satisfying $0 < \gamma < \frac{1}{\lambda}$, with λ being the spectral radius of the operator AA^* .*

Then a sequence $\{x_n\}$ by the following algorithm:

$$\begin{cases} y_n = x_n + \gamma A^*(S - I)Ax_n, \\ x_{n+1} = (1 - \alpha_n)x_n + \alpha_n y_n, \quad n \geq 0, \end{cases}$$

converges weakly to a split common fixed point $p \in \Gamma_4$.

4. Strong convergence to split common fixed point of nonexpansive mappings

In this section, we suggest a Halpern-type iterative algorithm to solve the SCFP (3) involved in nonexpansive mappings, and prove its strong convergence.

Theorem 4.1. *Let H_1 and H_2 be two real Hilbert spaces. Let $A : H_1 \rightarrow H_2$ be a bounded linear operator. Let $S : H_2 \rightarrow H_2$ and $T : H_1 \rightarrow H_1$ are two nonexpansive mappings. Assume that $\Gamma_0 \neq \emptyset$. Given $\{\alpha_n\}, \{\beta_n\} \subset (0,1)$ and constant δ satisfying the following conditions:*

- (i) $\lim_{n \rightarrow \infty} \alpha_n = 0, \sum_{n=0}^{\infty} \alpha_n = \infty;$
- (ii) $\sum_{n=0}^{\infty} |\alpha_n - \alpha_{n-1}| < \infty, \sum_{n=0}^{\infty} |\beta_n - \beta_{n-1}| < \infty,$
- (iii) $0 < \beta_n < b < 1, 0 < \delta < \frac{1-b}{\lambda},$ with λ being the spectral radius of the operator AA^* .

For fixed $u, x_0 \in H_1$ arbitrarily, define a sequence $\{x_n\}$ by algorithm (6), then $\{x_n\}$ converges strongly to x^* , given by $x^* = P_{\Gamma_0}(u)$.

Proof. First, we observe that $\{x_n\}$ is bounded, if we take an arbitrary fixed point x^* of Γ_0 . Then we get $Ax^* \in \text{Fix}(S)$ and $x^* \in \text{Fix}(T)$, noting that S is nonexpansive mapping, using (8) and condition (iii), we have

$$\begin{aligned} \|y_n - x^*\|^2 &\leq \|Tx_n - x^*\|^2 + 2(1 - \beta_n)\delta \langle Tx_n - Tx^*, A^*(S - I)ATx_n \rangle + \delta^2 \|A^*(S - I)ATx_n\|^2 \\ &\leq \|x_n - x^*\|^2 + 2(1 - \beta_n)\delta \langle ATx_n - ATx^*, (S - I)ATx_n \rangle + \lambda\delta^2 \|(S - I)ATx_n\|^2 \\ &= \|x_n - x^*\|^2 + \lambda\delta^2 \|SATx_n - ATx_n\|^2 \\ &\quad + (1 - \beta_n)\delta (\|SATx_n - ATx^*\|^2 - \|ATx_n - ATx^*\|^2 - \|SATx_n - ATx_n\|^2) \\ &= \|x_n - x^*\|^2 - \delta((1 - \beta_n) - \lambda\delta) \|SATx_n - ATx_n\|^2 \\ &\leq \|x_n - x^*\|^2. \end{aligned} \tag{25}$$

From (6) and (25), we obtain that

$$\begin{aligned} \|x_{n+1} - x^*\| &\leq \alpha_n \|u - x^*\| + (1 - \alpha_n) \|y_n - x^*\| \\ &\leq \alpha_n \|u - x^*\| + (1 - \alpha_n) \|x_n - x^*\| \\ &\leq \max\{\|u - x^*\|, \|x_n - x^*\|\}. \end{aligned}$$

The boundedness of the sequence $\{x_n\}$ yields by the induction. So are $\{y_n\}$ and $\{Tx_n\}$. These boundedness play a prominent role in proving the strong convergence theorem.

From (6), (7) and (25), we have

$$\begin{aligned} \|x_{n+1} - x^*\|^2 &\leq \alpha_n \|u - x^*\|^2 + (1 - \alpha_n) \|y_n - x^*\|^2 \\ &\leq \alpha_n \|u - x^*\|^2 + \|x_n - x^*\|^2 - \delta((1 - b) - \lambda\delta) \|SATx_n - ATx_n\|^2. \end{aligned}$$

It follows that

$$\delta(1 - b - \lambda\delta) \|SATx_n - ATx_n\|^2 \leq \alpha_n \|u - x^*\|^2 + \|x_n - x^*\|^2 - \|x_{n+1} - x^*\|^2. \tag{26}$$

Next, we consider two possible cases.

Case 1. Assume there exists some integer $m > 0$ such that $\{\|x_n - x^*\|\}$ is decreasing for all $n > m$, in this case, we know that $\lim_{n \rightarrow \infty} \|x_n - x^*\|$ exists. From (26) and condition (i), (iii), we deduce

$$\lim_{n \rightarrow \infty} \|SATx_n - ATx_n\| = 0. \tag{27}$$

As a result, by setting $y_n = Tx_n + (1 - \beta_n)\delta A^*(S - I)ATx_n$, we obtain that

$$\lim_{n \rightarrow \infty} \|y_n - Tx_n\| = 0. \tag{28}$$

Since $\{y_n\}$ is bounded, from (6) and condition (i), we have

$$\lim_{n \rightarrow \infty} \|x_{n+1} - y_n\| = 0. \tag{29}$$

We next show that

$$\lim_{n \rightarrow \infty} \|x_n - Tx_n\| = 0. \tag{30}$$

As a matter of fact $\|x_n - Tx_n\| \leq \|x_n - x_{n+1}\| + \|x_{n+1} - y_n\| + \|y_n - Tx_n\|$, so, it suffices to show that

$$\lim_{n \rightarrow \infty} \|x_n - x_{n+1}\| = 0, \tag{31}$$

if (31) holds, then (30) holds from (28) and (29).

Before calculate $x_n - x_{n+1}$, letting $z_n = Tx_n + \delta A^*(S - I)ATx_n$, on the lines similar to (11), we get the boundedness of the sequence $\{z_n\}$.

For every integer $n \geq 1$, we have

$$\begin{aligned} \|z_n - z_{n-1}\|^2 &= \|Tx_n - Tx_{n-1}\|^2 + \delta^2 \|A^*(S - I)ATx_n - A^*(S - I)ATx_{n-1}\|^2 \\ &\quad + 2\delta \langle Tx_n - Tx_{n-1}, A^*(S - I)ATx_n - A^*(S - I)ATx_{n-1} \rangle \\ &\leq \|x_n - x_{n-1}\|^2 + \lambda \delta^2 \|(S - I)ATx_n - (S - I)ATx_{n-1}\|^2 \\ &\quad + 2\delta \langle ATx_n - ATx_{n-1}, (S - I)ATx_n - (S - I)ATx_{n-1} \rangle \\ &= \|x_n - x_{n-1}\|^2 + \lambda \delta^2 \|(S - I)ATx_n - (S - I)ATx_{n-1}\|^2 \\ &\quad + \delta (\|SATx_n - SATx_{n-1}\|^2 - \|ATx_n - ATx_{n-1}\|^2 - \|(S - I)ATx_n - (S - I)ATx_{n-1}\|^2) \\ &\leq \|x_n - x_{n-1}\|^2 - \delta(1 - \lambda\delta) \|(S - I)ATx_n - (S - I)ATx_{n-1}\|^2 \\ &\leq \|x_n - x_{n-1}\|^2. \end{aligned} \tag{32}$$

In order to prove (30), we now calculate $x_{n+1} - x_n$ for every integer $n \geq 1$,

$$\begin{aligned} x_{n+1} - x_n &= (\alpha_n u + (1 - \alpha_n)((1 - \beta_n)z_n + \beta_n Tx_n)) - (\alpha_{n-1} u + (1 - \alpha_{n-1})((1 - \beta_{n-1})z_{n-1} + \beta_{n-1} Tx_{n-1})) \\ &= (\alpha_n - \alpha_{n-1})u + (1 - \alpha_n)(1 - \beta_n)(z_n - z_{n-1}) + (1 - \alpha_n)(1 - \beta_n)z_{n-1} - (1 - \alpha_{n-1})\beta_{n-1}Tx_{n-1} \\ &\quad + (1 - \alpha_n)\beta_n(Tx_n - Tx_{n-1}) + (1 - \alpha_n)\beta_n Tx_{n-1} - (1 - \alpha_{n-1})(1 - \beta_{n-1})z_{n-1} \\ &= (\alpha_n - \alpha_{n-1})u + (1 - \alpha_n)(1 - \beta_n)(z_n - z_{n-1}) + (1 - \alpha_n)(1 - \beta_n)Tx_{n-1} \\ &\quad + (1 - \alpha_n)(1 - \beta_n)(z_{n-1} - Tx_{n-1}) - (1 - \alpha_{n-1})(1 - \beta_{n-1})(z_{n-1} - Tx_{n-1}) + (1 - \alpha_n)\beta_n Tx_{n-1} \\ &\quad - (1 - \alpha_{n-1})(1 - \beta_{n-1})Tx_{n-1} + (1 - \alpha_n)\beta_n(Tx_n - Tx_{n-1}) - (1 - \alpha_{n-1})\beta_{n-1}Tx_{n-1} \\ &= (\alpha_n - \alpha_{n-1})u + (1 - \alpha_n)Tx_{n-1} - (1 - \alpha_{n-1})Tx_{n-1} + (1 - \alpha_n)(1 - \beta_n)(z_n - z_{n-1}) \\ &\quad + ((1 - \alpha_n)(1 - \beta_n) - (1 - \alpha_{n-1})(1 - \beta_{n-1}))(z_{n-1} - Tx_{n-1}) + (1 - \alpha_n)\beta_n(Tx_n - Tx_{n-1}) \\ &= (\alpha_n - \alpha_{n-1})(u - Tx_{n-1}) + (1 - \alpha_n)(1 - \beta_n)(z_n - z_{n-1}) \\ &\quad + ((\beta_{n-1} - \beta_n)(1 - \alpha_n) + (\alpha_{n-1} - \alpha_n)(1 - \beta_{n-1}))(z_{n-1} - Tx_{n-1}) + (1 - \alpha_n)\beta_n(Tx_n - Tx_{n-1}). \end{aligned}$$

It follows from (32) that

$$\begin{aligned} \|x_{n+1} - x_n\| &\leq (1 - \alpha_n)\beta_n \|Tx_n - Tx_{n-1}\| + (1 - \alpha_n)(1 - \beta_n) \|z_n - z_{n-1}\| \\ &\quad + (|\beta_n - \beta_{n-1}| + |\alpha_n - \alpha_{n-1}|) \|z_{n-1} - Tx_{n-1}\| + |\alpha_n - \alpha_{n-1}| \|u - Tx_{n-1}\| \\ &\leq (1 - \alpha_n) \|x_n - x_{n-1}\| + M(|\beta_n - \beta_{n-1}| + 2|\alpha_n - \alpha_{n-1}|), \end{aligned} \tag{33}$$

where M is a constant such that $M = \max\{\|z_{n-1} - Tx_{n-1}\|, \|u - Tx_{n-1}\|\}$ for all $n \geq 1$. By assumptions (i) and (ii), we have $\sum_{n=0}^{\infty} \alpha_n = \infty$ and $\sum_{n=0}^{\infty} (|\beta_n - \beta_{n-1}| + 2|\alpha_n - \alpha_{n-1}|) < \infty$. Hence, Lemma 2.4 is applicable to (33) and we obtain $\lim_{n \rightarrow \infty} \|x_n - x_{n+1}\| = 0$.

Since the sequence $\{x_n\}$ is bounded, we can choose a subsequence $\{x_{n_i}\}$ of $\{x_n\}$ such that $x_{n_i} \rightarrow x$. Consequently, we derive from (27) and (30) that

$$Tx_{n_i} \rightarrow x, ATx_{n_i} \rightarrow Ax.$$

Applying demiclosedness principle, we deduce

$$x \in \text{Fix}(T) \text{ and } Ax \in \text{Fix}(S).$$

That is to say, $x \in \Gamma_0$.

Therefore,

$$\limsup_{n \rightarrow \infty} \langle u - x^*, x_{n+1} - x^* \rangle = \lim_{i \rightarrow \infty} \langle u - x^*, x_{n_i} - x^* \rangle = \langle u - x^*, x - x^* \rangle \leq 0. \tag{34}$$

Using (9), we have

$$\begin{aligned} \|x_{n+1} - x^*\|^2 &\leq (1 - \alpha_n) \|y_n - x^*\|^2 + 2\alpha_n \langle u - x^*, x_{n+1} - x^* \rangle \\ &\leq (1 - \alpha_n) \|x_n - x^*\|^2 + 2\alpha_n \langle u - x^*, x_{n+1} - x^* \rangle. \end{aligned} \tag{35}$$

Applying Lemma 2.4 and (34) and (35), we deduce $x_n \rightarrow x^*$, i.e., sequence $\{x_n\}$ converges strongly to a split common fixed point x^* , given by $x^* = P_{\Gamma_0}(u)$.

Case 2. Assume there exists an integers n_0 , such that

$$\|x_{n_0} - x^*\| \leq \|x_{n_0+1} - x^*\|.$$

Set $u_n = \{\|x_n - x^*\|\}$, then we have

$$u_{n_0} \leq u_{n_0+1}.$$

Define an integer sequence $\{\tau_n\}$ for all $n \geq n_0$ as follows:

$$\tau(n) = \max\{l \geq 1 : n_0 \leq l \leq n, u_l \leq u_{l+1}\}.$$

It is clear that $\tau(n)$ is non-decreasing sequence satisfying

$$\lim_{n \rightarrow \infty} \tau(n) = \infty,$$

and

$$u_{\tau(n)} \leq u_{\tau(n)+1},$$

for all $n \geq n_0$.

By a similar argument to that of Case 1, we can obtain that

$$\lim_{n \rightarrow \infty} \|SATx_{\tau(n)} - ATx_{\tau(n)}\| = 0 \quad \text{and} \quad \lim_{n \rightarrow \infty} \|x_{\tau(n)} - Tx_{\tau(n)}\| = 0.$$

This implies that

$$\omega(x_{\tau(n)}) \subset \Gamma_0.$$

Thus, we obtain

$$\limsup_{n \rightarrow \infty} \langle u - x^*, x_{\tau(n)+1} - x^* \rangle \leq 0. \tag{36}$$

Since $u_{\tau(n)} \leq u_{\tau(n)+1}$, we have from (35) that

$$u_{\tau(n)}^2 \leq u_{\tau(n)+1}^2 \leq (1 - \alpha_{\tau(n)})u_{\tau(n)}^2 + 2\alpha_{\tau(n)} \langle u - x^*, x_{\tau(n)+1} - x^* \rangle. \tag{37}$$

It follows that

$$u_{\tau(n)}^2 \leq 2 \langle u - x^*, x_{\tau(n)+1} - x^* \rangle. \tag{38}$$

Combining (36) and (38), we have $\limsup_{n \rightarrow \infty} u_{\tau(n)} \leq 0$, and hence

$$\lim_{n \rightarrow \infty} u_{\tau(n)} = 0. \tag{39}$$

By (37), we obtain

$$\limsup_{n \rightarrow \infty} u_{\tau(n)+1}^2 \leq \limsup_{n \rightarrow \infty} u_{\tau(n)}^2.$$

This together with (39) implies that

$$\lim_{n \rightarrow \infty} u_{\tau(n)+1} = 0.$$

Applying Lemma 2.5 to get

$$0 \leq u_n \leq \max\{u_{\tau(n)}, u_{\tau(n)+1}\}.$$

therefore, $u_n \rightarrow 0$. That is, $x_n \rightarrow x^*$, i.e., sequence $\{x_n\}$ converges strongly to a split common fixed point x^* , given by $x^* = P_{\Gamma_0}(u)$. This completes the proof. \square

If taking $u = 0$ in the algorithm in Theorem 4.1, we get the following corollary immediately.

Corollary 4.2. *Let H_1 and H_2 be two real Hilbert spaces. Let $A : H_1 \rightarrow H_2$ be a bounded linear operator. Let $S : H_2 \rightarrow H_2$ and $T : H_1 \rightarrow H_1$ are two nonexpansive mappings. Assume that $\Gamma_0 \neq \emptyset$. Given $\{\alpha_n\}, \{\beta_n\} \subset (0,1)$ and constant δ satisfying the following conditions:*

- (1) $\lim_{n \rightarrow \infty} \alpha_n = 0, \sum_{n=0}^{\infty} \alpha_n = \infty$;
 - (2) $\sum_{n=0}^{\infty} |\alpha_n - \alpha_{n-1}| < \infty, \sum_{n=0}^{\infty} |\beta_n - \beta_{n-1}| < \infty$,
 - (3) $0 < \beta_n < b < 1, 0 < \delta < \frac{1-b}{\lambda}$, with λ being the spectral radius of the operator AA^* .
- For fixed $x_0 \in H_1$ arbitrarily, define a sequence $\{x_n\}$ by the following algorithm,

$$\begin{cases} y_n = Tx_n + (1 - \beta_n)\delta A^*(S - I)ATx_n, \\ x_{n+1} = (1 - \alpha_n)y_n, \quad n \geq 0, \end{cases}$$

then $\{x_n\}$ converges strongly to a x^* , given by $x^* = P_{\Gamma_0}(0)$, which is the minimum norm in Γ_0 .

5. Acknowledgment

This work was supported in part by Applied Basic Research Project of Sichuan Province (2018JY0169, Chengdu Univ. Info. Tech.).

References

- [1] H. Bauschke and J. Borwein, *On projection algorithms for solving convex feasibility problems*, SIAM Rev., **38** (1996), 367–426.
- [2] C. Byrne, *Iterative oblique projection onto convex subsets and the split feasibility problem*, Inverse Probl., **18** (2002), 441–453.
- [3] C. Byrne, *A unified treatment of some iterative algorithm in signal processing and image reconstruction*, Inverse Probl., **20** (2004), 103–120.
- [4] L. C. Ceng, A. Petrusel, J. C. Yao and Y. Yao, *Hybrid viscosity extragradient method for systems of variational inequalities, fixed Points of nonexpansive mappings, zero points of accretive operators in Banach spaces*, Fixed Point Theory, **19** (2018), 487–502.
- [5] L. C. Ceng, A. Petrusel, J. C. Yao and Y. Yao, *Systems of variational inequalities with hierarchical variational inequality constraints for Lipschitzian pseudocontractions*, Fixed Point Theory, **20** (2019), 113–134.
- [6] Y. Censor, T. Bortfeld, B. Martin and A. Trofimov, *A unified approach for inversion problems in intensity modulated radiation therapy*, Phys. Med. Biol., **51** (2006), 2353–2365.
- [7] Y. Censor and T. Elfving, *A multiprojection algorithm using Bregman projections in a product space*, Numer. Algor., **8** (1994), 221–239.
- [8] Y. Censor and A. Segal, *The split common fixed point problem for directed operators*, J. Convex. Anal., **16** (2009), 587–600.
- [9] J. Chen and D. Wu, *A new iterative scheme for approximating common fixed points of two nonexpansive mappings in Banach space*, J. Appl. Math. Bio., **3** (2013), 285–300.
- [10] J. Chen, D. Wu and C. Zhang, *A new iterative scheme of modified Mann iteration in Banach space*, Abstr. Appl. Anal., **2014** (2014), Art. ID. 264909.

- [11] Y. Dang and Y. Gao, *The strong convergence of a KM-CQ-like algorithm for a split feasibility problem*, Inverse Probl., **27** (2011), Art. ID 015007.
- [12] A. Moudafi, *A note on the split common fixed-point problem for quasi-nonexpansive operators*, Nonlinear. Anal., **74** (2011), 4083–4087.
- [13] A. Moudafi and E. Al-Shemas, *Simultaneous iterative methods for split equality problem*, Trans. Math. Program. Appl., **1** (2013), 1–11.
- [14] B. Qu and N. Xiu, *A note on the CQ algorithm for the split feasibility problem*, Inverse Probl., **21** (2005), 1655–1665.
- [15] M. Sezan and H. Stark, *Applications of convex projection theory to image recovery in tomography and related areas*, in Image Recovery Theory and Applications. H.Stark, Ed. pp.415-462, Academic Press, Orlando, Fla, USA, 1987.
- [16] B. S. Thakur, D. Thakur and M. Postolache, *A new iterative scheme for numerical reckoning fixed points of Suzuki's generalized nonexpansive mappings*, Appl. Math. Comput., **275** (2016), 147–155.
- [17] F. Wang and H. K. Xu, *Approximating curve and strong convergence of the CQ algorithm for the split feasibility problem*, J. Inequal. Appl., **2010** (2010), Article ID 102085.
- [18] F. Wang and H. K. Xu, *Cyclic algorithms for the split feasibility problems in Hilbert spaces*, Nonlinear. Anal., **74** (2011), 4105–4111.
- [19] Z. Wang, Q. Yang and Y. Yang, *The relaxed inexact projection methods for the split feasibility problem*, Appl. Math. Comput., **217** (2011), 5347–5359.
- [20] H. K. Xu, *A variable Krasnosel'skii-Mann algorithm and the multiple-set split feasibility problem*, Inverse Probl., **22** (2006), 2021–2034.
- [21] H. K. Xu, *Iterative methods for the split feasibility problem in infinite-dimensional Hilbert spaces*, Inverse Probl., **26** (2010), Art. ID 105018.
- [22] Q. Yang, *The relaxed CQ algorithm solving the split feasibility problem*, Inverse Probl., **20** (2004), 1261–1266.
- [23] Y. Yao, R. Agarwal, M. Postolache and Y. Liou, *Algorithms with strong convergence for the split common solution of the feasibility problem and fixed point problem*, Fixed Point Theory Appl., **2014** (2014), Article ID 183.
- [24] Y. Yao, L. Leng, M. Postolache and X. Zheng, *Mann-type iteration method for solving the split common fixed point problem*, J. Nonlinear Convex Anal., **18** (2017), 875–882.
- [25] Y. H. Yao, Y. C. Liou and J. C. Yao, *Split common fixed point problem for two quasi-pseudocontractive operators and its algorithm construction*, Fixed Point Theory Appl., **2015** (2015), Art. No. 127.
- [26] Y. Yao, Y. C. Liou and J. C. Yao, *Iterative algorithms for the split variational inequality and fixed point problems under nonlinear transformations*, J. Nonlinear Sci. Appl., **10** (2017), 843–854.
- [27] Y. Yao, M. Postolache and Y. C. Liou, *Strong convergence of a self-adaptive method for the split feasibility problem*, Fixed Point Theory Appl., **2013** (2013), Art. ID. 201.
- [28] Y. Yao, M. Postolache and J. C. Yao, *An iterative algorithm for solving the generalized variational inequalities and fixed points problems*, Mathematics, **7** (2019), Art. ID. 61.
- [29] Y. Yao, M. Postolache and Z. Zhu, *Gradient methods with selection technique for the multiple-sets split feasibility problem*, Optim., **69** (2020), 269-281.
- [30] Y. Yao, J. Wu and Y.C. Liou, *Regularized methods for the split feasibility problem*, Abstr. Appl. Anal., **2012** (2012), Art. ID. 140679.
- [31] Y. Yao, J. C. Yao, Y. C. Liou and M. Postolache, *Iterative algorithms for split common fixed points of demicontractive operators without priori knowledge of operator norms*, Carpathian J. Math., **34** (2018), 459–466.
- [32] J. Zhao and Q. Yang, *Seveal solution methods for the split feasibility problem*, Inverse Probl., **21** (2005), 1791–1799.
- [33] X. P. Zhao, J. C. Yao and Y. Yao, *A proximal algorithm for solving split monotone variational inclusions*, U.P.B. Sci. Bull., Series A, **82**(3) (2020), 43-52.
- [34] X. P. Zhao and Y. H. Yao, *Modified extragradient algorithms for solving monotone variational inequalities and fixed point problems*, Optim., in press.