



Complete Convergence and Complete Moment Convergence for Negatively Dependent Random Variables Under Sub-Linear Expectations

Qunying Wu^a, Yuanying Jiang^a

^aCollege of Science, Guilin University of Technology

Abstract. This paper we study and establish the complete convergence and complete moment convergence theorems under a sub-linear expectation space. As applications, the complete convergence and complete moment convergence for negatively dependent random variables with $C_V(\exp(\ln^\alpha |X|)) < \infty$, $\alpha > 1$ have been generalized to the sub-linear expectation space context. We extend some complete convergence and complete moment convergence theorems for the traditional probability space to the sub-linear expectation space. Our results generalize corresponding results obtained by Gut and Stadtmüller (2011), Qiu and Chen (2014) and Wu and Jiang (2016). There is no report on the complete moment convergence under sub-linear expectation, and we provide the method to study this subject.

1. Introduction

The sub-linear expectation space have advantages of modelling the uncertainty of probability and distribution. Therefore, limit theorems for sub-linear expectations have raised a large number of issues of interest recently. Limit theorems are important research topics in probability and statistics. They were widely used in finance and other fields. Classical limit theorems only hold in the case of model certainty. However, in practice, such model certainty assumption is not realistic in many areas of applications because the uncertainty phenomena cannot be modeled using model certainty. Motivated by modeling uncertainty in practice, Peng (2006 [14]) introduced a new notion of sub-linear expectation. As an alternative to the traditional probability/expectation, capacity/sub-linear expectation has been studied in many fields such as statistics, finance, economics, and measures of risk (see Denis and Martini (2006 [5]), Gilboa (1987 [6]), Marinacci (1999 [11]), Peng (1997 [12], 1999 [13], 2006 [14], 2008 [15]) etc). The general framework of the sub-linear expectation in a general function space was introduced by Peng (2006 [14], 2008 [15], 2009 [16]), and sub-linear expectation is a natural extension of the classical linear expectation.

Because the sub-linear expectation provides a very flexible framework to model sub-linear probability problems, the limit theorems of the sub-linear expectation have received more and more attention and research recently. A series of useful results have been established. Peng (2006 [14], 2008 [15], 2009

2010 *Mathematics Subject Classification.* Primary 60F15

Keywords. sub-linear expectation; complete convergence; complete moment convergence; negatively dependent random variables.

Received: 07 January 2019; Accepted: 19 January 2020

Communicated by Miljana Jovanović

Research supported by the National Natural Science Foundation of China (12061028, 71963008), and the Support Program of the Guangxi China Science Foundation (2018GXNSFAA281011, 2018GXNSFAA294131).

Email addresses: wqy666@glut.edu.cn (Qunying Wu), jyy@glut.edu.cn (Yuanying Jiang)

[16]) constructed the basic framework, basic properties and the central limit theorem under sub-linear expectations, Zhang (2016a [26], 2016b [27], 2016c [28]) established the exponential inequalities, Rosenthal's inequalities, strong law of larger numbers and law of iterated logarithm, Cheng (2016 [3]), Chen (2016 [2]) and Wu and Jiang (2017 [24]) also studied strong law of larger numbers, and so on. In general, extending the limit properties of conventional probability space to the cases of sub-linear expectation is highly desirable and of considerable significance in the theory and application. Because sub-linear expectation and capacity are not additive, many powerful tools and common methods for linear expectations and probabilities are no longer valid, so that the study of the limit theorems under sub-linear expectation becomes much more complex and difficult.

Complete convergence and complete moment convergence are the most important problems in probability theory. Many of their related results have been obtained in the probabilistic space. However, the complete moment convergence under sub-linear expectation has not been reported. In this paper, we establish the complete convergence and complete moment convergence for negatively dependent random variables under sub-linear expectation. As a result, the corresponding results obtained by Gut and Stadtmüller (2011 [8]), Qiu and Chen (2014 [17]) and Wu and Jiang (2016 [23]) have been generalized to the sub-linear expectation space context.

In the next section, we summarize some basic notations and concepts, related properties under the sub-linear expectations and present the preliminary lemmas that are useful to prove the main results. In Section 3, complete convergence and complete moment convergence theorems for negatively dependence random variables under sub-linear expectation space are established.

2. Preliminaries

We use the framework and notations of Peng (2009 [16]). Let (Ω, \mathcal{F}) be a given measurable space and let \mathcal{H} be a linear space of real functions defined on (Ω, \mathcal{F}) such that if $X_1, \dots, X_n \in \mathcal{H}$ then $\varphi(X_1, \dots, X_n) \in \mathcal{H}$ for each $\varphi \in C_{l,Lip}(\mathbb{R}_n)$, where $C_{l,Lip}(\mathbb{R}_n)$ denotes the linear space of (local Lipschitz) functions φ satisfying

$$|\varphi(\mathbf{x}) - \varphi(\mathbf{y})| \leq c(1 + |\mathbf{x}|^m + |\mathbf{y}|^m)|\mathbf{x} - \mathbf{y}|, \quad \forall \mathbf{x}, \mathbf{y} \in \mathbb{R}_n,$$

for some $c > 0, m \in \mathbb{N}$ depending on φ . \mathcal{H} is considered as a space of random variables. In this case we denote $X \in \mathcal{H}$.

Definition 2.1. A sub-linear expectation $\hat{\mathbb{E}}$ on \mathcal{H} is a function $\hat{\mathbb{E}} : \mathcal{H} \rightarrow \bar{\mathbb{R}}$ satisfying the following properties: for all $X, Y \in \mathcal{H}$, we have

- (a) *Monotonicity:* If $X \geq Y$ then $\hat{\mathbb{E}}X \geq \hat{\mathbb{E}}Y$;
- (b) *Constant preserving:* $\hat{\mathbb{E}}c = c$;
- (c) *Sub-additivity:* $\hat{\mathbb{E}}(X + Y) \leq \hat{\mathbb{E}}X + \hat{\mathbb{E}}Y$ whenever $\hat{\mathbb{E}}X + \hat{\mathbb{E}}Y$ is not of the form $+\infty - \infty$ or $-\infty + \infty$;
- (d) *Positive homogeneity:* $\hat{\mathbb{E}}(\lambda X) = \lambda \hat{\mathbb{E}}X, \lambda \geq 0$.

Here $\bar{\mathbb{R}} = [-\infty, \infty]$. The triple $(\Omega, \mathcal{H}, \hat{\mathbb{E}})$ is called a sub-linear expectation space.

Give a sub-linear expectation $\hat{\mathbb{E}}$, let us denote the conjugate expectation $\hat{\varepsilon}$ of $\hat{\mathbb{E}}$ by

$$\hat{\varepsilon}X := -\hat{\mathbb{E}}(-X), \quad \forall X \in \mathcal{H}.$$

From the definition, it is easily shown that for all $X, Y \in \mathcal{H}$

$$\hat{\varepsilon}X \leq \hat{\mathbb{E}}X, \quad \hat{\mathbb{E}}(X + c) = \hat{\mathbb{E}}X + c, \quad |\hat{\mathbb{E}}(X - Y)| \leq \hat{\mathbb{E}}|X - Y| \quad \text{and} \quad \hat{\mathbb{E}}(X - Y) \geq \hat{\mathbb{E}}X - \hat{\mathbb{E}}Y.$$

If $\hat{\mathbb{E}}Y = \hat{\varepsilon}Y$, then $\hat{\mathbb{E}}(X + aY) = \hat{\mathbb{E}}X + a\hat{\mathbb{E}}Y$ for any $a \in \mathbb{R}$.

Next, we consider the capacities corresponding to the sub-linear expectations. Let $\mathcal{G} \subset \mathcal{F}$. A function $V : \mathcal{G} \rightarrow [0, 1]$ is called a capacity if

$$V(\emptyset) = 0, \quad V(\Omega) = 1 \quad \text{and} \quad V(A) \leq V(B) \quad \text{for} \quad \forall A \subseteq B, A, B \in \mathcal{G}.$$

It is called to be sub-additive if $V(A \cup B) \leq V(A) + V(B)$ for all $A, B \in \mathcal{G}$ with $A \cup B \in \mathcal{G}$. In the sub-linear space $(\Omega, \mathcal{H}, \hat{\mathbb{E}})$, we denote a pair (\mathbb{V}, ν) of capacities by

$$\mathbb{V}(A) := \inf\{\hat{\mathbb{E}}\xi; I(A) \leq \xi, \xi \in \mathcal{H}\}, \quad \nu(A) := 1 - \mathbb{V}(A^c), \forall A \in \mathcal{F},$$

where A^c is the complement set of A . By definition of \mathbb{V} and ν , it is obvious that \mathbb{V} is sub-additive, and

$$\nu(A) \leq \mathbb{V}(A), \quad \forall A \in \mathcal{F}; \quad \mathbb{V}(A) = \hat{\mathbb{E}}(I(A)), \quad \nu(A) = \hat{\mathbb{E}}(I(A)), \quad \text{if } I(A) \in \mathcal{H},$$

$$\hat{\mathbb{E}}f \leq \mathbb{V}(A) \leq \hat{\mathbb{E}}g, \quad \varepsilon f \leq \nu(A) \leq \varepsilon g, \quad \text{if } f \leq I(A) \leq g, \quad f, g \in \mathcal{H}. \tag{1}$$

This implies Markov inequality: $\forall X \in \mathcal{H}$,

$$\mathbb{V}(|X| \geq x) \leq \hat{\mathbb{E}}(|X|^p)/x^p, \quad \forall x > 0, p > 0$$

from $I(|X| \geq x) \leq |X|^p/x^p \in \mathcal{H}$. By Lemma 4.1 in Zhang (2016b [27]), we have Hölder inequality: $\forall X, Y \in \mathcal{H}, p, q > 1$ satisfying $p^{-1} + q^{-1} = 1$,

$$\hat{\mathbb{E}}(|XY|) \leq (\hat{\mathbb{E}}(|X|^p))^{1/p} (\hat{\mathbb{E}}(|Y|^q))^{1/q},$$

particularly, Jensen inequality: $\forall X \in \mathcal{H}$,

$$(\hat{\mathbb{E}}(|X|^r))^{1/r} \leq (\hat{\mathbb{E}}(|X|^s))^{1/s} \quad \text{for } 0 < r \leq s.$$

Also, we define the Choquet integrals/expectations $(C_{\mathbb{V}}, C_{\nu})$ by

$$C_{\nu}(X) := \int_0^{\infty} V(X > x)dx + \int_{-\infty}^0 (V(X > x) - 1)dx$$

with V being replaced by \mathbb{V} and ν respectively.

Definition 2.2. (Peng 2006 [13], Zhang 2016a [26])

(i) (Identical distribution) Let \mathbf{X}_1 and \mathbf{X}_2 be two n -dimensional random vectors defined respectively in sub-linear expectation spaces $(\Omega_1, \mathcal{H}_1, \hat{\mathbb{E}}_1)$ and $(\Omega_2, \mathcal{H}_2, \hat{\mathbb{E}}_2)$. They are called identically distributed if

$$\hat{\mathbb{E}}_1(\varphi(\mathbf{X}_1)) = \hat{\mathbb{E}}_2(\varphi(\mathbf{X}_2)), \quad \forall \varphi \in C_{l,Lip}(\mathbb{R}^n),$$

whenever the sub-expectations are finite. A sequence $\{X_n; n \geq 1\}$ of random variables is said to be identically distributed if for each $i \geq 1$, X_i and X_1 are identically distributed.

(ii) (Negative dependence) In a sub-linear expectation space $(\Omega, \mathcal{H}, \hat{\mathbb{E}})$, a random vector $\mathbf{Y} = (Y_1, \dots, Y_n)$, $Y_i \in \mathcal{H}$ is said to be negatively dependent (ND) to another random vector $\mathbf{X} = (X_1, \dots, X_m)$, $X_i \in \mathcal{H}$ under $\hat{\mathbb{E}}$ if for each pair of test functions $\varphi_1 \in C_{l,Lip}(\mathbb{R}_m)$ and $\varphi_2 \in C_{l,Lip}(\mathbb{R}_n)$ we have $\hat{\mathbb{E}}(\varphi_1(\mathbf{X})\varphi_2(\mathbf{Y})) \leq \hat{\mathbb{E}}(\varphi_1(\mathbf{X}))\hat{\mathbb{E}}(\varphi_2(\mathbf{Y}))$, whenever φ_1, φ_2 are coordinatewise nondecreasing or φ_1, φ_2 are coordinatewise non-increasing with $\varphi_1(\mathbf{X}) \geq 0, \hat{\mathbb{E}}\varphi_2(\mathbf{Y}) \geq 0, \hat{\mathbb{E}}|\varphi_1(\mathbf{X})\varphi_2(\mathbf{Y})| < \infty, \hat{\mathbb{E}}|\varphi_1(\mathbf{X})| < \infty, \hat{\mathbb{E}}|\varphi_2(\mathbf{Y})| < \infty$.

A sequence of random variables $\{X_n; n \geq 1\}$ is said to be negatively dependent if X_{i+1} is negatively dependent to (X_1, \dots, X_i) for each $i \geq 1$.

It is obvious that, if $\{X_n; n \geq 1\}$ is a sequence of negatively dependent random variables and functions $f_1(x), f_2(x), \dots \in C_{l,Lip}(\mathbb{R})$ are all non-decreasing (resp. all non-increasing), then $\{f_n(X_n); n \geq 1\}$ is also a sequence of negatively dependent random variables.

In the following, let $\{X_n; n \geq 1\}$ be a sequence of random variables in $(\Omega, \mathcal{H}, \hat{\mathbb{E}})$, and $S_n = \sum_{i=1}^n X_i$. The symbol c stands for a generic positive constant which may differ from one place to another. Let $a_x \sim b_x$ denote $\lim_{x \rightarrow \infty} a_x/b_x = 1$, $a_x \ll b_x$ denote that there exists a constant $c > 0$ such that $a_x \leq cb_x$ for sufficiently large x , and $I(\cdot)$ denote an indicator function.

To prove our results, we need the following two lemmas.

Lemma 2.3. (Zhang 2016b, Theorem 3.1 [27]) Let $\{X_k; k \geq 1\}$ be a sequence of negatively dependent random variables in $(\Omega, \mathcal{H}, \mathbb{E})$ with $\hat{\mathbb{E}}X_k \leq 0$. Then for any $x, y > 0$

$$\mathbb{V}(S_n \geq y) \leq \mathbb{V}\left(\max_{1 \leq k \leq n} X_k > x\right) + \exp\left(-\frac{y^2}{2(xy + B_n)} \left\{1 + \frac{2}{3} \ln\left(1 + \frac{xy}{B_n}\right)\right\}\right),$$

where $B_n = \sum_{k=1}^n \hat{\mathbb{E}}X_k^2$.

Lemma 2.4. Suppose $X \in \mathcal{H}, \alpha > 1$.

(i) Then

$$C_{\mathbb{V}}(\exp(\ln^\alpha |X|)) < \infty \Leftrightarrow \sum_{n=1}^{\infty} \exp(\ln^\alpha n) \frac{\ln^{\alpha-1} n}{n} \mathbb{V}(|X| > n) < \infty. \tag{2}$$

(ii) If $C_{\mathbb{V}}(\exp(\ln^\alpha |X|)) < \infty$, and $f(x) := \frac{cx}{\ln^\alpha x}$ for any $c > 0$ then

$$\lim_{n \rightarrow \infty} \exp(\ln^\alpha f(n)) \mathbb{V}(|X| > f(n)) = 0. \tag{3}$$

Proof (i) Note that

$$\begin{aligned} C_{\mathbb{V}}(\exp(\ln^\alpha |X|)) &\sim \int_1^\infty \mathbb{V}(\exp(\ln^\alpha |X|) > x) dx \\ &= \int_1^\infty \mathbb{V}(|X| > \exp(\ln^{1/\alpha} x)) dx \quad (\text{let } \exp(\ln^{1/\alpha} x) = y) \\ &= \int_1^\infty \frac{\alpha \ln^{\alpha-1} y \exp(\ln^\alpha y)}{y} \mathbb{V}(|X| > y) dy, \end{aligned} \tag{4}$$

and

$$\begin{aligned} \frac{\exp(\ln^\alpha(n+1))}{\exp(\ln^\alpha n)} &= \exp\left(\ln^\alpha n \left[\left(\frac{\ln(n+1)}{\ln n}\right)^\alpha - 1\right]\right) \\ &= \exp\left(\ln^\alpha n \left[\left(1 + \frac{\ln(1+1/n)}{\ln n}\right)^\alpha - 1\right]\right) \\ &\sim \exp\left(\ln^\alpha n \frac{\alpha \ln(1+1/n)}{\ln n}\right) \sim \exp\left(\frac{\alpha \ln^{\alpha-1} n}{n}\right) \\ &\rightarrow 1. \end{aligned}$$

Therefore, (2) follows from (4) and $n^{-1} \ln^{\alpha-1} n \exp(\ln^\alpha n) \sim (n+1)^{-1} \ln^{\alpha-1}(n+1) \exp(\ln^\alpha(n+1))$.

(ii) By $\int_1^\infty \mathbb{V}(|X| > \exp(\ln^{1/\alpha} x)) dx < \infty$ and $\mathbb{V}(|X| > \exp(\ln^{1/\alpha} x)) \downarrow$, we get

$$\lim_{x \rightarrow \infty} x \mathbb{V}(|X| > \exp(\ln^{1/\alpha} x)) = 0.$$

This is equivalent to

$$\lim_{y \rightarrow \infty} \exp(\ln^\alpha f(y)) \mathbb{V}(|X| > f(y)) = 0.$$

Therefore, (3) holds.

3. Complete Convergence and Complete Moment Convergence Theorems

The concept of complete convergence of a sequence of random variables was introduced by Hsu and Robbins (1947 [9]). Chow (1988 [4]) first investigated the complete moment convergence, which is more

exact than complete convergence. Complete convergence and complete moment convergence are two of the most important problems in probability theory. Their recent results can be found in Wu (2015 [22]), Wu and Jiang (2016 [23]), Xu and Tang (2014 [25]), Guo et al. (2014 [7]), Gut and Stadtmüller (2011 [8]), Qiu and Chen (2014 [17]), Wang and Hu (2014 [20]), Wang et al (2015 [21]), Liu et al (2015 [10]), Chen and Sung (2016 [1]), Tan et al (2016 [19]), and Shen et al (2017 [18]). In sub-linear expectations, due to the uncertainty of expectation and capacity, the complete convergence is essentially different from the ordinary probability space. The study of complete convergence and complete moment convergence for sub-linear expectations are much more complex and difficult. The purpose of this paper is to extend corresponding results obtained by Gut and Stadtmüller (2011 [8]), Qiu and Chen (2014 [17]), and and Wu and Jiang (2016 [23]) from the probabilistic space to sub-linear expectation space. Our results are as follows.

Theorem 3.1. *Let $\alpha > 1$, $\{X, X_n; n \geq 1\}$ be a sequence of negatively dependent and identically distributed random variables with $\hat{\mathbb{E}}(X^2) < \infty$. Suppose that*

$$C_V(\exp(\ln^\alpha |X|)) < \infty, \tag{5}$$

then

$$\sum_{n=1}^{\infty} \exp(\ln^\alpha n) \frac{\ln^{\alpha-1} n}{n^2} \mathbb{V} \left(\sum_{i=1}^n (X_i - \hat{\mathbb{E}}X_i) > n\beta \right) < \infty \text{ for all } \beta > 1, \tag{6}$$

and

$$\sum_{n=1}^{\infty} \exp(\ln^\alpha n) \frac{\ln^{\alpha-1} n}{n^2} \mathbb{V} \left(\sum_{i=1}^n (X_i - \hat{\varepsilon}X_i) < -n\beta \right) < \infty \text{ for all } \beta > 1. \tag{7}$$

In particular, if $\hat{\mathbb{E}}X_i = \hat{\varepsilon}X_i$, then

$$\sum_{n=1}^{\infty} \exp(\ln^\alpha n) \frac{\ln^{\alpha-1} n}{n^2} \mathbb{V} \left(\left| \sum_{i=1}^n (X_i - \hat{\mathbb{E}}X_i) \right| > n\beta \right) < \infty \text{ for all } \beta > 1, \tag{8}$$

Theorem 3.2. *Assume that the conditions of Theorem 3.1 hold and $\hat{\mathbb{E}}X_i = \hat{\varepsilon}X_i$. Then*

$$\sum_{n=1}^{\infty} \exp(\ln^\alpha n) \frac{\ln^{\alpha-1} n}{n^{2+q}} C_V \left\{ \left| \sum_{i=1}^n (X_i - \hat{\mathbb{E}}X_i) \right| - \beta n \right\}_+^q < \infty \text{ for all } \beta > 1 \text{ and all } q > 0. \tag{9}$$

Remark 3.3. *Theorems 3.1-3.2 extend the corresponding results obtained by Gut and Stadtmüller (2011 [8]), Qiu and Chen (2014 [17]), and Wu and Jiang (2016 [23]) from the probability space to sub-linear expectation space.*

Proof of Theorem 3.1. Without loss of generality, we can assume that $\hat{\mathbb{E}}X_1 = 0$.

For negatively dependent random variables $\{X_n; n \geq 1\}$, in order to ensure that the truncated random variables are also negatively dependent, we need that truncated functions belong to $C_{l,Lip}$ and are non-decreasing. Let $\beta > 1$ be arbitrary, set, $\beta^{-1} < \mu < 1$, for $n \geq 1$, $b_n = \beta n / (8 \ln^\alpha n)$, define, $f_c(x) = xI(x \leq c) + cI(x > c)$ for any $c > 0$, for $1 \leq k \leq n$,

$$X'_k := f_{b_n}(X_k) = X_k I(X_k \leq b_n) + b_n I(X_k > b_n), \quad S'_n := \sum_{k=1}^n X'_k,$$

$$X''_k := (X_k - b_n)I(b_n < X_k \leq n/\mu), \quad X'''_k := (X_k - b_n)I(X_k > n/\mu).$$

Obviously, $X_k = X'_k + X''_k + X'''_k$ and $\{X'_k; k \geq 1\}$ is also a sequence of negatively dependent random variables by $f_c(x) \in C_{l,Lip}$ and $f_c(x)$ being non-decreasing. Note that

$$\begin{aligned} & \{S_n > n\beta\} \\ \subseteq & \{S_n > n\beta \text{ and } X_k \leq b_n \text{ for all } k \leq n\} \\ & \cup \{S_n > n\beta \text{ and } b_n < X_{k_0} \leq n/\mu \text{ for exactly one } k_0 \leq n \text{ and } X_j \leq b_n \text{ for all } j \neq k_0\} \\ & \cup \{X''_k \neq 0 \text{ for at least two } k \leq n\} \cup \{X'''_k \neq 0 \text{ for at least one } k \leq n\} \\ := & A_n \cup B_n \cup C_n \cup D_n. \end{aligned}$$

Therefore,

$$\mathbb{V}(S_n > n\beta) \leq \mathbb{V}(A_n) + \mathbb{V}(B_n) + \mathbb{V}(C_n) + \mathbb{V}(D_n). \tag{10}$$

It shall be noted that, in the probability space, there is an equality: $EI(|X| \leq a) = P(|X| \leq a)$, however, in the sub-linear expectation space, $\hat{\mathbb{E}}$ is defined through continuous functions in $C_{l,Lip}$ and the indicator function $I(|x| \leq a)$ is not continuous. Therefore, the expression $\hat{\mathbb{E}}I(|X| \leq a)$ does not exist. This needs to modify the indicator function by functions in $C_{l,Lip}$. To this end, we define the function $g(x) \in C_{l,Lip}(\mathbb{R})$ as follows.

For $0 < \mu < 1$, let $g(x) \in C_{l,Lip}(\mathbb{R})$ be a non-increasing function such that $0 \leq g(x) \leq 1$ for all x and $g(x) = 1$ if $x \leq \mu$, $g(x) = 0$ if $x > 1$. Then

$$I(x \leq \mu) \leq g(x) \leq I(x \leq 1), \quad I(x > 1) \leq 1 - g(x) \leq I(x > \mu). \tag{11}$$

Note that

$$|X_1 - X'_1| = (X_1 - b_n)I(X_1 > b_n) \leq |X_1|I(|X_1| > b_n) \leq \frac{X_1^2}{b_n}.$$

By $\hat{\mathbb{E}}X_1 = 0, \hat{\mathbb{E}}X_1^2 < \infty$,

$$\left| \sum_{k=1}^n \hat{\mathbb{E}}X'_k \right| = n|\hat{\mathbb{E}}X'_1| = n|\hat{\mathbb{E}}X_1 - \hat{\mathbb{E}}X'_1| \leq n\hat{\mathbb{E}}|X_1 - X'_1| \leq \frac{n}{b_n} \hat{\mathbb{E}}X^2 = c\beta \ln^\alpha n, \tag{12}$$

where $c = 8\beta^{-2}\hat{\mathbb{E}}X^2 > 0$, so that, taking $y = (n - c \ln^\alpha n)\beta, x = 2b_n$ in Lemma 2.3, noting that $X'_k - \hat{\mathbb{E}}X'_k \leq 2b_n, k \leq n$, for sufficiently large n , we get

$$\begin{aligned} \mathbb{V}(A_n) &= \mathbb{V}(S'_n > n\beta) \leq \mathbb{V}\left(\sum_{k=1}^n (X'_k - \hat{\mathbb{E}}X'_k) > (n - c \ln^\alpha n)\beta\right) \\ &\leq \exp\left(-\frac{(n - c \ln^\alpha n)^2 \beta^2}{2\left(\frac{\beta^2 n(n - c \ln^\alpha n)}{4 \ln^\alpha n} + n \hat{\mathbb{E}}X^2\right)}\right) \\ &= \exp\left(-\frac{2\beta^2 \left(1 - \frac{c \ln^\alpha n}{n}\right)^2}{\beta^2 \left(1 - \frac{c \ln^\alpha n}{n}\right) + \frac{4\hat{\mathbb{E}}X^2 \ln^\alpha n}{n}} \ln^\alpha n\right) \\ &\leq \exp(-\ln^\alpha n), \end{aligned} \tag{13}$$

from $\frac{2\beta^2 \left(1 - \frac{c \ln^\alpha n}{n}\right)^2}{\beta^2 \left(1 - \frac{c \ln^\alpha n}{n}\right) + \frac{4\hat{\mathbb{E}}X^2 \ln^\alpha n}{n}} \rightarrow 2 > 1$ as $n \rightarrow \infty$.

By $\sum_{1 \leq i \leq n, i \neq k_0} X'_i$ and X_{k_0} are negatively dependent random variable, and set, $0 < \delta := 1 - (\mu\beta)^{-1} < 1$, we get from (1) and (11)

$$\begin{aligned} \mathbb{V}(B_n) &\leq \mathbb{V}\left(\exists 1 \leq k_0 \leq n \text{ such that } \sum_{1 \leq i \leq n, i \neq k_0} X'_i > \beta n - n/\mu, X_{k_0} > b_n\right) \\ &\leq \sum_{k_0=1}^n \mathbb{V}\left(\sum_{1 \leq i \leq n, i \neq k_0} X'_i > \beta n - n/\mu = \beta\delta n, X_{k_0} > b_n\right) \\ &\leq \sum_{k_0=1}^n \hat{\mathbb{E}}\left[\left(1 - g\left(\frac{\sum_{1 \leq i \leq n, i \neq k_0} X'_i}{\beta\delta n}\right)\right)\left(1 - g\left(\frac{X_{k_0}}{b_n}\right)\right)\right] \\ &\leq \sum_{k_0=1}^n \hat{\mathbb{E}}\left(1 - g\left(\frac{\sum_{1 \leq i \leq n, i \neq k_0} X'_i}{\beta\delta n}\right)\right) \hat{\mathbb{E}}\left(1 - g\left(\frac{X}{b_n}\right)\right) \\ &\leq \sum_{k_0=1}^n \mathbb{V}\left(\sum_{1 \leq i \leq n, i \neq k_0} X'_i > \mu\beta\delta n\right) \mathbb{V}(|X| > \mu b_n). \end{aligned} \tag{14}$$

Similarly to the proof of (12), we have $\left|\sum_{1 \leq i \leq n, i \neq k_0} \hat{\mathbb{E}}X'_i\right| \leq \mu\beta c \ln^\alpha n$, where $c = 8\beta^{-1}\mu^{-1}\hat{\mathbb{E}}X^2 > 0$, so that, taking $y = \mu\beta(\delta n - c \ln^\alpha n)$, $x = 2b_n$ in Lemma 2.3, using the fact that $\frac{2\beta^2(\delta - \frac{c \ln^\alpha n}{n})^2}{\beta^2(\delta - \frac{c \ln^\alpha n}{n}) + \frac{4\hat{\mathbb{E}}X^2(n-1)\ln^\alpha n}{n^2}} \rightarrow 2 > 1$ as $n \rightarrow \infty$, for sufficiently large n , we get

$$\begin{aligned} &\mathbb{V}\left(\sum_{1 \leq i \leq n, i \neq k_0} X'_i > \mu\beta\delta n\right) \\ &\leq \mathbb{V}\left(\sum_{1 \leq i \leq n, i \neq k_0} (X'_i - \hat{\mathbb{E}}X'_i) > \mu\beta(\delta n - c \ln^\alpha n)\right) \\ &\leq \exp\left(-\frac{\mu^2\beta^2(\delta n - c \ln^\alpha n)^2}{2\left(\frac{\mu\beta^2 n(\delta n - c \ln^\alpha n)}{4 \ln^\alpha n} + (n-1)\hat{\mathbb{E}}X^2\right)}\right) \\ &= \exp\left(-\frac{2\mu^2\beta^2\left(\delta - \frac{c \ln^\alpha n}{n}\right)^2}{\mu\beta^2\left(\delta - \frac{c \ln^\alpha n}{n}\right) + \frac{4\hat{\mathbb{E}}X^2(n-1)\ln^\alpha n}{n^2}} \ln^\alpha n\right) \\ &\leq \exp(-\mu\delta \ln^\alpha n). \end{aligned} \tag{15}$$

By (3), and for any $\theta > 0$, $(\ln n + \ln(\mu\beta/8) - \alpha \ln \ln n)^\alpha / \ln^\alpha n \rightarrow 1 > 1 - \theta$ as $n \rightarrow \infty$, for sufficiently large n , $\ln^\alpha(\mu b_n) = (\ln n + \ln(\mu\beta/8) - \alpha \ln \ln n)^\alpha \geq (1 - \theta) \ln^\alpha n$, thus,

$$\mathbb{V}(|X| > \mu b_n) \ll \exp(-\ln^\alpha(\mu b_n)) \leq \exp(-(1 - \theta) \ln^\alpha n) \text{ for any } \theta > 0, \tag{16}$$

Substituting (15) and (16) in (14), we obtain

$$\mathbb{V}(B_n) \ll n \exp(-\mu\delta \ln^\alpha n) \exp(-(1 - \mu\delta/2) \ln^\alpha n) \leq \exp(-\ln^\alpha n). \tag{17}$$

By (1), (11), and (16),

$$\begin{aligned}
 \mathbb{V}(C_n) &= \mathbb{V}\left(\exists 1 \leq k_1 < k_2 \leq n \text{ such that } X_{k_1}'' \neq 0, X_{k_2}'' \neq 0\right) \\
 &\leq \sum_{1 \leq k_1 < k_2 \leq n} \mathbb{V}(X_{k_1} > b_n, X_{k_2} > b_n) \\
 &\leq \sum_{1 \leq k_1 < k_2 \leq n} \hat{\mathbb{E}}\left[\left(1 - g\left(\frac{X_{k_1}}{b_n}\right)\right)\left(1 - g\left(\frac{X_{k_2}}{b_n}\right)\right)\right] \\
 &\leq \sum_{1 \leq k_1 < k_2 \leq n} \hat{\mathbb{E}}\left(1 - g\left(\frac{X_{k_1}}{b_n}\right)\right) \hat{\mathbb{E}}\left(1 - g\left(\frac{X_{k_2}}{b_n}\right)\right) \\
 &\leq n^2 \hat{\mathbb{E}}^2\left(1 - g\left(\frac{X}{b_n}\right)\right) \leq n^2 \mathbb{V}(|X| > \mu b_n) \\
 &\leq n^2 \exp(-2(1 - \delta/2) \ln^\alpha n) = n^2 \exp(-1 - (1 - \delta) \ln^\alpha n) \\
 &= \exp(-\ln^\alpha n) \frac{n^2}{(e^{\ln n})^{(1-\delta) \ln^{\alpha-1} n}} \\
 &\leq \exp(-\ln^\alpha n).
 \end{aligned} \tag{18}$$

From (1) and (11),

$$\begin{aligned}
 \mathbb{V}(D_n) &\leq \sum_{k=1}^n \mathbb{V}(X_k > n/\mu) \leq \sum_{k=1}^n \hat{\mathbb{E}}\left(1 - g\left(\frac{\mu X_k}{n}\right)\right) \\
 &= n \hat{\mathbb{E}}\left(1 - g\left(\frac{\mu X}{n}\right)\right) \leq n \mathbb{V}(|X| > n).
 \end{aligned} \tag{19}$$

This, together with (2), (5), (10), (13), (17), and (18), shows

$$\begin{aligned}
 &\sum_{n=1}^{\infty} \exp(\ln^\alpha n) \frac{\ln^{\alpha-1} n}{n^2} \mathbb{V}(S_n > \beta n) \\
 &\ll \sum_{n=1}^{\infty} \frac{\ln^{\alpha-1} n}{n^2} + \sum_{n=1}^{\infty} \exp(\ln^\alpha n) \frac{\ln^{\alpha-1} n}{n} \mathbb{V}(|X| > n) \\
 &< \infty.
 \end{aligned}$$

That is, (6) holds.

Obviously, $\{-X, -X_k; k \geq 1\}$ also satisfies the conditions of Theorem 3.1. Considering $\{-X_n; n \geq 1\}$ instead of $\{X_n; n \geq 1\}$ in (6), we can obtain (7).

In particular, if $\hat{\mathbb{E}}X_k = \varepsilon X_k$, then (8) follows from (6), (7), and

$$\begin{aligned}
 &\mathbb{V}\left(\left|\sum_{i=1}^n (X_i - \hat{\mathbb{E}}X_i)\right| > n\beta\right) \\
 &\leq \mathbb{V}\left(\sum_{i=1}^n (X_i - \hat{\mathbb{E}}X_i) > n\beta\right) + \mathbb{V}\left(\sum_{i=1}^n (X_i - \hat{\mathbb{E}}X_i) < -n\beta\right).
 \end{aligned}$$

This completes the proof of Theorem 3.1.

Proof of Theorem 3.2. Without loss of generality, we still assume that $\hat{\mathbb{E}}X_1 = 0$. Note that

$$\begin{aligned} & \sum_{n=1}^{\infty} \exp(\ln^\alpha n) \frac{\ln^{\alpha-1} n}{n^{2+q}} C_V(|S_n| - \beta n)_+^q \\ = & \beta^q \sum_{n=1}^{\infty} \exp(\ln^\alpha n) \frac{\ln^{\alpha-1} n}{n^{2+q}} \int_0^n qx^{q-1} \mathbb{V}(|S_n| - \beta n > \beta x) dx \\ & + \beta^q \sum_{n=1}^{\infty} \exp(\ln^\alpha n) \frac{\ln^{\alpha-1} n}{n^{2+q}} \int_n^\infty qx^{q-1} \mathbb{V}(|S_n| - \beta n > \beta x) dx \\ \ll & \sum_{n=1}^{\infty} \exp(\ln^\alpha n) \frac{\ln^{\alpha-1} n}{n^2} \mathbb{V}(|S_n| > \beta n) \\ & + \sum_{n=1}^{\infty} \exp(\ln^\alpha n) \frac{\ln^{\alpha-1} n}{n^{2+q}} \int_n^\infty x^{q-1} \mathbb{V}(|S_n| > \beta x) dx. \end{aligned}$$

Hence, by (8), in order to establish (9), it suffices to prove that

$$\sum_{n=1}^{\infty} \exp(\ln^\alpha n) \frac{\ln^{\alpha-1} n}{n^{2+q}} \int_n^\infty x^{q-1} \mathbb{V}(|S_n| > \beta x) dx < \infty. \tag{20}$$

Let $\beta > 1$ be an arbitrary, and $\beta^{-1} < \mu < 1$, set, for $x \geq n$, $b_x = \beta x / (8 \ln^\alpha x)$, define, for $1 \leq k \leq n$,

$$Y'_k := X_k I(X_k \leq b_x) + b_x I(X_k > b_x), \quad U'_n := \sum_{k=1}^n Y'_k,$$

$$Y''_k := (X_k - b_x) I(b_x < X_k \leq x/\mu), \quad Y'''_k := (X_k - b_x) I(X_k > x/\mu).$$

By similar methods to the proof of (10), we have

$$\mathbb{V}(S_n > x\beta) \leq \mathbb{V}(A_x) + \mathbb{V}(B_x) + \mathbb{V}(C_x) + \mathbb{V}(D_x), \tag{21}$$

which leads to

$$A_x = \{U'_n > x\beta\},$$

$$B_x = \{S_n > x\beta \text{ and } b_x < X_{k_0} \leq x/\mu \text{ for exactly one } k_0 \leq n \text{ and } X_j \leq b_x \text{ for all } j \neq k_0\},$$

$$C_x = \{Y''_k \neq 0 \text{ for at least two } k \leq n\},$$

$$D_x = \{Y'''_k \neq 0 \text{ for at least one } k \leq n\}.$$

Using similar methods to those used in the proof of (12), (13), and (16)-(19), for $0 < \delta := 1 - \beta^{-1} < 1$ and $x \geq n$, we have $|\sum_{k=1}^n \hat{\mathbb{E}}Y'_k| \leq c\beta \ln^\alpha x$, where $c = 8\beta^{-2} \hat{\mathbb{E}}X^2$, and

$$\mathbb{V}(A_x) \ll \exp(-\ln^\alpha x),$$

$$\mathbb{V}(|X| > \mu b_x) \ll \exp(-(1 - \theta) \ln^\alpha x) \text{ for any } \theta > 0,$$

$$\begin{aligned} \mathbb{V}(B_x) & \ll n \exp(-\mu\delta \ln^\alpha x - (1 - \mu\delta/2) \ln^\alpha x) \\ & = \exp(-\ln^\alpha x) \frac{n}{\exp(\mu\delta \ln^\alpha x/2)} \leq \exp(-\ln^\alpha x), \end{aligned}$$

$$\begin{aligned} \mathbb{V}(C_x) & \leq n^2 \mathbb{V}^2(|X| > \mu b_x) \ll n^2 \exp(-2(1 - \delta/2) \ln^\alpha x) \\ & = \exp(-\ln^\alpha x) n^2 \exp(-(1 - \delta) \ln^\alpha x) \leq \exp(-\ln^\alpha x), \end{aligned}$$

$$\mathbb{V}(D_x) \leq n\mathbb{V}(|X| > x),$$

which, combining with (21), shows

$$\mathbb{V}(S_n > x\beta) \ll \exp(-\ln^\alpha x) + n\mathbb{V}(|X| > x).$$

Because $\{-X, -X_k; k \geq 1\}$ is also a sequence of negatively dependent random variables. Obviously, $\{-X, -X_k; k \geq 1\}$ also satisfies the condition (5) and $\hat{\mathbb{E}}(-X_k) = -\varepsilon X_k = -\hat{\mathbb{E}}(X_k) = 0$ from the assumption: $\hat{\mathbb{E}}(X_k) = \varepsilon X_k$. Therefore, replacing X_k by $-X_k$ in the above inequality, we get

$$\mathbb{V}(S_n < -x\beta) = \mathbb{V}(-S_n > x\beta) \ll \exp(-\ln^\alpha x) + n\mathbb{V}(|X| > x).$$

Therefore,

$$\begin{aligned} \mathbb{V}(|S_n| > x\beta) &\leq \mathbb{V}(S_n > x\beta) + \mathbb{V}(S_n < -x\beta) \\ &\ll \exp(-\ln^\alpha x) + n\mathbb{V}(|X| > x). \end{aligned}$$

Hence,

$$\begin{aligned} &\int_n^\infty x^{q-1} \mathbb{V}(|S_n| > x\beta) \, dx \\ &\ll \int_n^\infty x^{q-1} \exp(-\ln^\alpha x) \, dx + \int_n^\infty x^{q-1} n \mathbb{V}(|X| > x) \, dx \\ &:= I_1 + I_2. \end{aligned} \tag{22}$$

By the fact that $(a + b)^\alpha \geq a^\alpha + b^\alpha$ for any $a, b > 0$ and $\alpha > 1$, let $t = x/n$, we get

$$\begin{aligned} \sum_{n=1}^\infty \exp(\ln^\alpha n) \frac{\ln^{\alpha-1} n}{n^{2+q}} I_1 &= \sum_{n=1}^\infty \exp(\ln^\alpha n) \frac{\ln^{\alpha-1} n}{n^{2+q}} \int_1^\infty n^q t^{q-1} \exp(-(\ln n + \ln t)^\alpha) \, dt \\ &\leq \sum_{n=1}^\infty \exp(\ln^\alpha n) \frac{\ln^{\alpha-1} n}{n^{2+q}} n^q \exp(-\ln^\alpha n) \int_1^\infty t^{q-1} \exp(-\ln^\alpha t) \, dt \\ &\ll \sum_{n=1}^\infty \exp(\ln^\alpha n) \frac{\ln^{\alpha-1} n}{n^{2+q}} n^q \exp(-\ln^\alpha n) \\ &= \sum_{n=1}^\infty \frac{\ln^{\alpha-1} n}{n^2} < \infty. \end{aligned} \tag{23}$$

By (5) and (2),

$$\begin{aligned}
& \sum_{n=1}^{\infty} \exp(\ln^{\alpha} n) \frac{\ln^{\alpha-1} n}{n^{2+q}} I_2 \\
&= \sum_{n=1}^{\infty} \exp(\ln^{\alpha} n) \frac{\ln^{\alpha-1} n}{n^{1+q}} \int_n^{\infty} x^{q-1} \mathbb{V}(|X| > x) dx \\
&= \sum_{n=1}^{\infty} \exp(\ln^{\alpha} n) \frac{\ln^{\alpha-1} n}{n^{1+q}} \sum_{j=n}^{\infty} \int_j^{j+1} x^{q-1} \mathbb{V}(|X| > x) dx \\
&\ll \sum_{n=1}^{\infty} \exp(\ln^{\alpha} n) \frac{\ln^{\alpha-1} n}{n^{1+q}} \sum_{j=n}^{\infty} \mathbb{V}(|X| > j) j^{q-1} \\
&= \sum_{j=1}^{\infty} \mathbb{V}(|X| > j) j^{q-1} \sum_{n=1}^j \exp(\ln^{\alpha} n) \frac{\ln^{\alpha-1} n}{n^{1+q}} \\
&\ll \sum_{j=1}^{\infty} \exp(\ln^{\alpha} j) \frac{\ln^{\alpha-1} j}{j} \mathbb{V}(|X| > j) \\
&< \infty,
\end{aligned}$$

from which, combining with (22) and (23), we see that (20) holds. This completes the proof of Theorem 3.2.

References

- [1] Chen, P.Y. and Sung, S. H., 2016. Complete convergence and strong laws of large numbers for weighted sums of negatively orthant dependent random variables. *Acta Math. Hungar.*, 148(1): 83-95.
- [2] Chen, Z.J., 2016. Strong laws of large numbers for sub-linear expectations. *Science China Mathematics*, 59 (5): 945-954.
- [3] Cheng, H., 2016. A strong law of large numbers for sub-linear expectation under a general moment condition. *Statistics and Probability Letters*. 119, 248-258.
- [4] Chow, Y., 1988. On the rate of moment convergence of sample sums and extremes. *Bull. Inst. Math. Acad. Sin.* 16, 177-201.
- [5] Denis, L. and Martini, C., 2006. A theoretical framework for the pricing of contingent claims in the presence of model uncertainty. *Ann Appl Probab*, 16(2): 827-852.
- [6] Gilboa, I., 1987. Expected utility theory with purely subjective non-additive probabilities. *J Math Econom*, 16: 65-68.
- [7] Guo, M.L., Xu, C.Y., Zhu, D.J., 2014. Complete convergence of weighted sums for arrays of rowwise m -negatively associated random variables, *Communications in Mathematical Research*, 30(1), 41-50.
- [8] Gut, A., Stadtmüller, U., 2011. An intermediate Baum-Katz theorem. *Statist. Probab. Lett.* 81, 1486-1492.
- [9] Hsu, P.L., Robbins, H., 1947. Complete convergence and the law of large numbers. *Proc Nat Acad Sci USA*. 33, 25-31. doi:10.1073/pnas.33.2.25
- [10] Liu, C.C., Guo, M.L., Zhu, D., 2015. Equivalent conditions of complete convergence for weighted sums of sequences of extended negatively dependent random variables. *Communications in Mathematical Research*, 31(1): 40-50.
- [11] Marinacci, M., 1999. Limit laws for non-additive probabilities and their frequentist interpretation. *J Econom Theory*. 84: 145-195.
- [12] Peng, S., 1997. BSDE and related g -expectation. *Pitman Research Notes in Mathematics Series*, 364: 141-159.
- [13] Peng, S., 1999. Monotonic limit theorem of BSDE and nonlinear decomposition theorem of Doob-Meyer type. *Probab Theory Related Fields*, 113: 473-499.
- [14] Peng, S., 2006. G -expectation, G -Brownian motion and related stochastic calculus of Ito type. *Proceedings of the 2005 Abel Symposium*.
- [15] Peng, S., 2008. Multi-dimensional G -Brownian motion and related stochastic calculus under G -expectation. *Stochastic Process Appl*, 118(12): 2223-2253.
- [16] Peng, S., 2009. Survey on normal distributions, central limit theorem, Brownian motion and the related stochastic calculus under sublinear expectations. *Science in China Series A-Mathematics*, 52(7): 1391-1411.
- [17] Qiu D.H. and Chen P.Y., 2014. Complete moment convergence for i.i.d. random variables, *Statistics and Probability Letters* 91, 76-82
- [18] Shen, A.T., Xue, M.X., Wang, W.J., 2017. Complete convergence for weighted sums of extended negatively dependent random variables, *Communications in Statistics-Theory and Methods*, 46(3): 1433-1444.
- [19] Tan, X.L., Wang, H., Zhang, Y., 2016. Complete convergence of the non-identically distributed pairwise NQD random sequences. *Communications in Statistics Theory and Methods*, 45(9): 2626-2637.
- [20] Wang, X.J., and Hu, S.H., 2014. Complete convergence and complete moment convergence for martingale difference sequence. *Acta Mathematica Sinica-English Series*, 30(1): 119-132.

- [21] Wang, X.J., Zheng, L.L., Xu, C., Hu, S.H., 2015. *Complete consistency for the estimator of nonparametric regression models based on extended negatively dependent errors*. *Statistics: A Journal of Theoretical and Applied Statistics*, 49, 396-407.
- [22] Wu, Q.Y., 2015. *Further study complete convergence for weighted sums of PNQD random variables*, *Journal of Inequalities and Applications*, 2015:289, DOI 10.1186/s13660-015-0814-1
- [23] Wu, Q.Y., and Jiang, Y.Y., 2016. *Complete convergence and complete moment convergence for negatively associated sequences of random variables*. *Journal of Inequalities and Applications*, 2016:157 DOI 10.1186/s13660-016-1107-z
- [24] Wu, Q.Y., and Jiang, Y.Y., 2018. *Strong law of large numbers and Chover's law of the iterated logarithm under sub-linear expectations*. *Mathematical Analysis and Applications*, 460(1): 252-270.
- [25] Xu, H. and Tang, L., 2014. *On complete convergence for arrays of rowwise AANA random variables*, *Stochastics: An International Journal of Probability and Stochastic Processes*, 86 (3): 371-381.
- [26] Zhang, L.X., 2016a. *Strong limit theorems for extended independent and extended negatively dependent random variables under non-linear expectations*. arXiv preprint arXiv:1608.00710.
- [27] Zhang, L.X., 2016b. *Exponential inequalities under the sub-linear expectations with applications to laws of the iterated logarithm*. *Sci China Math*, 59 (2): 2503-2526.
- [28] Zhang, L.X., 2016c. *Rosenthal's inequalities for independent and negatively dependent random variables under sub-linear expectations with applications*. *Sci China Math*, 59 (4): 751-768.