



Applications of the Horadam Polynomials Involving λ -Pseudo-Starlike Bi-Univalent Functions Associated with a Certain Convolution Operator

H. M. Srivastava^a, A. K. Wanas^b

^aDepartment of Mathematics and Statistics, University of Victoria,
Victoria, British Columbia V8W 3R4, Canada

and

Department of Medical Research, China Medical University Hospital,
China Medical University, Taichung 40402, Taiwan, Republic of China

and

Department of Mathematics and Informatics, Azerbaijan University,
71 Jeyhun Hajibeyli Street, AZ1007 Baku, Azerbaijan

and

Section of Mathematics, International Telematic University Uninettuno, I-00186 Rome, Italy

^bDepartment of Mathematics, College of Science, University of Al-Qadisiyah, Al-Diwaniyah, Al-Qadisiyah, Iraq

Abstract. In this article, we introduce and study a new family $\mathcal{P}_{\Sigma}(\delta, \lambda, k, \gamma, \alpha, \beta, r)$ of normalized analytic and λ -pseudo-starlike bi-univalent functions by using the Horadam polynomials, which is associated with a certain convolution operator defined in the open unit disk \mathbb{U} . We establish the bounds for $|a_2|$ and $|a_3|$, where a_2, a_3 are the initial Taylor-Maclaurin coefficients. Furthermore, we obtain the Fekete-Szegő inequality for functions in the class $\mathcal{P}_{\Sigma}(\delta, \lambda, k, \gamma, \alpha, \beta, r)$, which we have introduced here. We indicate several special cases and consequences for our results. Finally, we comment on the recent usages, especially in Geometric Function Theory of Complex Analysis, of the basic (or q -) calculus and also of its trivial and inconsequential (p, q) -variation involving an obviously redundant parameter p .

1. Introduction, Motivation and Preliminaries

We denote by \mathcal{A} the class of functions which are analytic in the open unit disk

$$\mathbb{U} = \{z : z \in \mathbb{C} \text{ and } |z| < 1\}$$

and have the following normalized form:

$$f(z) = z + \sum_{n=2}^{\infty} a_n z^n. \quad (1)$$

2020 *Mathematics Subject Classification.* Primary 30C45; Secondary 11B39, 30C50, 33C05.

Keywords. Analytic functions; Univalent functions; Bi-univalent functions; Horadam polynomials; Upper bounds; Fekete-Szegő problem; Convolution operator; Subordination between analytic functions; Basic (or q -) calculus and its so-called (p, q) -variation.

Received: 03 November 2020; Accepted: 10 March 2021

Communicated by Dragan S. Djordjević

Email addresses: harimsri@math.uvic.ca (H. M. Srivastava), abbas.kareem.w@qu.edu.iq (A. K. Wanas)

We also denote by \mathcal{S} the subclass of \mathcal{A} consisting of functions which are also univalent in \mathbb{U} . According to the Koebe one-quarter theorem [10], every function $f \in \mathcal{S}$ has an inverse f^{-1} defined by

$$f^{-1}(f(z)) = z \quad (z \in \mathbb{U})$$

and

$$f(f^{-1}(w)) = w \quad \left(|w| < r_0(f); r_0(f) \geq \frac{1}{4} \right),$$

where

$$g(w) = f^{-1}(w) = w - a_2w^2 + (2a_2^2 - a_3)w^3 - (5a_2^3 - 5a_2a_3 + a_4)w^4 + \dots \tag{2}$$

A function $f \in \mathcal{A}$ is said to be bi-univalent in \mathbb{U} if both f and f^{-1} are univalent in \mathbb{U} . Let Σ stand for the class of bi-univalent functions in \mathbb{U} given by (1). For a brief historical account and for several interesting examples of functions in the class Σ , see the pioneering work on this subject by Srivastava *et al.* [47], which actually revived the study of bi-univalent functions in recent years. From the work of Srivastava *et al.* [47], we choose to recall the following examples of functions in the class Σ :

$$\frac{z}{1-z}, \quad -\log(1-z) \quad \text{and} \quad \frac{1}{2} \log\left(\frac{1+z}{1-z}\right).$$

We notice that the class Σ is not empty. However, the Koebe function is not a member of Σ .

In a considerably large number of sequels to the aforementioned work of Srivastava *et al.* [47], several different subclasses of the bi-univalent function class Σ were introduced and studied analogously by the many authors (see, for example, [8, 12, 29, 36, 39, 40, 45, 48, 53, 54, 61, 62, 64, 65]), but only non-sharp estimates on the initial coefficients $|a_2|$ and $|a_3|$ in the Taylor-Maclaurin expansion (1) were obtained in most (if not all) of the recent papers. The problem to find the general coefficient bounds on the Taylor-Maclaurin coefficients

$$|a_n| \quad (n \in \mathbb{N}; n \geq 3)$$

for functions $f \in \Sigma$ is still not completely addressed for many of the subclasses of the bi-univalent function class Σ (see, for example, [36, 48, 53]).

The Fekete-Szegő functional $|a_3 - \mu a_2^2|$ for $f \in \mathcal{S}$ is well known for its rich history in the field of Geometric Function Theory. Its origin was in the disproof by Fekete and Szegő [11] of the Littlewood-Paley conjecture that the coefficients of odd univalent functions are bounded by unity. The functional has since received great attention, particularly in the study of many subclasses of the family of univalent functions. This topic has become of considerable interest among researchers in Geometric Function Theory (see, for example, [41, 46, 52]).

Definition 1. A function $f \in \mathcal{A}$ is said to be λ -pseudo-starlike function of order ρ ($0 \leq \rho < 1$) in \mathbb{U} if (see [6])

$$\Re \left(\frac{z(f'(z))^\lambda}{f(z)} \right) > \rho \quad (z \in \mathbb{U}; \lambda \geq 1).$$

With a view to recalling the principle of subordination between analytic functions, let the functions f and g be analytic in \mathbb{U} . We say that the function f is subordinate to g , if there exists a Schwarz function ω , which is analytic in \mathbb{U} with

$$\omega(0) = 0 \quad \text{and} \quad |\omega(z)| < 1 \quad (z \in \mathbb{U}),$$

such that

$$f(z) = g(\omega(z)).$$

This subordination is denoted by

$$f < g \quad \text{or} \quad f(z) < g(z) \quad (z \in \mathbb{U}).$$

It is well known that, if the function g is univalent in \mathbb{U} , then (see [21])

$$f < g \quad (z \in \mathbb{U}) \iff f(0) = g(0) \quad \text{and} \quad f(\mathbb{U}) \subseteq g(\mathbb{U}).$$

Recently, Hörçum and Koçer [15] considered the so-called Horadam polynomials $h_n(r)$, which are given by the following recurrence relation (see also Horadam and Mahon [14]):

$$h_n(r) = prh_{n-1}(r) + qh_{n-2}(r) \quad (r \in \mathbb{R}; n \in \mathbb{N} = \{1, 2, 3, \dots\}) \tag{3}$$

with

$$h_1(r) = a \quad \text{and} \quad h_2(r) = br,$$

for some real constants a, b, p and q . The characteristic equation of the recurrence relation (3) is given by

$$t^2 - prt - q = 0.$$

This equation has the following two real roots:

$$\alpha = \frac{pr + \sqrt{p^2r^2 + 4q}}{2} \quad \text{and} \quad \beta = \frac{pr - \sqrt{p^2r^2 + 4q}}{2}.$$

Remark 1. By selecting particular values of a, b, p and q , the Horadam polynomial $h_n(r)$ reduces to several known polynomials. Some of these special cases are recorded below.

1. Taking $a = b = p = q = 1$, we obtain the Fibonacci polynomials $F_n(r)$.
2. Taking $a = 2$ and $b = p = q = 1$, we get the Lucas polynomials $L_n(r)$.
3. Taking $a = q = 1$ and $b = p = 2$, we have the Pell polynomials $P_n(r)$.
4. Taking $a = b = p = 2$ and $q = 1$, we find the Pell-Lucas polynomials $Q_n(r)$.
5. Taking $a = b = 1, p = 2$ and $q = -1$, we obtain the Chebyshev polynomials $T_n(r)$ of the first kind.
6. Taking $a = 1, b = p = 2$ and $q = -1$, we have the Chebyshev polynomials $U_n(r)$ of the second kind.

These polynomials, the families of orthogonal polynomials and other special polynomials, as well as their extensions and generalizations, are potentially important in a variety of disciplines in many branches of science, especially in the mathematical, statistical and physical sciences. For more information associated with these polynomials, see [13, 14, 18, 19].

The generating function of the Horadam polynomials $h_n(r)$ is given as follows (see [15]):

$$\Pi(r, z) = \sum_{n=1}^{\infty} h_n(r)z^{n-1} = \frac{a + (b - ap)rz}{1 - prz - qz^2}. \tag{4}$$

Srivastava *et al.* [33] have already applied the Horadam polynomials in a context involving analytic and bi-univalent functions. The investigation by Srivastava *et al.* [33] was followed by such works as those by Al-Amoush [2], Wanas and Alina [63] and Abirami *et al.* [1].

Recently, Wanas [60] introduced the following convolution operator $W_{\alpha, \beta}^{k, \gamma} : \mathcal{A} \rightarrow \mathcal{A}$ defined by

$$W_{\alpha, \beta}^{k, \gamma} f(z) = z + \sum_{n=2}^{\infty} [\Phi_n(k, \alpha, \beta)]^\gamma a_n z^n, \tag{5}$$

where

$$\Phi_n(k, \alpha, \beta) = \sum_{m=1}^k \binom{k}{m} (-1)^{m+1} \left(\frac{\alpha^m + n\beta^m}{\alpha^m + \beta^m} \right) \tag{6}$$

$$(\alpha \in \mathbb{R}; \beta \geq 0 \text{ with } \alpha + \beta > 0; m, \gamma \in \mathbb{N}_0 = \mathbb{N} \cup \{0\}).$$

Remark 2. The operator $W_{\alpha,\beta}^{k,\gamma}$ is a generalization of several known operators considered in earlier investigations which are being recalled below.

1. For $k = 1$, the operator $W_{\alpha,\beta}^{1,\gamma} \equiv I_{\alpha,\beta}^\gamma$ was introduced and studied by Swamy [58].
2. For $k = \beta = 1$, $\gamma = -\mu$, $\Re(\mu) > 1$ and $\alpha \in \mathbb{C} \setminus \mathbb{Z}_0^-$, the operator $W_{\alpha,1}^{1,-\mu} \equiv J_{\mu,\alpha}$ was investigated by Srivastava and Attiya [35]. The operator $J_{\mu,\alpha}$ is now popularly known in the literature as the Srivastava-Attiya operator. Various applications of the Srivastava-Attiya operator are found in [25, 37, 38, 42, 51] and in the references cited in each of these earlier works.
3. For $k = \beta = 1$ and $\alpha > -1$, the operator $W_{\alpha,1}^{1,\gamma} \equiv I_\alpha^\gamma$ was investigated by Cho and Srivastava [9].
4. For $k = \alpha = \beta = 1$, the operator $W_{1,1}^{1,\gamma} \equiv I^\gamma$ was considered by Uralegaddi and Somanatha [59].
5. For $k = \alpha = \beta = 1$, $\gamma = -\sigma$ and $\sigma > 0$, the operator $W_{1,1}^{1,-\sigma} \equiv I^\sigma$ was introduced by Jung *et al.* [16]. The operator I^σ is widely known as the Jung-Kim-Srivastava integral operator.
6. For $k = \beta = 1$, $\gamma = -1$ and $\alpha > -1$, the operator $W_{\alpha,1}^{1,-1} \equiv L_\alpha$ was studied by Bernardi [7].
7. For $\alpha = 0, k = \beta = 1$ and $\gamma = -1$, the operator $W_{0,1}^{1,-1} \equiv u$ was investigated by Alexander [3].
8. For $k = 1, \alpha = 1 - \beta$ and $\beta \geq 0$, the operator $W_{1-\beta,\beta}^{1,\gamma} \equiv D_\beta^\gamma$ was given by Al-Oboudi [4].
9. For $k = 1, \alpha = 0$ and $\beta = 1$, the operator $W_{0,1}^{1,\gamma} \equiv S^\gamma$ was considered by Sălăgean [24].

Remark 3. In a recently-published survey-cum-expository review article by Srivastava [29], the so-called (p, q) -calculus was exposed to be a rather trivial and inconsequential variation of the classical q -calculus, the additional parameter p being redundant or superfluous (see, for details, [29, p. 340]). Srivastava [29] also pointed out how the Hurwitz-Lerch Zeta function as well as its *multi-parameter* extension, which is popularly known as the λ -generalized Hurwitz-Lerch Zeta function (see, for details, [26]), have motivated the studies of several other families of extensively- and widely-investigated linear convolution operators which emerge essentially from the Srivastava-Attiya operator [35] (see also [27], [28], [30] and [31]).

2. A Set of Main Results

We begin this section by defining the new subclass $\mathcal{P}_\Sigma(\delta, \lambda, k, \gamma, \alpha, \beta, r)$.

Definition 2. For $\delta \in \mathbb{C} \setminus \{0\}$, $\lambda \geq 1$ and $r \in \mathbb{R}$, a function $f \in \Sigma$ is said to be in the class $\mathcal{P}_\Sigma(\delta, \lambda, k, \gamma, \alpha, \beta, r)$ if it satisfies the following subordination conditions:

$$1 + \frac{1}{\delta} \left(\frac{z \left(\left(W_{\alpha,\beta}^{k,\gamma} f(z) \right)' \right)^\lambda}{W_{\alpha,\beta}^{k,\gamma} f(z)} - 1 \right) < \Pi(r, z) + 1 - a$$

and

$$1 + \frac{1}{\delta} \left(\frac{w \left(\left(W_{\alpha, \beta}^{k, \gamma} g(w) \right)' \right)^\lambda}{W_{\alpha, \beta}^{k, \gamma} g(w)} - 1 \right) < \Pi(r, w) + 1 - a,$$

where a is a real constant and the function $g = f^{-1}$ is given by (2).

Our first main result is asserted by Theorem 1 below.

Theorem 1. For $\delta \in \mathbb{C} \setminus \{0\}$, $\lambda \geq 1$ and $r \in \mathbb{R}$, let $f \in \mathcal{A}$ be in the class $\mathcal{P}_\Sigma(\delta, \lambda, k, \gamma, \alpha, \beta, r)$. Then

$$|a_2| \leq \frac{\delta |br| \sqrt{|br|}}{\sqrt{\left| \delta \left[(3\lambda - 1)\Phi_3^\gamma(k, \alpha, \beta) + (2\lambda(\lambda - 2) + 1)\Phi_2^{2\gamma}(k, \alpha, \beta) \right] b - p(2\lambda - 1)^2 br^2 - qa(2\lambda - 1)^2 \right|}}$$

and

$$|a_3| \leq \frac{\delta |br|}{(3\lambda - 1) |\Phi_3^\gamma(k, \alpha, \beta)|} + \frac{\delta^2 b^2 r^2}{(2\lambda - 1)^2 \Phi_2^{2\gamma}(k, \alpha, \beta)}.$$

Proof. Let $f \in \mathcal{P}_\Sigma(\delta, \lambda, k, \gamma, \alpha, \beta, r)$. Then there are two analytic functions $u, v : \mathbb{U} \rightarrow \mathbb{U}$ given by

$$u(z) = u_1 z + u_2 z^2 + u_3 z^3 + \dots \quad (z \in \mathbb{U}) \tag{7}$$

and

$$v(w) = v_1 w + v_2 w^2 + v_3 w^3 + \dots \quad (w \in \mathbb{U}), \tag{8}$$

with

$$u(0) = v(0) = 0 \quad \text{and} \quad \max \{ |u(z)|, |v(w)| \} < 1 \quad (z, w \in \mathbb{U}),$$

such that

$$\frac{1}{\delta} \left(\frac{z \left(\left(W_{\alpha, \beta}^{k, \gamma} f(z) \right)' \right)^\lambda}{W_{\alpha, \beta}^{k, \gamma} f(z)} - 1 \right) = \Pi(r, u(z)) - a$$

and

$$\frac{1}{\delta} \left(\frac{w \left(\left(W_{\alpha, \beta}^{k, \gamma} g(w) \right)' \right)^\lambda}{W_{\alpha, \beta}^{k, \gamma} g(w)} - 1 \right) = \Pi(r, v(w)) - a.$$

Equivalently, we have

$$\frac{1}{\delta} \left(\frac{z \left(\left(W_{\alpha, \beta}^{k, \gamma} f(z) \right)' \right)^\lambda}{W_{\alpha, \beta}^{k, \gamma} f(z)} - 1 \right) = h_1(r) + h_2(r)u(z) + h_3(r)u^2(z) + \dots \tag{9}$$

and

$$\frac{1}{\delta} \left(\frac{w \left(\left(W_{\alpha, \beta}^{k, \gamma} g(w) \right)' \right)^\lambda}{W_{\alpha, \beta}^{k, \gamma} g(w)} - 1 \right) = h_1(r) + h_2(r)v(w) + h_3(r)v^2(w) + \dots \tag{10}$$

Combining (7), (8), (9) and (10), we find that

$$\frac{1}{\delta} \left(\frac{z \left((W_{\alpha, \beta}^{k, \gamma} f(z))' \right)^\lambda}{W_{\alpha, \beta}^{k, \gamma} f(z)} - 1 \right) = h_2(r)u_1z + [h_2(r)u_2 + h_3(r)u_1^2]z^2 + \dots \tag{11}$$

and

$$\frac{1}{\delta} \left(\frac{w \left((W_{\alpha, \beta}^{k, \gamma} g(w))' \right)^\lambda}{W_{\alpha, \beta}^{k, \gamma} g(w)} - 1 \right) = h_2(r)v_1w + [h_2(r)v_2 + h_3(r)v_1^2]w^2 + \dots \tag{12}$$

It is well known that, if

$$\max \{|u(z)|, |v(w)|\} < 1 \quad (z, w \in \mathbb{U}),$$

then

$$|u_j| \leq 1 \quad \text{and} \quad |v_j| \leq 1 \quad (\forall j \in \mathbb{N}). \tag{13}$$

Now, by comparing the corresponding coefficients in (11) and (12), and after some simplification, we have

$$\frac{(2\lambda - 1)\Phi_2^\gamma(k, \alpha, \beta)}{\delta} a_2 = h_2(r)u_1, \tag{14}$$

$$\begin{aligned} \frac{(3\lambda - 1)\Phi_3^\gamma(k, \alpha, \beta)}{\delta} a_3 + \frac{(2\lambda(\lambda - 2) + 1)\Phi_2^{2\gamma}(k, \alpha, \beta)}{\delta} a_2^2 \\ = h_2(r)u_2 + h_3(r)u_1^2, \end{aligned} \tag{15}$$

$$-\frac{(2\lambda - 1)\Phi_2^\gamma(k, \alpha, \beta)}{\delta} a_2 = h_2(r)v_1 \tag{16}$$

and

$$\begin{aligned} \frac{(3\lambda - 1)\Phi_3^\gamma(k, \alpha, \beta)}{\delta} (2a_2^2 - a_3) + \frac{(2\lambda(\lambda - 2) + 1)\Phi_2^{2\gamma}(k, \alpha, \beta)}{\delta} a_2^2 \\ = h_2(r)v_2 + h_3(r)v_1^2. \end{aligned} \tag{17}$$

It follows from (14) and (16) that

$$u_1 = -v_1 \tag{18}$$

and

$$\frac{2(2\lambda - 1)^2 \Phi_2^{2\gamma}(k, \alpha, \beta)}{\delta^2} a_2^2 = h_2^2(r)(u_1^2 + v_1^2). \tag{19}$$

If we add (15) to (17), we find that

$$\begin{aligned} \frac{2 \left[(3\lambda - 1)\Phi_3^\gamma(k, \alpha, \beta) + (2\lambda(\lambda - 2) + 1)\Phi_2^{2\gamma}(k, \alpha, \beta) \right]}{\delta} a_2^2 \\ = h_2(r)(u_2 + v_2) + h_3(r)(u_1^2 + v_1^2). \end{aligned} \tag{20}$$

Upon substituting the value of $u_1^2 + v_1^2$ from (19) into the right-hand side of (20), we deduce that

$$a_2^2 = \frac{\delta^2 h_2^3(r)(u_2 + v_2)}{2 \left[\delta h_2^2(r) \left((3\lambda - 1)\Phi_3^\gamma(k, \alpha, \beta) + (2\lambda(\lambda - 2) + 1)\Phi_2^{2\gamma}(k, \alpha, \beta) \right) - h_3(r)(2\lambda - 1)^2 \right]}. \tag{21}$$

By further computations using (3), (13) and (21), we obtain

$$|a_2| \leq \frac{\delta |br| \sqrt{|br|}}{\sqrt{\left| \left[\delta \left((3\lambda - 1)\Phi_3^\gamma(k, \alpha, \beta) + (2\lambda(\lambda - 2) + 1)\Phi_2^{2\gamma}(k, \alpha, \beta) \right) b - p(2\lambda - 1)^2 \right] br^2 - qa(2\lambda - 1)^2 \right|}}.$$

Next, if we subtract (17) from (18), we can easily see that

$$\frac{2(3\lambda - 1)\Phi_3^\gamma(k, \alpha, \beta)}{\delta} (a_3 - a_2^2) = h_2(r)(u_2 - v_2) + h_3(r)(u_1^2 - v_1^2). \tag{22}$$

Also, in view of (18) and (19), we find from (22) that

$$a_3 = \frac{\delta h_2(r)(u_2 - v_2)}{2(3\lambda - 1)\Phi_3^\gamma(k, \alpha, \beta)} + \frac{\delta^2 h_2^2(r)(u_1^2 + v_1^2)}{2(2\lambda - 1)^2 \Phi_2^{2\gamma}(k, \alpha, \beta)}.$$

Thus, by applying (3), we obtain

$$|a_3| \leq \frac{\delta |br|}{(3\lambda - 1) |\Phi_3^\gamma(k, \alpha, \beta)|} + \frac{\delta^2 b^2 r^2}{(2\lambda - 1)^2 \Phi_2^{2\gamma}(k, \alpha, \beta)}.$$

This completes the proof of Theorem 1. \square

In the next theorem, we present the Fekete-Szegő inequality for functions in the class $\mathcal{P}_\Sigma(\delta, \lambda, k, \gamma, \alpha, \beta, r)$.

Theorem 2. For $\delta \in \mathbb{C} \setminus \{0\}$, $\lambda \geq 1$ and $r, \mu \in \mathbb{R}$, let $f \in \mathcal{A}$ be in the class $\mathcal{P}_\Sigma(\delta, \lambda, k, \gamma, \alpha, \beta, r)$. Then

$$|a_3 - \mu a_2^2| \leq \begin{cases} \frac{\delta |br|}{(3\lambda - 1) |\Phi_3^\gamma(k, \alpha, \beta)|} \\ \left(\left| \mu - 1 \right| \leq \frac{\left| \left[\delta \left((3\lambda - 1)\Phi_3^\gamma(k, \alpha, \beta) + (2\lambda(\lambda - 2) + 1)\Phi_2^{2\gamma}(k, \alpha, \beta) \right) b - p(2\lambda - 1)^2 \right] br^2 - qa(2\lambda - 1)^2 \right|}{\delta b^2 r^2 (3\lambda - 1) |\Phi_3^\gamma(k, \alpha, \beta)|} \right) \\ \frac{\delta^2 |br|^3 |\mu - 1|}{\left| \left[\delta \left((3\lambda - 1)\Phi_3^\gamma(k, \alpha, \beta) + (2\lambda(\lambda - 2) + 1)\Phi_2^{2\gamma}(k, \alpha, \beta) \right) b - p(2\lambda - 1)^2 \right] br^2 - qa(2\lambda - 1)^2 \right|} \\ \left(\left| \mu - 1 \right| \geq \frac{\left| \left[\delta \left((3\lambda - 1)\Phi_3^\gamma(k, \alpha, \beta) + (2\lambda(\lambda - 2) + 1)\Phi_2^{2\gamma}(k, \alpha, \beta) \right) b - p(2\lambda - 1)^2 \right] br^2 - qa(2\lambda - 1)^2 \right|}{\delta b^2 r^2 (3\lambda - 1) |\Phi_3^\gamma(k, \alpha, \beta)|} \right) \end{cases}.$$

Proof. It follows from (21) and (22) that

$$\begin{aligned} a_3 - \mu a_2^2 &= \frac{\delta h_2(r)(u_2 - v_2)}{2(3\lambda - 1)\Phi_3^\gamma(k, \alpha, \beta)} + (1 - \mu)a_2^2 \\ &= \frac{\delta h_2(r)(u_2 - v_2)}{2(3\lambda - 1)\Phi_3^\gamma(k, \alpha, \beta)} \\ &\quad + \frac{\delta^2 h_2^3(r)(u_2 + v_2)(1 - \mu)}{2 \left[\delta h_2^2(r) \left((3\lambda - 1)\Phi_3^\gamma(k, \alpha, \beta) + (2\lambda(\lambda - 2) + 1)\Phi_2^{2\gamma}(k, \alpha, \beta) \right) - h_3(r)(2\lambda - 1)^2 \right]} \\ &= \frac{h_2(x)}{2} \left[\left(\psi(\mu, r) + \frac{\delta}{(3\lambda - 1)\Phi_3^\gamma(k, \alpha, \beta)} \right) u_2 + \left(\psi(\mu, r) - \frac{\delta}{(3\lambda - 1)\Phi_3^\gamma(k, \alpha, \beta)} \right) v_2 \right], \end{aligned}$$

where

$$\psi(\mu, r) = \frac{\delta^2 h_2^2(r)(1 - \mu)}{\delta h_2^2(r) \left((3\lambda - 1)\Phi_3^\gamma(k, \alpha, \beta) + (2\lambda(\lambda - 2) + 1)\Phi_2^{2\gamma}(k, \alpha, \beta) \right) - h_3(r)(2\lambda - 1)^2}.$$

Thus, according to (3), we have

$$|a_3 - \mu a_2^2| \leq \begin{cases} \frac{\delta |br|}{(3\lambda - 1) |\Phi_3^\gamma(k, \alpha, \beta)|} \\ \left(0 \leq |\psi(\mu, r)| \leq \frac{\delta}{(3\lambda - 1) |\Phi_3^\gamma(k, \alpha, \beta)|} \right) \\ |br| \cdot |\psi(\mu, r)| \\ \left(|\psi(\mu, r)| \geq \frac{\delta}{(3\lambda - 1) |\Phi_3^\gamma(k, \alpha, \beta)|} \right), \end{cases}$$

which, after simple computation, yields

$$|a_3 - \mu a_2^2| \leq \begin{cases} \frac{\delta |br|}{(3\lambda - 1) |\Phi_3^\gamma(k, \alpha, \beta)|} \\ \left(|\mu - 1| \leq \frac{\left| \left[\delta \left((3\lambda - 1)\Phi_3^\gamma(k, \alpha, \beta) + (2\lambda(\lambda - 2) + 1)\Phi_2^{2\gamma}(k, \alpha, \beta) \right) b - p(2\lambda - 1)^2 \right] br^2 - qa(2\lambda - 1)^2 \right|}{\delta b^2 r^2 (3\lambda - 1) |\Phi_3^\gamma(k, \alpha, \beta)|} \right) \\ \frac{\delta^2 |br|^3 |\mu - 1|}{\left| \left[\delta \left((3\lambda - 1)\Phi_3^\gamma(k, \alpha, \beta) + (2\lambda(\lambda - 2) + 1)\Phi_2^{2\gamma}(k, \alpha, \beta) \right) b - p(2\lambda - 1)^2 \right] br^2 - qa(2\lambda - 1)^2 \right|} \\ \left(|\mu - 1| \geq \frac{\left| \left[\delta \left((3\lambda - 1)\Phi_3^\gamma(k, \alpha, \beta) + (2\lambda(\lambda - 2) + 1)\Phi_2^{2\gamma}(k, \alpha, \beta) \right) b - p(2\lambda - 1)^2 \right] br^2 - qa(2\lambda - 1)^2 \right|}{\delta b^2 r^2 (3\lambda - 1) |\Phi_3^\gamma(k, \alpha, \beta)|} \right). \end{cases}$$

We have thus completed the proof of Theorem 2. \square

3. Corollaries and Consequences

Our main results (Theorem 1 and Theorem 2) can be specialized to deduce a number of known or new results as their corollaries and consequences dealing with the initial Taylor-Maclaurin coefficient inequalities and the Fekete-Szegő inequalities. We choose to record here one example in which, by putting $\mu = 1$ in Theorem 2, we are led to the following corollary.

Corollary. For $\delta \in \mathbb{C} \setminus \{0\}$, $\lambda \geq 1$ and $r \in \mathbb{R}$, let $f \in \mathcal{A}$ be in the class $\mathcal{P}_\Sigma(\delta, \lambda, k, \gamma, \alpha, \beta, r)$. Then

$$|a_3 - a_2^2| \leq \frac{\delta |br|}{(3\lambda - 1) |\Phi_3^\gamma(k, \alpha, \beta)|}.$$

Remark 4. By taking particular values of the parameters $\delta, \lambda, \gamma, a, b, p$ and q , in our main results (Theorem 1 and Theorem 2), we can derive a number of known results. Some of these special cases are recorded below.

1. If we put $\gamma = 0$ and $\delta = \lambda = 1$ in Theorem 1 and Theorem 2, we have the results for the well-known class $S_\Sigma^*(r)$ of bi-starlike functions which was studied recently by Srivastava *et al.* [33].
2. If we put $\gamma = 0, \delta = a = 1, b = p = 2, q = -1$ and $r \rightarrow t$ in Theorem 1 and Theorem 2, we have the results for the class $\mathcal{LB}_\Sigma(t)$ of λ -Pseudo bi-starlike functions which was considered recently by Magesh and Bulut [20].
3. If we put $\gamma = 0, \delta = \lambda = a = 1, b = p = 2, q = -1$ and $r \rightarrow t$ in Theorem 1 and Theorem 2, we obtain the results for the class $S_\Sigma^*(t)$ of bi-starlike functions which was considered recently by Altinkaya and Yalçın [5].

4. Concluding Remarks and Observations

In our present investigation, we have introduced and studied a (presumably new) class $\mathcal{P}_\Sigma(\delta, \lambda, k, \gamma, \alpha, \beta, r)$ of normalized analytic and λ -pseudo-starlike bi-univalent functions in the open unit disk \mathbb{U} by applying the Horadam polynomials. This λ -pseudo-starlike bi-univalent function class is associated with a certain convolution operator. We have established the bounds for the initial Taylor-Maclaurin coefficients $|a_2|$ and $|a_3|$ of functions belonging to the λ -pseudo-starlike bi-univalent function class which we have defined in this article. Furthermore, we have successfully solved the Fekete-Szegő problem for functions in the same class $\mathcal{P}_\Sigma(\delta, \lambda, k, \gamma, \alpha, \beta, r)$ λ -pseudo-starlike bi-univalent functions. We have also considered a number of special cases and consequences for our main results (Theorem 1 and Theorem 2).

We conclude our investigation by remarking that, in order to motivate further researches on the subject-matter of this article, we have chosen to draw the attention of the interested readers toward a considerably large number of related recent publications on the subjects which we have discussed here. One of these directions for further researches should be motivated by a recently-published survey-cum-expository review article by Srivastava [29]. With this point in view, the attention of the interested reader is drawn toward the possibility of investigating the basic (or q -) extensions of the results which are presented in this paper. However, as already pointed out by Srivastava [29], their further extensions using the so-called (p, q) -calculus will be rather trivial and inconsequential variations of the suggested extensions which are based upon the classical q -calculus, the additional parameter p being redundant or superfluous (see, for details, [29, p. 340] and [32, pp. 1511–1512]). With a view to aiding the interested reader, we choose to cite several recent developments (see [17], [22], [23], [34], [44], [55] and [57]) on various usages of the basic (or q -) calculus in Geometric Function Theory of Complex Analysis.

Conflicts of Interest: The authors declares that they have no conflicts of interest.

References

- [1] C. Abirami, N. Magesh and J. Yamini, Initial bounds for certain classes of bi-univalent functions defined by Horadam polynomials, *Abstr. Appl. Anal.* **2020** (2020), Article ID 7391058, 1–8.
- [2] A. G. Al-Amoush, Certain subclasses of bi-univalent functions involving the Poisson distribution associated with Horadam polynomials, *Malaya J. Mat.* **7** (2019), 618–624.
- [3] J. W. Alexander, Functions which map the interior of the unit circle upon simple region, *Ann. Math.* **17** (1915), 12–22.
- [4] F. M. Al-Oboudi, On univalent functions defined by a generalized Sălăgean operator, *Internat. J. Math. Math. Sci.* **27** (2004), 1429–1436.
- [5] Ş. Altınkaya and S. Yalçın, On the Chebyshev polynomial coefficient problem of some subclasses of bi-univalent functions, *Gulf J. Math.* **5** (2017), 34–40.
- [6] K. O. Babalola, On λ -pseudo-starlike functions, *J. Class. Anal.* **3** (2013), 137–147.
- [7] S. D. Bernardi, Convex and starlike univalent functions, *Trans. Amer. Math. Soc.* **135** (1969), 429–446.
- [8] S. Bulut, Coefficient estimates for general subclasses of m -fold symmetric analytic bi-univalent functions, *Turkish J. Math.* **40** (2016), 1386–1397.
- [9] N. E. Cho and H. M. Srivastava, Argument estimates of certain analytic functions defined by a class of multiplier transformations, *Math. Comput. Model.* **37** (2003), 39–49.
- [10] P. L. Duren, *Univalent Functions*, Grundlehren der Mathematischen Wissenschaften, Band **259**, Springer-Verlag, New York, Berlin, Heidelberg and Tokyo, 1983.
- [11] M. Fekete and G. Szegő, Eine bemerkung uber ungerade schlichte funktionen, *J. London Math. Soc.* **2** (1933), 85–89.
- [12] H. Ö. Güney, G. Murugusundaramoorthy and J. Sokół, Subclasses of bi-univalent functions related to shell-like curves connected with Fibonacci numbers, *Acta Univ. Sapient. Math.* **10** (2018), 70–84.
- [13] A. F. Horadam, Jacobsthal representation polynomials, *Fibonacci Quart.* **35** (1997), 137–148.
- [14] A. F. Horadam and J. M. Mahon, Pell and Pell-Lucas polynomials, *Fibonacci Quart.* **23** (1985), 7–20.
- [15] T. Hörcüm and E. G. Koçer, On some properties of Horadam polynomials, *Internat. Math. Forum.* **4** (2009), 1243–1252.
- [16] I. B. Jung, Y. C. Kim and H. M. Srivastava, The Hardy space of analytic functions associated with certain one-parameter families of integral operators, *J. Math. Anal. Appl.* **176** (1993), 138–147.
- [17] B. Khan, H. M. Srivastava, M. Tahir, M. Darus, Q. Z. Ahmad and N. Khan, Applications of a certain q -integral operator to the subclasses of analytic and bi-univalent functions, *AIMS Mathematics.* **6** (2021), 1024–1039.
- [18] T. Koshy, *Fibonacci and Lucas Numbers with Applications*, A Wiley Interscience Publication, John Wiley and Sons, New York, Chichester, Brisbane and Toronto, 2001.
- [19] A. Lupas, A guide of Fibonacci and Lucas polynomials, *Octagon Math. Mag.* **7** (1999), 2–12.
- [20] N. Magesh and S. Bulut, Chebyshev polynomial coefficient estimates for a class of analytic bi-univalent functions related to pseudo-starlike functions, *Afrika Mat.* **29** (2018), 203–209.
- [21] S. S. Miller and P. T. Mocanu, *Differential Subordinations: Theory and Applications*, Series on Monographs and Textbooks in Pure and Applied Mathematics, Vol. **225**, Marcel Dekker Incorporated, New York and Basel, 2000.
- [22] M. Raza, H. M. Srivastava, M. Arif and K. Ahmad, Coefficient estimates for a certain family of analytic functions involving a q -derivative operator, *Ramanujan J.* **54** (2021), 501–519.
- [23] M. S. U. Rehman, Q. Z. Ahmad, H. M. Srivastava, N. Khan, M. Darus and B. Khan, Applications of higher-order q -derivatives to the subclass of q -starlike functions associated with the Janowski functions, *AIMS Mathematics.* **6** (2021), 1110–1125.
- [24] G. St. Sălăgean, Subclasses of univalent functions, *Lecture Notes in Mathematics*, Vol. **1013**, pp. 362–372, Springer Verlag, Berlin, Heidelberg and New York, 1983.
- [25] Y. J. Sim, O. S. Kwon, N. E. Cho and H. M. Srivastava, Bounds for the real parts and arguments of normalized analytic functions defined by the Srivastava-Attiya operator, *J. Comput. Anal. Appl.* **28** (2020), 628–645.
- [26] H. M. Srivastava, A new family of the λ -generalized Hurwitz-Lerch zeta functions with applications, *Appl. Math. Inform. Sci.* **8** (2014), 1485–1500.
- [27] H. M. Srivastava, Some general families of the Hurwitz-Lerch Zeta functions and their applications: Recent developments and directions for further researches, *Proc. Inst. Math. Mech. Nat. Acad. Sci. Azerbaijan.* **45** (2019), 234–269.
- [28] H. M. Srivastava, The Zeta and related functions: Recent developments, *J. Adv. Engrg. Comput.* **3** (2019), 329–354.
- [29] H. M. Srivastava, Operators of basic (or q -) calculus and fractional q -calculus and their applications in geometric function theory of complex analysis, *Iran. J. Sci. Technol. Trans. A: Sci.* **44** (2020), 327–344.
- [30] H. M. Srivastava, A survey of some recent developments on higher transcendental functions of analytic number theory and applied mathematics, *Symmetry* **13** (2021), Article ID 2294, 1–22.
- [31] H. M. Srivastava, An introductory overview of fractional-calculus operators based upon the Fox-Wright and related higher transcendental functions, *J. Adv. Engrg. Comput.* **5** (2021), 135–166.
- [32] H. M. Srivastava, Some parametric and argument variations of the operators of fractional calculus and related special functions and integral transformations, *J. Nonlinear Convex Anal.* **22** (2021), 1501–1520.
- [33] H. M. Srivastava, Ş. Altınkaya and S. Yalçın, Certain subclasses of bi-univalent functions associated with the Horadam polynomials, *Iran. J. Sci. Technol. Trans. A: Sci.* **43** (2019), 1873–1879.
- [34] H. M. Srivastava, M. Arif and M. Raza, Convolution properties of meromorphically harmonic functions defined by a generalized convolution q -derivative operator, *AIMS Mathematics.* **6** (2021), 5869–5885.
- [35] H. M. Srivastava and A. A. Attiya, An integral operator associated with the Hurwitz-Lerch zeta function and differential subordination, *Integral Transforms Spec. Funct.* **18** (2007), 207–216.
- [36] H. M. Srivastava, S. S. Eker, S. G. Hamidi and J. M. Jahangiri, Faber polynomial coefficient estimates for bi-univalent functions defined by the Tremblay fractional derivative operator, *Bull. Iran. Math. Soc.* **44** (2018), 149–157.

- [37] H. M. Srivastava and S. Gaboury, A new class of analytic functions defined by means of a generalization of the Srivastava-Attiya operator, *J. Inequal. Appl.* **2015** (2015), Article ID 39, 1–15.
- [38] H. M. Srivastava, S. Gaboury and F. Ghanim, A unified class of analytic functions involving a generalization of the Srivastava-Attiya operator, *Appl. Math. Comput.* **251** (2015), 35–45.
- [39] H. M. Srivastava, S. Gaboury and F. Ghanim, Coefficient estimates for some general subclasses of analytic and bi-univalent functions, *Afrika Mat.* **28** (2017), 693–706.
- [40] H. M. Srivastava, S. Gaboury and F. Ghanim, Coefficient estimates for a general subclass of analytic and bi-univalent functions of the Ma-Minda type, *Rev. Real Acad. Cienc. Exactas Fis. Natur. Ser. A Mat. (RACSAM)*. **112** (2018), 1157–1168.
- [41] H. M. Srivastava, S. Hussain, A. Raziq and M. Raza, The Fekete-Szegő functional for a subclass of analytic functions associated with quasi-subordination, *Carpathian J. Math.* **34** (2018), 103–113.
- [42] H. M. Srivastava, A. S. Juma and H. M. Zayed, Univalence conditions for an integral operator defined by a generalization of the Srivastava-Attiya operator, *Filomat*. **32** (2018), 2101–2114.
- [43] H. M. Srivastava, G. Kaur and G. Singh, Estimates of the fourth Hankel determinant for a class of analytic functions with bounded turnings involving cardioid domains, *J. Nonlinear Convex Anal.* **22** (2021), 511–526.
- [44] H. M. Srivastava, B. Khan, N. Khan, M. Tahir, S. Ahmad and Nasir Khan, Upper bound of the third Hankel determinant for a subclass of q -starlike functions associated with the q -exponential function, *Bull. Sci. Math.* **167** (2021), Article ID 102942, 1–16.
- [45] H. M. Srivastava, S. Khan, Q. Z. Ahmad, N. Khan and S. Hussain, The Faber polynomial expansion method and its application to the general coefficient problem for some subclasses of bi-univalent functions associated with a certain q -integral operator, *Stud. Univ. Babeş-Bolyai Math.* **63** (2018), 419–436.
- [46] H. M. Srivastava, A. K. Mishra and M. K. Das, The Fekete-Szegő problem for a subclass of close-to-convex functions, *Complex Variables Theory Appl.* **44** (2001), 145–163.
- [47] H. M. Srivastava, A. K. Mishra and P. Gochhayat, Certain subclasses of analytic and bi-univalent functions, *Appl. Math. Lett.* **23** (2010), 1188–1192.
- [48] H. M. Srivastava, A. Motamednezhad and E. A. Adegani, Faber polynomial coefficient estimates for bi-univalent functions defined by using differential subordination and a certain fractional derivative operator, *Mathematics*. **8** (2020), Article ID 172, 1–12.
- [49] H. M. Srivastava, A. Motamednezhad and S. Salehian, Coefficients of a comprehensive subclass of meromorphic bi-univalent functions associated with the Faber polynomial expansion, *Axioms*. **10** (2021), Article ID 27, 1–13.
- [50] H. M. Srivastava, G. Murugusundaramoorthy and S. M. El-Deeb, Faber polynomial coefficient estimates of bi-close-to-convex functions connected with the Borel distribution of the Mittag-Leffler type, *J. Nonlinear Var. Anal.* **5** (2021), 103–118.
- [51] H. M. Srivastava, A. Prajapati and P. Gochhayat, Third order differential subordination and differential superordination results for analytic functions involving the Srivastava-Attiya operator, *Appl. Math. Inform. Sci.* **12** (2018), 469–481.
- [52] H. M. Srivastava, N. Raza, E. S. A. AbuJarad, G. Srivastava and M. H. AbuJarad, Fekete-Szegő inequality for classes of (p, q) -starlike and (p, q) -convex functions, *Rev. Real Acad. Cienc. Exactas Fis. Natur. Ser. A Mat. (RACSAM)*. **113** (2019), 3563–3584.
- [53] H. M. Srivastava, F. M. Sakar and H. Ö. Güneş, Some general coefficient estimates for a new class of analytic and bi-univalent functions defined by a linear combination, *Filomat*. **32** (2018), 1313–1322.
- [54] H. M. Srivastava and A. K. Wanas, Initial Maclaurin coefficient bounds for new subclasses of analytic and m -fold symmetric bi-univalent functions defined by a linear combination, *Kyungpook Math. J.* **59** (2019), 493–503.
- [55] H. M. Srivastava, A. K. Wanas and H. Ö. Güneş, New families of bi-univalent functions associated with the Bazilevič functions and the λ -pseudo-starlike functions, *Iran. J. Sci. Technol. Trans. A: Sci.* **45** (2021), 1799–1804.
- [56] H. M. Srivastava, A. K. Wanas and G. Murugusundaramoorthy, A certain family of bi-univalent functions associated with the Pascal distribution series based upon the Horadam polynomials, *Surveys Math. Appl.* **16** (2021), 193–205.
- [57] H. M. Srivastava, A. K. Wanas and R. Srivastava, Applications of the q -Srivastava-Attiya operator involving a certain family of bi-univalent functions associated with the Horadam polynomials, *Symmetry* **13** (2021), Article ID 1230, 1–14.
- [58] S. R. Swamy, Inclusion properties of certain subclasses of analytic functions, *Internat. Math. Forum* **7** (2012), 1751–1760.
- [59] B. A. Uraleghaddi and C. Somanatha, Certain classes of univalent functions, in *Current Topics in Analytic Function Theory*, (Edited by H. M. Srivastava and S. Owa), pp. 371–374, World Scientific Publishing Company, Singapore, New Jersey, London and Hong Kong, 1992.
- [60] A. K. Wanas, New differential operator for holomorphic functions, *Earthline J. Math. Sci.* **2** (2019), 527–537.
- [61] A. K. Wanas, Bounds for initial Maclaurin coefficients for a new subclasses of analytic and m -fold symmetric bi-univalent functions, *TWMS J. App. Eng. Math.* **10** (2020), 305–311.
- [62] A. K. Wanas, Coefficient estimates of bi-starlike and bi-convex functions with respect to symmetrical points associated with the second kind Chebyshev polynomials, *Earthline J. Math. Sci.* **3** (2020), 191–198.
- [63] A. K. Wanas and A. L. Alina, Applications of Horadam polynomials on Bazilevič bi-univalent function satisfying subordinate conditions, *J. Phys.: Conf. Ser.* **1294** (2019), 1–6.
- [64] A. K. Wanas and A. H. Majeed, Chebyshev polynomial bounded for analytic and bi-univalent functions with respect to symmetric conjugate points, *Appl. Math. E-Notes* **19** (2019), 14–21.
- [65] A. K. Wanas and S. Yalçın, Initial coefficient estimates for a new subclasses of analytic and m -fold symmetric bi-univalent functions, *Malaya J. Mat.* **7** (2019), 472–476.