



Approximation by trigonometric polynomials and Faber-Laurent rational functions in grand Morrey spaces

Sadulla Z. Jafarov^a

^aDepartment of Mathematics and Science Education, Faculty of Education, Muş Alparslan University, 49250, Muş, Turkey; Institute of Mathematics and Mechanics, National Academy of Sciences of Azerbaijan, 9 B. Vahabzadeh str., AZ1141, Baku, Azerbaijan

Abstract. Let G be finite Jordan domain bounded a Dini smoth curve Γ in the complex plane \mathbb{C} . We investigate the approximation properties of the partial sums of the Fourier series and prove direct theorem for approximation by polynomials in the subspace of Morrey spaces associated with grand Lebesgue spaces. Also, approximation properties of the Faber-Laurent rational series expansions in spaces $L^{p,\lambda}(\Gamma)$ are studied. Direct theorems of approximation theory in grand Morrey-Smirnov classes, defined in domains with a Dini- smooth boundary, are proved.

1. Introduction and main results

Let \mathbb{T} denotes the interval $[0, 2\pi]$. Let $L^p(\mathbb{T})$, $1 \leq p < \infty$ be the Lebesgue space of all measurable 2π -periodic functions defined on \mathbb{T} such that

$$\|f\|_p := \left(\int_{\mathbb{T}} |f(x)|^p dx \right)^{\frac{1}{p}} < \infty.$$

The Morrey spaces $L_0^{p,\lambda}(\mathbb{T})$ for a given $0 \leq \lambda \leq 1$ and $p \geq 1$, we define as the set of functions $f \in L_{loc}^p(\mathbb{T})$ such that

$$\|f\|_{L_0^{p,\lambda}(\mathbb{T})} := \left\{ \sup_I \frac{1}{|I|^{1-\lambda}} \int_I |f(t)|^p dt \right\}^{\frac{1}{p}} < \infty,$$

where the supremum is taken over all intervals $I \subset [0, 2\pi]$. Note that $L_0^{p,\lambda}(\mathbb{T})$ becomes a Banach spaces, $\lambda = 1$ coincides with $L^p(\mathbb{T})$ and for $\lambda = 0$ with $L^\infty(\mathbb{T})$. If $0 \leq \lambda_1 \leq \lambda_2 \leq 1$, then $L_0^{p,\lambda_1}(\mathbb{T}) \subset L_0^{p,\lambda_2}(\mathbb{T})$. Also, if $f \in L_0^{p,\lambda}(\mathbb{T})$, then $f \in L^p(\mathbb{T})$ and hence $f \in L^1(\mathbb{T})$. The Morrey spaces, were introduced by C. B. Morrey in

2020 Mathematics Subject Classification. 30E10, 41A10, 41A17, 41A20, 41A50, 42A05, 42A10

Keywords. Morrey spaces associated with grand Lebesgue spaces; Modulus of smoothness; Direct theorem; Faber-Laurent rational functions; Dini-smooth curve; Best approximation.

Received: 26 September 2022; Accepted: 07 November 2022

Communicated by Dragan S. Djordjević

Email address: s.jafarov@alparslan.edu.tr (Sadulla Z. Jafarov)

1938. The properties of these spaces have been investigated intensively by various authors and together with Lebesgue spaces L^p play an important role in the theory of partial equations, especially in the study of local behavior of the solutions of elliptic differential equations and describe local regularity more precisely than Lebesgue spaces L^p . The properties of the Morrey spaces have been investigated by several authors (see, for example, [13], [22], [24], [35], [44], [45], [46], [51] and [56]).

We denote by $L^{p,\theta}(\mathbb{T})$, $\theta \geq 0$, the Lebesgue space of all measurable functions f on \mathbb{T} , that is, the space of all such functions for which

$$\|f\|_{L^{p,\theta}(\mathbb{T})} := \sup_{0 < \varepsilon < p-1} \left(\frac{\varepsilon^\theta}{2\pi} \int_{\mathbb{T}} |f(x)|^{p-\varepsilon} dx \right)^{\frac{1}{p-\varepsilon}} < \infty.$$

The space $L^{p,\theta}(\mathbb{T})$, $\theta > 0$ is called the *generalized grand Lebesgue space*. $L^{p,\theta}(\mathbb{T})$ is Banach function space, nonreflexive and nonseparable. Note that if $\theta = 0$ then $L^{p,\theta}(\mathbb{T})$ turns into the grand Lebesgue space $L^p(\mathbb{T})$. If $\theta = 1$ then we obtain grand Lebesgue space $L^p(\mathbb{T}) := L^{p,1}(\mathbb{T})$. The grand and generalized grand Lebesgue space were introduced in the works [27] and [23], respectively. Note that the space $L^p(\mathbb{T})$ is a rearrangement invariant Banach function space, but is not reflexive. We can write the following embeddings:

$$L^p(\mathbb{T}) \subset L^{p,1}(\mathbb{T}) \subset L^{p-\varepsilon}(\mathbb{T}), \quad 1 < p < \infty$$

If $\theta_1 < \theta_2$ then for $0 < \varepsilon < p - 1$ the embeddings

$$L^p(\mathbb{T}) \subset L^{p,\theta_1}(\mathbb{T}) \subset L^{p,\theta_2}(\mathbb{T}) \subset L^{p-\varepsilon}(\mathbb{T}), \quad 1 < p < \infty$$

hold. Note that the space $L^p(\mathbb{T})$ is not dense in $L^p(\mathbb{T})$ [18]. The informations about properties and applications of the grand Lebesgue spaces can be found in [18], [23], [27], [39], [53] and [54].

We denote by $L^{p,\lambda}(\mathbb{T})$, $1 < p < \infty$, $0 \leq \lambda < 1$ the Lebesgue space of all measurable functions f on \mathbb{T} , that is, the space of all such functions for which

$$\|f\|_{L^{p,\lambda}(\mathbb{T})} := \sup_{0 < \varepsilon < p-1} \left(\sup_I \frac{\varepsilon}{|I|^\lambda} \int_I |f(x)|^{p-\varepsilon} dx \right)^{\frac{1}{p-\varepsilon}} < \infty,$$

where the supremum is taken over all intervals $I \subset [0, 2\pi]$. The space $L^{p,\lambda}(\mathbb{T})$ is called *grand Morrey space*.

It is easy to verify that the following embeddings are hold:

$$L^{p,\lambda}(\mathbb{T}) \subset L^{p,\lambda}(\mathbb{T}) \subset L^{p-\varepsilon}(\mathbb{T}), \quad 0 < \varepsilon \leq p - 1.$$

Denote by $C^\infty(\mathbb{T})$ the set of all functions that are realized as the restriction to \mathbb{T} of elements in $C^\infty(\mathbb{R})$. We define $\widetilde{L}^{p,\lambda}(\mathbb{T})$ as the closure of $C^\infty(\mathbb{T})$ in $L^{p,\lambda}(\mathbb{T})$.

Let $W_r^{p,\lambda}(\mathbb{T})$ ($r = 1, 2, \dots$) (resp. $\widetilde{W}_r^{p,\lambda}(\mathbb{T})$ ($r = 1, 2, \dots$)) be the linear space of functions for which $f^{(r-1)}$ is absolutely continuous and $f^{(r)} \in L^{p,\lambda}(\mathbb{T})$ (resp. $\widetilde{L}^{p,\lambda}(\mathbb{T})$) be the linear space of functions for which $f^{(r-1)}$ is absolutely continuous and $f^{(r)} \in \widetilde{L}^{p,\lambda}(\mathbb{T})$.

Let G be a finite domain in the complex plane \mathbb{C} , bounded by the rectifiable Jordan curve Γ . Without loss of generality we assume $0 \in \text{Int } \Gamma$. Let $G^- = \text{Ext } \Gamma$. Let also $\mathbb{T} := \{w \in \mathbb{C} : |w| = 1\}$, $\mathbb{D} = \text{Int } \mathbb{T}$ and $\mathbb{D}^- = \text{Ext } \mathbb{T}$. We recall that if for a given analytic function f on G , there exists a sequence of rectifiable Jordan curves (Γ_n) in G tending to the boundary Γ in the sense that Γ_n eventually surrounds each compact subdomain of G such that

$$\int_{\Gamma_n} |f(z)|^p |dz| \leq M < \infty,$$

then we say that f belongs to the Smirnov class $E^p(G)$, $1 \leq p < \infty$. Each function $f \in E^p(G)$ has non-tangential limit almost everywhere (a.e.) on Γ and the boundary function belongs to $L^p(\Gamma)$.

We denote by φ the conformal mapping of G^- onto \mathbb{D}^- normalized by

$$\varphi(\infty) = \infty, \quad \lim_{z \rightarrow \infty} \frac{\varphi(z)}{z} > 0.$$

Let ψ be the inverse of φ . The function φ and ψ have continuous extensions to Γ and \mathbb{T} , their derivatives φ' and ψ' have definite non-tangential limit values on Γ and \mathbb{T} a.e., and they are integrable with respect to the Lebesgue measure on Γ and \mathbb{T} , respectively. It is known that $\varphi' \in E^1(G^-)$ and $\psi' \in E^1(\mathbb{D}^-)$. Note that the general information about Smirnov classes can be found in [19, pp. 168-185] and [26 pp. 438-453].

Let $\Gamma \subset \mathbb{C}$ be a rectifiable Jordan curve in the complex plane. We denote $\Gamma(t, r) = \Gamma \cap B(t, r)$, $t \in \Gamma, r > 0$, where $B(t, r) = \{z \in \mathbb{C} : |z - t| < r\}$. The Morrey spaces $L^{p,\lambda}(\Gamma)$ for a given $0 \leq \lambda \leq 1$ and $p \geq 1$, are defined as the set of functions $f \in L^p_{loc}(\Gamma)$ such that

$$\|f\|_{L^{p,\lambda}(\Gamma)} := \sup_{z \in \Gamma, 0 < r < L} r^{-\frac{\lambda}{p}} \|f\|_{L^p(\Gamma(t,r))} < \infty,$$

where L is the length of the curve Γ .

Note that $L^{p,0}(\Gamma) = L^p(\Gamma)$, and if $\lambda < 0$ or $\lambda > 1$, then $L^{p,\lambda}(\Gamma) = \Theta$, where Θ is the set of all functions equivalent to 0 on Γ .

We define also the Morrey–Smirnov classes $E^{p,\lambda}(G)$, $0 \leq \lambda \leq 1$, and $p \geq 1$, of analytic functions in G as

$$E^{p,\lambda}(G) := \{f \in E^1(G) : f \in L^{p,\lambda}(\Gamma)\}.$$

If we define the norm of $f \in E^{p,\lambda}(G)$, $0 \leq \lambda \leq 1$ and $p \geq 1$ by

$$\|f\|_{E^{p,\lambda}(G)} := \|f\|_{L^{p,\lambda}(\Gamma)},$$

then the class $E^{p,\lambda}(G)$, $0 \leq \lambda \leq 1$ and $p \geq 1$ will be a Banach space. Note that for $\lambda = 1$ the class $E^{p,\lambda}(G)$ coincides with the classical Smirnov class $E^p(G)$ and for $\lambda = 0$ with $E^\infty(G)$. If $G = D := \{z : |z| < 1\}$, then we have Morrey–Hardy space $H^{p,\lambda}(D) := E^{p,\lambda}(D)$ on the unit disk D .

Let $\Gamma \subset \mathbb{C}$ be a rectifiable curve and let ν be arc-length measure on Γ . Let $1 < p < \infty$ and let $0 \leq \lambda < 1$. We say that a measurable locally integrable function f on Γ belongs to the class $L^{p,\lambda}(\Gamma)$ if the following condition holds:

$$\|f\|_{L^{p,\lambda}(\Gamma)} := \sup_{0 < \varepsilon < p-1} \left(\sup_{\substack{z \in \Gamma \\ 0 < r < L}} \frac{\varepsilon}{\nu(B(z,r) \cap \Gamma)^\lambda} \int_{B(z,r) \cap \Gamma} |f(t)|^{p-\varepsilon} d\nu(t) \right)^{\frac{1}{p-\varepsilon}} < \infty.$$

We define also the grand Morrey–Smirnov classes $E^{p,\lambda}(G)$, $0 \leq \lambda < 1$, and $1 < p < \infty$, of analytic functions f in G as

$$E^{p,\lambda}(G) := \{f \in E^1(G) : f \in L^{p,\lambda}(\Gamma)\}.$$

We define the norm of $f \in E^{p,\lambda}(G)$, $0 \leq \lambda < 1$ and $1 < p < \infty$ by

$$\|f\|_{E^{p,\lambda}(G)} := \|f\|_{L^{p,\lambda}(\Gamma)}.$$

The closure of Smirnov class $E^p(G)$ in the space $E^{p,\lambda}(G)$, $0 \leq \lambda < 1$ and $1 < p < \infty$ we denote by $\widetilde{E}^{p,\lambda}(G)$, $1 < p < \infty$ and $0 \leq \lambda < 1$.

Let $1 < p < \infty$ and $0 \leq \lambda < 1$, and let

$$\Delta_t^r f(x) := \sum_{s=0}^r (-1)^{r+s+1} \binom{r}{s} f(x+st), \quad t > 0,$$

for a given $r \in \mathbb{N}$. For $f \in \widetilde{L}^{p,\lambda}(\mathbb{T})$, we define the operator

$$(v_h^r f)(x) := \frac{1}{h} \int_0^h |\Delta_t^r f(x)| dt.$$

The function

$$\Omega_r(f, \delta)_{p,\lambda} := \sup_{0 < h \leq \delta} \left\| (v_h^r f)(x) \right\|_{L^{p,\lambda}(\mathbb{T})}, \quad \delta > 0,$$

is called the *r*th mean modulus of $f \in \widetilde{L}^{p,\lambda}(\mathbb{T})$, $1 < p < \infty$, $0 \leq \lambda < 1$.

It can easily be shown that $\Omega_r(f, \cdot)_{p,\lambda}$ is a continuous, nonnegative and nondecreasing function satisfying the conditions

$$\lim_{\delta \rightarrow 0} \Omega_r(f, \delta)_{p,\lambda} = 0, \quad \Omega_r(f+g, \delta)_{p,\lambda} \leq \Omega_r(f, \delta)_{p,\lambda} + \Omega_r(g, \delta)_{p,\lambda}, \quad \delta > 0$$

for $f, g \in L^{p,\lambda}(\mathbb{T})$.

Let $f \in L^1(\mathbb{T})$ and let \widetilde{f} be its conjugate function, with Fourier series

$$f(x) \sim \sum_{k=1}^{\infty} (a_k \cos kx + b_k \sin kx), \quad \widetilde{f}(x) \sim \sum_{k=1}^{\infty} (a_k \sin kx - b_k \cos kx).$$

Let $S_n(\cdot, f)$ ($n = 1, 2, \dots$) be the *n*th partial sum of the Fourier series of $f \in L^1(\mathbb{T})$, i.e.

$$S_n(x, f) = \sum_{k=1}^n (a_k \cos kx + b_k \sin kx).$$

Using the method of proof of [17, Theorem 2.1 and Proposition 2.2] one can show that

$$\|S_n(\cdot, f)\|_{L^{p,\lambda}(\mathbb{T})} \leq c_1 \|f\|_{L^{p,\lambda}(\mathbb{T})}, \quad \|\widetilde{f}\|_{L^{p,\lambda}(\mathbb{T})} \leq c_2 \|f\|_{L^{p,\lambda}(\mathbb{T})},$$

as a corollary we obtain

$$\|f - S_n(\cdot, f)\|_{L^{p,\lambda}(\mathbb{T})} \leq c_3 E_n(f)_{p,\lambda}, \quad E_n(\widetilde{f})_{p,\lambda} \leq c_4 E_n(f)_{p,\lambda}. \tag{1}$$

Let $f \in \widetilde{L}^{p,\lambda}(\mathbb{T})$ and $r \in \mathbb{N}$. We define *K*-functional as

$$K_r(f, \delta)_{p,\lambda} := \inf \left\{ \|f - \psi\|_{L^{p,\lambda}} + \delta^r \|\psi^{(r)}\|_{L^{p,\lambda}(\mathbb{T})} : \psi \in \widetilde{W}_r^{p,\lambda}(\mathbb{T}), \delta > 0 \right\}.$$

We denote by \prod_n the class of trigonometric polynomials of degree not exceeding $n \in \mathbb{N}$. The best approximation of $f \in \widetilde{L}^{p,\lambda}$, $1 < p < \infty$, $0 \leq \lambda < 1$ in the class \prod_n is defined by

$$E_n(f)_{p,\lambda} := \inf \left\{ \|f - T_n\|_{L^{p,\lambda}(\mathbb{T})} : T_n \in \prod_n \right\}.$$

We denote by $w = \phi(z)$ the conformal mapping of G^- onto domain $\mathbb{D} := \{w \in \mathbb{C} : |w| > 1\}$ normalized by the conditions

$$\phi(\infty) = \infty, \lim_{z \rightarrow \infty} \frac{\phi(z)}{z} > 0$$

and let ψ be the inverse mapping of ϕ .

We denote by $w = \phi_1(z)$ the conformal mapping of G onto the domain $\mathbb{D}^- := \{w \in \mathbb{C} : |w| > 1\}$ normalized by the conditions

$$\phi_1(0) = \infty, \lim_{z \rightarrow 0} (z \cdot \phi_1(z)) > 0,$$

and let ψ_1 be the inverse mapping of ϕ_1 .

The functions ψ and ψ_1 have in some deleted neighborhood of the point $w = \infty$ the following representations

$$\psi(w) = \gamma w + \gamma_0 + \frac{\gamma_1}{w} + \frac{\gamma_2}{w^2} + \dots, \gamma > 0,$$

and

$$\psi_1(w) = \frac{\alpha_1}{w} + \frac{\alpha_2}{w^2} + \dots + \frac{\alpha_k}{w^k} + \dots, \alpha_1 > 0.$$

The following expansions hold [14], [19] and [57] :

$$\frac{\psi'(w)}{\psi(w) - z} = \sum_{k=0}^{\infty} \frac{\Phi_k(z)}{w^{k+1}}, \quad z \in G \text{ and } w \in \mathbb{D}^-, \tag{2}$$

and

$$\frac{\psi_1'(w)}{\psi_1(w) - z} = \sum_{k=0}^{\infty} -\frac{F_k\left(\frac{1}{z}\right)}{w^{k+1}}, \quad z \in G^- \text{ and } w \in \mathbb{D}^-, \tag{3}$$

where $\Phi_k(z)$ and $F_k\left(\frac{1}{z}\right)$ are the Faber polynomials of degree k with respect to z and $\frac{1}{z}$ for the continuums \overline{G} and $\overline{\mathbb{C}} \setminus G$, respectively. Also, for the Faber polynomials $\Phi_k(z)$ and rational functions $F_k\left(\frac{1}{z}\right)$ the integral representations

$$\Phi_k(z) = [\phi(z)]^k + \frac{1}{2\pi i} \int_{\Gamma} \frac{[\phi(\zeta)]^k}{\zeta - z} d\zeta, \quad k = 0, 1, 2, \dots, z \in G, \tag{4}$$

$$F_k\left(\frac{1}{z}\right) = [\phi_1(z)]^k - \frac{1}{2\pi i} \int_{\Gamma} \frac{[\phi_1(\zeta)]^k}{\zeta - z} d\zeta, \quad k = 0, 1, 2, \dots, z \in G \tag{5}$$

hold [14], [57].

Let also χ be a continuous function on 2π . Its modulus of continuity is defined by

$$\omega(t, \chi) := \sup_{t_1, t_2 \in [0, 2\pi], |t_1 - t_2| < t} |\chi(t_1) - \chi(t_2)|, \quad t \geq 0.$$

The curve Γ is called Dini-smooth curve if it has the parametrization

$$\Gamma : \chi(t), \quad 0 \leq t \leq 2\pi,$$

such that $\chi'(t)$ satisfied the Dini-continuous, i.e.

$$\int_0^\pi \frac{\omega(t, \chi')}{t} dt < \infty$$

and

$$\chi'(t) \neq 0$$

[50, p.48].

Let $f \in L_1(\Gamma)$. Then the functions f^+ and f^- defined by

$$f^+(z) = \frac{1}{2\pi i} \int_\Gamma \frac{f(\zeta)}{\zeta - z} d\zeta = \frac{1}{2\pi i} \int_{\mathbb{T}} \frac{f(\psi(w)) \psi'(w)}{\psi(w) - z} dw, \quad z \in G \tag{6}$$

and

$$f^-(z) = \frac{1}{2\pi i} \int_\Gamma \frac{f(\zeta)}{\zeta - z} d\zeta = \frac{1}{2\pi i} \int_{\mathbb{T}} \frac{f(\psi_1(w)) \psi_1'(w)}{\psi_1(w) - z} dw, \quad z \in G^- \tag{7}$$

are analytic in G and G^- , respectively, and $f^-(\infty) = 0$. Thus the limit

$$S_\Gamma(f)(z) := \lim_{\varepsilon \rightarrow \infty} \frac{1}{2\pi i} \int_{\Gamma \cap \{\zeta: |\zeta - z| > \varepsilon\}} \frac{f(\zeta)}{\zeta - z} d\zeta$$

exists and is finite for almost all $z \in \Gamma$.

The quantity $S_\Gamma(f)(z)$ is called the *Cauchy singular integral* of f at $z \in \Gamma$.

According to the Privalov theorem [26, p.431] if one of the functions f^+ or f^- has the non-tangential limits a.e. on Γ , then $S_\Gamma(f)(z)$ exists a.e. on Γ and also the other one has the non-tangential limits a.e. on Γ . Conversely, if $S_\Gamma(f)(z)$ exists a.e. on Γ , then the functions $f^+(z)$ and $f^-(z)$ have non-tangential limits a.e. on Γ . In both cases, the formulas

$$f^+(z) = S_\Gamma(f)(z) + \frac{1}{2}f(z), \quad f^-(z) = S_\Gamma(f)(z) - \frac{1}{2}f(z) \tag{8}$$

and hence

$$f(z) = f^+(z) - f^-(z) \tag{9}$$

holds a.e. on Γ . From the results in [47], it follows that if Γ is a Dini-smooth curve S_Γ is bounded on $L^{p,\lambda}(\Gamma)$. Note that some properties of the Cauchy singular integral in the different spaces were investigated in [12], [17], [20], [24], [38], [40], [41], [43] and [47].

Let $f \in L^{p(\cdot),\lambda(\cdot)}(\Gamma)$. Using (3), (4), (7), (8) and (9) we can associate Faber- Laurent series

$$f(z) \sim \sum_{k=0}^\infty a_k \Phi_k(z) + \sum_{k=1}^\infty b_k F_k\left(\frac{1}{z}\right),$$

where the coefficients a_k and b_k are defined by

$$a_k := \frac{1}{2\pi i} \int_{\mathbb{T}} \frac{f[\psi(w)]}{w^{k+1}} d\omega, \quad k = 0, 1, 2, \dots$$

and

$$b_k := \frac{1}{2\pi i} \int_{\mathbb{T}} \frac{f[\psi_1(w)]}{w^{k+1}} dw, \quad k = 0, 1, 2, \dots$$

The coefficients a_k and b_k are said to be the Faber- Laurent coefficients of f .

If Γ is a Dini-smooth curve, then from the results in [59], it follows that

$$\left. \begin{aligned} 0 < c_5 < |\phi'(w)| < c_6 < \infty, \quad 0 < c_7 < |\phi'_1(w)| < c_8 < \infty; \\ 0 < c_9 < |\psi'(w)| < c_{10} < \infty, \quad 0 < c_{11} < |\psi'_1(w)| < c_{12} < \infty \end{aligned} \right\} \tag{10}$$

where the constants c_5, c_6, c_7, c_8 and $c_9, c_{10}, c_{11}, c_{12}$ are independent of $z \in \bar{G}^-$ and $|w| \geq 1$, respectively.

Let Γ be a Dini-smooth curve and let $f_0(w) := f[\psi(w)]$ for $f \in L^{p,\lambda}(\Gamma)$ and let $f_1(w) := f[\psi_1(w)]$ for $f \in L^{p,\lambda}(\Gamma)$. Then using (10) and the method, applied for the proof of the similar result in [33, Lemma 1], we obtain $f_0 \in L^{p,\lambda}(\mathbb{T})$ and $f_1 \in L^{p,\lambda}(\mathbb{T})$.

Moreover, $f_0^-(\infty) = f_1^-(\infty) = 0$ and by (10)

$$f_0(w) = f_0^+(w) - f_0^-(w), \quad f_1(w) = f_1^+(w) - f_1^-(w) \tag{11}$$

a.e. on \mathbb{T} .

We set

$$R_n(f, z) := \sum_{k=0}^n a_k \Phi_k(z) + \sum_{k=0}^n b_k F_k\left(\frac{1}{z}\right).$$

The rational function $R_n(f, z)$ is called the Faber-Laurent rational function of degree n of f .

Note that, in this paper we study the problems of approximation theory in space $\widetilde{L}^{p,\lambda}(\mathbb{T})$, the closure of the set of the trigonometric polynomials in $L^{p,\lambda}(\mathbb{T})$, with $1 < p < \infty$ and $0 \leq \lambda < 1$.

The problems of approximation of the functions in classical Morrey spaces and variable exponent Morrey spaces were investigated in [10], [11], [12], [14], [15], [21], [26], [27]. In this work we investigate the approximation properties of the partial sums of the Fourier series and some direct theorems for approximation by trigonometric polynomials in the subspace grand Morrey spaces. Similar results in different spaces have been investigated by several authors (see. for example, [1]-[4], [8]-[11], [15], [16], [15]-[17], [25], [30]-[35], [42]). Also, approximation of the functions by Faber-Laurent rational functions in the grand Morrey spaces defined on the Dini-smooth curve are investigated.. Direct theorems of approximation theory in grand Morrey- Smirnov classes, defined in domains with a Dini-smooth boundary, are proved. Similar problems of approximation of the functions by Faber polynomials and Faber- Laurent rational functions in different spaces were studied in [5]-[7], [28]-[33], [36], [37], [49], [58], [60].

Our main results are as follows.

Theorem 1.1. *Let $1 < p < \infty$, $0 \leq \lambda < 1$ and $r \in \mathbb{N}$. Then for every $f \in \widetilde{W}_r^{p,\lambda}(\mathbb{T})$ the inequality*

$$E_n(f)_{p,\lambda} \leq \frac{c_{13}}{n^r} E_n(f^{(r)})_{p,\lambda} \tag{12}$$

holds with a constant $c_{13} > 0$ independent of n .

From Theorem 1.1 we obtain the following Corollary.

Corollary 1.1. *Let $1 < p < \infty$, $0 \leq \lambda < 1$ and $r \in \mathbb{N}$. If $f \in \widetilde{W}_r^{p,\lambda}(\mathbb{T})$ then*

$$E_n(f)_{p,\lambda} \leq \frac{c_{14}}{n^r} \|f^{(r)}\|_{L^{p,\lambda}(\mathbb{T})}$$

with a constant $c_{14} > 0$, independent of n .

Theorem 1.2. *Let $1 < p < \infty$, $0 \leq \lambda < 1$ and $r \in \mathbb{N}$. Then for every $f \in \widetilde{L}^{p,\lambda}(\mathbb{T})$ the inequality*

$$E_n(f)_{p,\lambda} \leq c_{15} \Omega_r(f, \delta)_{p,\lambda}$$

holds with a constant $c_{15} > 0$, independent of n .

Theorem 1.3. Let Γ be a Dini-smooth curve. If $f \in L^{p,\lambda}(\Gamma)$, $1 < p < \infty$, $0 \leq \lambda < 1$, then for every natural number n there are a constant $c_{16} > 0$ and rational function

$$R_n(z, f) := \sum_{k=-n}^n a_k^{(n)} z^k$$

such that

$$\|f - R_n(\cdot, f)\|_{L^{p,\lambda}(\Gamma)} \leq c_{16} \left\{ \Omega_r(f_0, 1/n)_{p,\lambda} + \Omega_r(f_1, 1/n)_{p,\lambda} \right\},$$

where $R_n(\cdot, f)$ is the n -th partial sum of the Faber-Laurent series of f .

Theorem 1.4. Let Γ be a Dini-smooth curve. If $f \in \widetilde{E}^{p,\lambda}(G)$, $1 < p < \infty$, $0 \leq \lambda < 1$, then for every natural number n the inequality

$$\left\| f - \sum_{k=0}^n a_k \Phi_k(z) \right\|_{L^{p,\lambda}(\Gamma)} \leq c_{17} \Omega_r(f_0, 1/n)_{p,\lambda} \tag{13}$$

holds with a constant $c_{17} > 0$ independent of n .

Note that the order of polynomial approximation in $E^p(G)$, $p \geq 1$, has been investigated by several authors. In [58] Walsh and Rusel gave results when Γ is an analytic curve. When Γ is a Dini-smooth curve direct and inverse theorems were proved by S. Y. Alper [5], These results were later extended to domains with regular boundary for $p > 1$ by V.M. Kokilashvili [42] and for $p \geq 1$ by J. E. Andersson [6]. The approximation properties of the p -Faber series expansions in the ω -weighted Smirnov class $E^p(G, \omega)$ of analytic functions in G whose boundary is a regular Jordan curve are investigated in [28].

Theorem 1.5. Let Γ be a Dini-smooth curve. $f \in \widetilde{E}^{p,\lambda}(G^-)$, then for every natural number n the inequality

$$\left\| f - f(\infty) - \sum_{k=0}^n -b_k F_k\left(\frac{1}{z}\right) \right\|_{L^{p,\lambda}(\Gamma)} \leq c_{18} \Omega(f_1, 1/n)_{p,1,\lambda} \tag{14}$$

holds, with a constant $c_{18} > 0$, independent of n .

2. Auxiliary results

In the proof of Theorem 1.2 we use the following theorem.

Theorem 2.1. Let $1 < p < \infty$, $0 \leq \lambda < 1$ and $r \in \mathbb{N}$. Then for every $f \in \widetilde{L}^{p,\lambda}(\mathbb{T})$ the inequality

$$K_r(f, \delta) \leq c_{19} \Omega_r(f, \delta)_{p,\lambda}$$

holds, where the constant c_{19} independent of δ .

Proof of Theorem 2.1. Let $r \geq 1$ and $\delta > 0$. We define the function

$$f_{r,\delta}(x) := \frac{2}{\delta} \int_{\delta/2}^{\delta} \left\{ \frac{1}{h^r} \int_0^h \dots \int_0^h \sum_{s=0}^{r-1} (-1)^{r+s+1} \binom{r}{s} \right. \\ \left. \times f\left(x + \frac{r-s}{r}(t_1 + \dots + t_r)\right) dt_1 \dots dt_r \right\} dh \tag{15}$$

It is clear that

$$\Delta_{(t_1+\dots+t_r)/r}^r f(x) \\ = \sum_{s=0}^{r-1} (-1)^{r+s+1} \binom{r}{s} f\left(x + \frac{r-s}{r}(t_1 + \dots + t_r)\right) - f(x). \tag{16}$$

Use of (15) and (16) we obtain

$$\begin{aligned}
 & \|f_{r,\delta}(\cdot) - f(\cdot)\|_{L^{p,\lambda}(\mathbb{T})} \\
 & \leq \frac{2}{\delta} \int_{\delta/2}^{\delta} \left\| \frac{1}{h^r} \int_0^h \dots \int_0^h \Delta_{(t_1+\dots+t_r)/r}^r f(\cdot) dt_1 \dots dt_r \right\|_{L^{p,\lambda}(\mathbb{T})} dh \\
 & = \frac{2}{\delta} \int_{\delta/2}^{\delta} \sup_{\frac{\delta}{2} \leq h \leq \delta} \left\| \frac{1}{h^r} \int_0^h \dots \int_0^h \Delta_{(t_1+\dots+t_r)/r}^r f(\cdot) dt_1 \dots dt_r \right\|_{L^{p,\lambda}(\mathbb{T})} dh \\
 & = \sup_{\frac{\delta}{2} \leq h \leq \delta} \left\| \frac{1}{h^r} \int_0^h \dots \int_0^h \Delta_{(t_1+\dots+t_r)/r}^r f(\cdot) dt_1 \dots dt_r \right\|_{L^{p,\lambda}(\mathbb{T})} \\
 & = \sup_{\frac{\delta}{2} \leq h \leq \delta} \left\| \frac{1}{h^r} \int_0^h \dots \int_0^h \left(\int_{t_2+\dots+t_r}^{t_2+\dots+t_r+h} |\Delta_{t/r}^r f(\cdot) dt| \right) dt_2 \dots dt_r \right\|_{L^{p,\lambda}(\mathbb{T})} \\
 & \leq c_{20} \sup_{\frac{\delta}{2} \leq h \leq \delta} \left\| \frac{1}{h^{r-1}} \int_0^h \dots \int_0^h \left(\int_{0_r}^{rh} |\Delta_{t/r}^r f(\cdot) dt| \right) dt_2 \dots dt_r \right\|_{L^{p,\lambda}(\mathbb{T})} \\
 & \leq \dots \leq c_{21} \sup_{\frac{\delta}{2} \leq h \leq \delta} \left\| \frac{1}{h} \int_{0_r}^{rh} |\Delta_{t/r}^r f(\cdot) dt| \right\|_{L^{p,\lambda}(\mathbb{T})} \\
 & = c_{22} r \sup_{0 \leq h \leq \delta} \left\| \frac{1}{h} \int_{0_r}^h \Delta_t^r f(\cdot) dt \right\|_{L^{p,\lambda}(\mathbb{T})} = c_{23} \Omega_r(f, \delta)_{p,\lambda} \tag{17}
 \end{aligned}$$

It is clear that for the function $f_{r,\delta}^{(r)}(\cdot)$ the equality

$$f_{r,\delta}^{(r)}(x) = \frac{2}{\delta} \int_{\delta/2}^{\delta} \left(\frac{1}{h^r} \sum_{s=0}^{r-1} (-1)^{r+s} \binom{r}{s} \left(\frac{r}{r-s}\right)^r \Delta_{(r-s)h/r} f(x) \right) dh \tag{18}$$

holds. Then from (18), we have

$$\begin{aligned}
 & \left| f_{r,\delta}^{(r)}(x) \right| \\
 & \leq \frac{2}{\delta} \int_{\delta/2}^{\delta} \sum_{s=0}^{r-1} \binom{r}{s} \left(\frac{r}{r-s}\right)^r |\Delta_{(r-s)h/r} f(x)| dh \\
 & \leq 2^{r+1} \delta^{-r} \sum_{s=0}^{r-1} \binom{r}{s} \left(\frac{r}{r-s}\right)^r \frac{1}{\delta} \int_0^{\delta} |\Delta_{(r-s)h/r} f(x)| dh. \tag{19}
 \end{aligned}$$

From (19), we get:

$$\begin{aligned}
 & \left\| f_{r,\delta}^{(r)} \right\|_{L^{p,\theta,\lambda}(\mathbb{T})} \\
 & \leq 2^{r+1} \delta^{-r} \sum_{s=0}^{r-1} \binom{r}{s} \left(\frac{r}{r-s} \right)^r \left\| \frac{1}{\delta} \int_0^\delta |\Delta_{(r-s)h/r} f(\cdot)| dh \right\|_{L^{p,\theta,\lambda}(\mathbb{T})} \\
 & = 2^{r+1} \delta^{-r} \sum_{s=0}^{r-1} \binom{r}{s} \left(\frac{r}{r-s} \right)^r \left\| \frac{1}{(r-s)\delta/r} \int_0^{(r-s)\delta/r} |\Delta_t^r f(x)| dt \right\|_{L^{p,\theta,\lambda}(\mathbb{T})} \\
 & \leq 2^{r+1} \delta^{-r} \sum_{s=0}^{r-1} \binom{r}{s} \left(\frac{r}{r-s} \right)^r \Omega_r(f, \delta)_{p,\theta,\lambda} \\
 & = 2^{2r} \delta^{-r} \Omega_r(f, \delta)_{p,\theta,\lambda}. \tag{20}
 \end{aligned}$$

Taking into account the relations (17) and (20) we have

$$\begin{aligned}
 & K_r(f, \delta)_{p,\lambda} \\
 & \leq \|f_{r,\delta} - f\|_{L^{p,\lambda}(\mathbb{T})} + \delta^r \|f_{r,\delta}^{(r)}\|_{L^{p,\lambda}(\mathbb{T})} \\
 & \leq 2^{r-1} r \Omega_r(f, \delta)_{p,\lambda} + 2^{2r} \delta^{-r} \Omega_r(f, \delta)_{p,\lambda} \leq c_{24} \Omega_r(f, \delta)_{p,\lambda}.
 \end{aligned}$$

The proof of Theorem 2.1 is completed.

In the proof of Theorem 1.3 we use the following Lemmas.

Using Theorem 1.2 and the method applied for the proof of a similar result in [14] we can prove the following Lemma:

Lemma 2.1. *Let $g \in E^{p,\lambda}(\mathbb{D})$, $1 < p < \infty$, $0 \leq \lambda < 1$. If $\sum_{k=0}^n d_k(g)w^k$ is the n th partial sum of the Taylor series of g at the origin, then*

$$\left\| g(w) - \sum_{k=0}^n d_k w^k \right\|_{L^{p,\lambda}(\mathbb{T})} \leq c_{25} \Omega\left(g, \frac{1}{n}\right)_{p,\lambda}, \forall n \in \mathbb{N},$$

holds with some constant $c_{25} > 0$, independent of n .

Lemma 2.2. *Let $g \in L^{p(\cdot),\lambda(\cdot)}(\mathbb{T})$, $1 < p < \infty$, $0 \leq \lambda < 1$. Then the inequality*

$$\Omega(g^+, \cdot)_{p(\cdot),\lambda(\cdot)} \leq c_{26} \Omega(g, \cdot)_{p(\cdot),\lambda(\cdot)} \tag{21}$$

holds, with some constant $c_{26} > 0$, independent of n .

Proof of Lemma 2.2. It is clear that the equality

$$g^+ = S_{\mathbb{T}}(g) + \frac{1}{2}g \tag{22}$$

holds a.e. on \mathbb{T} . Using the method of proof of [14, Lemma 3.3] (see also, [33, Lemma 2]) and the boundedness of the singular operator $S_{\mathbb{T}}(g)$ in $L^{p,\lambda}(\mathbb{T})$ [47] we can prove that

$$\Omega_r(S_{\mathbb{T}}(g) \cdot)_{p,\lambda} \leq c_{27} \Omega_r(g, \cdot)_{p,\lambda}. \tag{23}$$

Then using the subadditivity of the modulus of smoothness $\Omega_r(g^+, \cdot)_{p,\lambda}$, (22) and (23) we obtain inequality (21) of Lemma 2.1.

3. Proof of the main results

Proof of Theorem 1.1. Let $f \in \widetilde{W}^{p,\lambda}(\mathbb{T})$ and let $\sum_{k=0}^{\infty} (a_k \cos kx) + b_k \sin kx$ be the Fourier series of f and

$$S_n(x, f) = \sum_{k=0}^n (a_k \cos kx) + b_k \sin kx$$

be its n th partial sum. Note that for the conjugate function \widetilde{f} the Fourier expansion

$$S_n(x, f) = \sum_{k=1}^{\infty} (a_k \sin kx) - b_k \cos kx$$

holds. We set

$$B_k(x, f) := a_k \cos kx + b_k \sin kx.$$

It is clear that

$$f(x) = \sum_{k=0}^{\infty} B_k(x, f) \tag{24}$$

Note that, for $k = 1, 2, \dots$, we can write the following equalities:

$$\begin{aligned} B_k(x, f) & : = a_k \cos kx + b_k \sin kx \\ & = a_k \cos k \left(x + \frac{r\pi}{2k} - \frac{r\pi}{2k} \right) + b_k \sin k \left(x + \frac{r\pi}{2k} - \frac{r\pi}{2k} \right) \\ & = a_k \cos \left(kx + \frac{r\pi}{2} - \frac{r\pi}{2} \right) + b_k \sin \left(kx + \frac{r\pi}{2} - \frac{r\pi}{2} \right) \\ & = a_k \left[\cos \left(kx + \frac{r\pi}{2} \right) \cos \frac{r\pi}{2} - \sin \left(kx + \frac{r\pi}{2} \right) \sin \frac{r\pi}{2} \right] \\ & \quad + b_k \left[\sin \left(kx + \frac{r\pi}{2} \right) \cos \frac{r\pi}{2} - \cos \left(kx + \frac{r\pi}{2} \right) \sin \frac{r\pi}{2} \right] \\ & = \cos \frac{r\pi}{2} \left[a_k \cos k \left(x + \frac{r\pi}{2} \right) + b_k \sin k \left(x + \frac{r\pi}{2} \right) \right] \\ & \quad + \sin \frac{r\pi}{2} \left[a_k \sin k \left(x + \frac{r\pi}{2} \right) - b_k \cos k \left(x + \frac{r\pi}{2} \right) \right] \\ & = B_k \left(x + \frac{r\pi}{2k}, f \right) \cos \frac{r\pi}{2} + B_k \left(x + \frac{r\pi}{2k}, \widetilde{f} \right) \sin \frac{r\pi}{2}, \end{aligned} \tag{25}$$

$$B_k(x, f^{(r)}) = k^r B_k \left(x + \frac{r\pi}{2k}, f \right). \tag{26}$$

From (25) and (26), we obtain:

$$\begin{aligned}
 \sum_{k=0}^{\infty} B_k(x, f) &= B_0(x, f) + \cos \frac{r\pi}{2} \sum_{k=1}^{\infty} B_k\left(x + \frac{r\pi}{2k}, f\right) \\
 &\quad + \sin \frac{r\pi}{2} \sum_{k=1}^{\infty} B_k\left(x + \frac{r\pi}{2k}, \widetilde{f}\right) \\
 &= B_0(x, f) + \cos \frac{r\pi}{2} \sum_{k=1}^{\infty} \frac{1}{r^k} r^k B_k\left(x + \frac{r\pi}{2k}, f\right) \\
 &\quad + \sin \frac{r\pi}{2} \sum_{k=1}^{\infty} \frac{1}{r^k} r^k B_k\left(x + \frac{r\pi}{2k}, \widetilde{f}\right) \\
 &= B_0(x, f) + \cos \frac{r\pi}{2} \sum_{k=1}^{\infty} \frac{1}{r^k} B_k(x, f^{(r)}) \\
 &\quad + \sin \frac{r\pi}{2} \sum_{k=1}^{\infty} \frac{1}{r^k} B_k(x, \widetilde{f}^{(r)}). \tag{27}
 \end{aligned}$$

Then, from equalities, (24) and (27), we have:

$$\begin{aligned}
 f(x) - S_n(x, f) &= \sum_{k=n+1}^{\infty} B_k(x, f) \\
 &= \cos \frac{r\pi}{2} \sum_{k=n+1}^{\infty} \frac{1}{r^k} B_k(x, f^{(r)}) \\
 &\quad + \sin \frac{r\pi}{2} \sum_{k=n+1}^{\infty} \frac{1}{r^k} B_k(x, \widetilde{f}^{(r)}). \tag{28}
 \end{aligned}$$

On the other hand the following equalities hold:

$$\begin{aligned}
 \sum_{k=n+1}^{\infty} \frac{1}{k^r} B_k(\alpha, f^{(r)}) &= \sum_{k=n+1}^{\infty} \frac{1}{k^r} [S_k(\alpha, f^{(r)}) - S_{k-1}(\alpha, f^{(r)})] \\
 &= \sum_{k=n+1}^n \frac{1}{k^r} \{ [S_k(\alpha, f^{(r)}) - f^{(r)}(\alpha)] - [S_{k-1}(\alpha, f^{(r)}) - f^{(r)}(\alpha)] \} \\
 &= \sum_{k=n+1}^n \left(\frac{1}{k^r} - \frac{1}{(k+1)^r} \right) [S_k(\alpha, f^{(r)}) - f^{(r)}(\alpha)] \\
 &\quad - \frac{1}{(n+1)^r} [S_n(\alpha, f^{(r)}) - f^{(r)}(\alpha)], \tag{29}
 \end{aligned}$$

$$\begin{aligned}
 \sum_{k=n+1}^{\infty} \frac{1}{k^r} B_k(\alpha, \widetilde{f}^{(r)}) &= \sum_{k=n+1}^n \left(\frac{1}{k^r} - \frac{1}{(k+1)^r} \right) [S_k(\alpha, \widetilde{f}^{(r)}) - \widetilde{f}^{(r)}] \\
 &\quad - \frac{1}{(n+1)^r} [S_n(\alpha, \widetilde{f}^{(r)}) - \widetilde{f}^{(r)}(\alpha)]. \tag{30}
 \end{aligned}$$

Use of (28), (29) , (30) and (1), we get:

$$\begin{aligned}
 & \|f - S_n(\cdot, f)\|_{L^{p,\lambda}(\mathbb{T})} \\
 \leq & \sum_{k=n+1}^n \left(\frac{1}{k^r} - \frac{1}{(k+1)^r}\right) \|S_k(\cdot, f^{(r)}) - f^{(r)}\|_{L^{p,\lambda}(\mathbb{T})} \\
 & + \frac{1}{(n+1)^r} \|S_n(\cdot, f^{(r)}) - f^{(r)}\|_{L^{p,\lambda}(\mathbb{T})} \\
 & + \sum_{k=n+1}^n \left(\frac{1}{k^r} - \frac{1}{(k+1)^r}\right) \|S_k(\cdot, \widetilde{f}^{(r)}) - \widetilde{f}^{(r)}\|_{L^{p,\lambda}(\mathbb{T})} \\
 & + \frac{1}{(n+1)^r} \|S_n(\cdot, \widetilde{f}^{(r)}) - \widetilde{f}^{(r)}\|_{L^{p,\lambda}(\mathbb{T})} \\
 \leq & c_{28} \left\{ \sum_{k=n+1}^n \left(\frac{1}{k^r} - \frac{1}{(k+1)^r}\right) E_n(f^{(r)})_{p,\lambda} + \frac{1}{(n+1)^r} E_n(f^{(r)})_{p,\lambda} \right\} \\
 & + c_{29} \sum_{k=n+1}^n \left(\frac{1}{k^r} - \frac{1}{(k+1)^r}\right) E_n(\widetilde{f}^{(r)})_{p,\lambda} \cdot \\
 & + c_{29} \frac{1}{(n+1)^r} E_n(\widetilde{f}^{(r)})_{p,\lambda} \tag{31}
 \end{aligned}$$

Note that the sequence $\{E_n(f^{(r)})_{p,\lambda}\}$ is decreasing. Then using (31) and (1) we have

$$\begin{aligned}
 & \|f - S_n(\cdot, f)\|_{L^{p,\lambda}(\mathbb{T})} \\
 \leq & c_{30} E_n(f^{(r)})_{p,\lambda} \left\{ \sum_{k=n+1}^n \left(\frac{1}{k^r} - \frac{1}{(k+1)^r}\right) + \frac{1}{(n+1)^r} \right\} \\
 & + c_{31} E_n(\widetilde{f}^{(r)})_{p,\lambda} \left\{ \sum_{k=n+1}^n \left(\frac{1}{k^r} - \frac{1}{(k+1)^r}\right) + \frac{1}{(n+1)^r} \right\} \\
 \leq & c_{32} E_n(f^{(r)})_{p,\lambda} \left\{ \sum_{k=n+1}^n \left(\frac{1}{k^r} - \frac{1}{(k+1)^r}\right) + \frac{1}{(n+1)^r} \right\} \\
 = & \frac{2c_{33}}{(n+1)^r} E_n(f^{(r)})_{p,\lambda} \cdot \tag{32}
 \end{aligned}$$

On the other hand the inequality

$$E_n(f)_{p,\lambda} \leq \|f - S_n(\cdot, f)\|_{L^{p,\lambda}(\mathbb{T})} \tag{33}$$

holds. According to (32) and (33), we have inequality (12) . The proof of Theorem 1.1 is completed.

Proof of Theorem 1.2. Let $f \in \mathcal{L}^{p,\lambda}(\mathbb{T})$ and $h \in \widetilde{W}_r^{p,\lambda}$. According to Theorem 1.1 and Corollary 1.1, we obtain:

$$\begin{aligned}
 E_n(f)_{p,\lambda} & \leq E_n(f - h + h)_{p,\lambda} \leq E_n(f - h)_{p,\lambda} + E_n(h)_{p,\lambda} \\
 & \leq c_{34} \left\{ \|f - h\|_{L^{p,\lambda}(\mathbb{T})} + \frac{1}{n^r} \|h^{(r)}\|_{L^{p,\lambda}(\mathbb{T})} \right\} \tag{34}
 \end{aligned}$$

Using (34) and definition of the K -functional we have

$$E_n(f)_{p,\lambda} \leq c_{35} K_r\left(f, \frac{1}{n}\right)_{p,\lambda} \cdot$$

The last inequality and Theorem 2.1 imply that:

$$E_n(f)_{p,\lambda} \leq c_{36} \Omega_r(f, \delta)_{p,\lambda}.$$

The proof of Theorem 1.2 is completed.

Proof of Theorem 1.3. Let $f \in L^{p,\lambda}(\Gamma)$. Then, from (11) we have $f_0 \in L^{p,\lambda}(\mathbb{T})$, $f_1 \in L^{p,\lambda}(\mathbb{T})$. According to (12), we obtain

$$f(\zeta) = f_0^+(\phi(\zeta)) - f_0^-(\phi(\zeta)), \quad f(\xi) = f_1^+(\phi_1(\xi)) - f_1^-(\phi_1(\xi)). \tag{35}$$

a.e. on Γ .

We prove that the rational function

$$R_n(f, z) = \sum_{k=0}^{\infty} a_k \Phi_k(z) + \sum_{k=1}^{\infty} b_k F_k\left(\frac{1}{z}\right)$$

satisfies the condition of Theorem 1.3 .

Let $z^* \in G^-$. Using the method of proof in [32], we can prove that $f_0^-(\phi(\zeta)) \in E^{p,\lambda}(G^-) \in E^1(G^-)$. Then it is clear that

$$\frac{1}{2\pi i} \int_{\Gamma} \frac{f_0^-(\phi(\zeta))}{\zeta - z^*} d\zeta = -f_0^-(\phi(z^*)).$$

Then last equality, (5) and (15) imply that

$$\begin{aligned} \sum_{k=0}^n a_k \Phi_k(z^*) &= \sum_{k=0}^n a_k [\phi(z^*)]^k + \frac{1}{2\pi i} \int_{\Gamma} \frac{\sum_{k=0}^n a_k [\phi(\zeta)]^k}{\zeta - z^*} d\zeta \\ &= \sum_{k=0}^n a_k [\phi(z^*)]^k + \frac{1}{2\pi i} \int_{\Gamma} \frac{\sum_{k=0}^n a_k [\phi(\zeta)]^k - f_0^+[\phi(\zeta)]}{\zeta - z^*} d\zeta \\ &\quad + \frac{1}{2\pi i} \int_{\Gamma} \frac{f(\zeta)}{\zeta - z^*} d\zeta - f_0^-[\phi(z^*)]. \end{aligned} \tag{36}$$

Using (7) and (36), we find:

$$\begin{aligned} \sum_{k=0}^n a_k \Phi_k(z^*) &= \sum_{k=0}^n a_k [\phi(z^*)]^k + \frac{1}{2\pi i} \int_{\Gamma} \frac{\sum_{k=0}^n a_k [\phi(\zeta)]^k}{\zeta - z^*} d\zeta \\ &= \sum_{k=0}^n a_k [\phi(z^*)]^k + \frac{1}{2\pi i} \int_{\Gamma} \frac{\sum_{k=0}^n a_k [\phi(\zeta)]^k - f_0^+[\phi(\zeta)]}{\zeta - z^*} d\zeta \\ &\quad + f^-(z^*) - f_0^-[\phi(z^*)]. \end{aligned} \tag{37}$$

Taking the limit as $z^* \rightarrow z \in \Gamma$ along all non-tangential paths outside Γ and considering (8),(9), (35) and (37), we have

$$f^+(z) - \sum_{k=0}^n a_k \Phi_k(z^*) = \frac{1}{2} \left[f_0^+ [\phi(z^*)] - \sum_{k=0}^n a_k [\phi(z^*)]^k \right] + S_\Gamma \left(\left[f_0^+ [\phi(z^*)] - \sum_{k=0}^n a_k [\phi(z^*)]^k \right] \right). \tag{38}$$

By [47] the singular operator $S_\Gamma : L^{p,\lambda}(\Gamma) \rightarrow L^{p,\lambda}(\Gamma)$ is bounded. Then using (35), Minkowski’s inequality, Lemma 2.1 and 2.2, we have:

$$\begin{aligned} & \left\| f^+(z) - \sum_{k=0}^n a_k \Phi_k(z^*) \right\|_{L^{p,\lambda}(\Gamma)} \\ & \leq \frac{1}{2} \left\| f_0^+ [\phi(z^*)] - \sum_{k=0}^n a_k [\phi(z^*)]^k \right\|_{L^{p,\lambda}(\Gamma)} + \left\| S_\Gamma \left(\left[f_0^+ [\phi(z^*)] - \sum_{k=0}^n a_k [\phi(z^*)]^k \right] \right) \right\|_{L^{p,\lambda}(\Gamma)} \\ & \leq \frac{1}{2} \left\| f_0^+(w) - \sum_{k=0}^n a_k w^k \right\|_{L^{p,\lambda}(\Gamma)} + c_{37} \left\| f_0^+(w) - \sum_{k=0}^n a_k w^k \right\|_{L^{p,\lambda}(\Gamma)} \\ & \leq c_{38} \left\| f_0^+(w) - \sum_{k=0}^n a_k w^k \right\|_{L^{p,\lambda}(\Gamma)} \leq c_{39} \left\| f_0^+(w) - \sum_{k=0}^n \alpha_k (f_0^+) w^k \right\|_{L^{p,\lambda}(\Gamma)} \\ & \leq c_{40} \Omega_r(f_0^+, 1/n)_{p,\lambda} \leq c_{41} \Omega_r(f_0, 1/n)_{p,\lambda}. \end{aligned} \tag{39}$$

Let $z^* \in G$. Using the method of proof in [32] we can prove that $f^-(\phi_1(\zeta)) \in E^{p,\lambda}(G^-) \in E^1(G^-)$. Therefore,

$$\frac{1}{2\pi i} \int_\Gamma \frac{f_1^-(\phi_1(\zeta))}{\zeta - z^*} d\zeta = f_1^{--}(\phi_1(z^*)),$$

Then the last equality, (5) and (35) imply that

$$\begin{aligned} \sum_{k=1}^n b_k F_k\left(\frac{1}{z^*}\right) &= \sum_{k=1}^n b_k [\phi_1(z^*)]^k - \frac{1}{2\pi i} \int_\Gamma \frac{\sum_{k=1}^n b_k [\phi_1(\xi)]^k}{\xi - z^*} d\xi \\ &= \sum_{k=1}^n b_k [\phi_1(z^*)]^k - \frac{1}{2\pi i} \int_\Gamma \frac{\left(\sum_{k=1}^n b_k [\phi_1(\xi)]^k - f_1^+[\phi_1(\xi)]\right)}{\xi - z^*} d\xi \\ &\quad - \frac{1}{2\pi i} \int_\Gamma \frac{f(\xi)}{\xi - z^*} d\xi - \frac{1}{2\pi i} \int_\Gamma \frac{f_1^-(\phi_1(\zeta))}{\zeta - z^*} d\zeta \\ &= \sum_{k=1}^n b_k [\phi_1(z^*)]^k - \frac{1}{2\pi i} \int_\Gamma \frac{\left(\sum_{k=1}^n b_k [\phi_1(\xi)]^k - f_1^+[\phi_1(\xi)]\right)}{\xi - z^*} d\xi \\ &\quad - f^+(z^*) - f_1^-[\phi_1(z^*)]. \end{aligned}$$

Taking the limit as $z^* \rightarrow z$ along all nontangential paths inside of Γ we have

$$\sum_{k=1}^n b_k F_k\left(\frac{1}{z}\right) = \sum_{k=1}^n b_k [\phi_1(z)]^k - \frac{1}{2} \left(\sum_{k=1}^n b_k [\phi_1(z)]^k - f_1^+[\phi_1(z)] \right)$$

$$-S_{\Gamma} \left(\sum_{k=1}^n b_k [\phi_1(z)]^k - f_1^+ [\phi_1(z)] \right) - f^+(z) - f_1^- [\phi_1(z)]$$

a.e. on Γ . Consideration of (9) and (35) gives us

$$f^-(z) + \sum_{k=1}^n b_k F_k \left(\frac{1}{z} \right) = \frac{1}{2} \left(\sum_{k=1}^n b_k [\phi_1(z)]^k - f_1^+ [\phi_1(z)] \right) - S_{\Gamma} \left(\sum_{k=1}^n b_k [\phi_1(z)]^k - f_1^+ [\phi_1(z)] \right) \tag{40}$$

Using (40), Minkowski’s inequality and the boundedness of S_{Γ} in $L^{p,\lambda}(\Gamma)$ [47], Lemma 2.1 and 2.2, we obtain:

$$\begin{aligned} & \left\| f^-(z) + \sum_{k=1}^n b_k F_k \left(\frac{1}{z} \right) \right\|_{L^{p,\lambda}(\Gamma)} \\ & \leq \left\| \frac{1}{2} \left(\sum_{k=1}^n b_k [\phi_1(z)]^k - f_1^+ [\phi_1(z)] \right) \right\|_{L^{p,\lambda}(\Gamma)} + \left\| S_{\Gamma} \left(\sum_{k=1}^n b_k [\phi_1(z)]^k - f_1^+ [\phi_1(z)] \right) \right\|_{L^{p,\lambda}(\Gamma)} \\ & \leq \frac{1}{2} \left\| \sum_{k=1}^n b_k w^k - f_1^+(w) \right\|_{L^{p,\lambda}(\mathbb{T})} + c_{42} \left\| \sum_{k=1}^n b_k w^k - f_1^+(w) \right\|_{L^{p,\lambda}(\mathbb{T})} \\ & \leq c_{43} \left\| \sum_{k=1}^n b_k w^k - f_1^+(w) \right\|_{L^{p,\lambda}(\mathbb{T})} \\ & = c_{44} \left\| \sum_{k=1}^n \beta_k (f_1^+) w^k - f_1^+(w) \right\|_{L^{p,\lambda}(\mathbb{T})} \\ & \leq c_{45} \Omega_r (f_1^+, 1/n)_{p,\lambda} \leq c_{46} \Omega_r (f_1, 1/n)_{p,\lambda}. \end{aligned} \tag{41}$$

Now combining (8), (39) and (41) we obtain

$$\|f - R_n(\cdot, f)\|_{L^{p,\lambda}(\Gamma)} \leq c_{47}(p) \{ \Omega_r (f_0, 1/n)_{p,\lambda} + \Omega_r (f_1, 1/n)_{p,\lambda} \}$$

The proof of Theorem 1.3 is completed.

Proof of Theorem 1.4. Let $z^* \in G^-$. If $f \in E^{p,\lambda}(G)$, then $f \in E^p(G)$ and $\frac{f(z)}{z-z^*} \in E^p(G)$. Therefore, $\int_{\Gamma} \frac{f(z)}{z-z^*} = 0$. That is $f^-(z) = 0$ a.e. on Γ . Then taking into account (11), we have:

$$\left\| f_0^+(w) - \sum_{k=0}^n a_k w^k \right\|_{L^{p(\cdot),\lambda(\cdot)}(\mathbb{T})} \leq c_{48} \Omega_r \left(f_0, \frac{1}{n} \right)_{p,\lambda}, \forall n \in \mathbb{N},$$

$$\left\| f_0^+(z) - \sum_{k=0}^n a_k \Phi_k(z) \right\|_{L^{p,\lambda}(\Gamma)} \leq c_{49} \left\| f_0^+(w) - \sum_{k=0}^n a_k w^k \right\|_{L^{p,\lambda}(\mathbb{T})}$$

we have the inequality (13) of Theorem 1.4.

Proof of Theorem 1.5. Let $z^* \in G$ and $f \in E^{p,\lambda}(G^-)$. It is clear that $\int_{\Gamma} \frac{f(\zeta)}{\zeta - z^*} = f(\infty)$. Then we obtain $f^+(z) = f(\infty)$ a.e. on Γ . Now combining (11), we get:

$$\left\| f_1^+(w) - \sum_{k=0}^n b_k w^k \right\|_{L^{p,\lambda}(\mathbb{T})} \leq c_{50}(p) \Omega_r \left(f_1, \frac{1}{n} \right)_{p,\lambda}, \forall n \in \mathbb{N},$$

$$\left\| f^-(z) - \sum_{k=0}^n b_k F_k \left(\frac{1}{z} \right) \right\|_{L^{p,\lambda}(\Gamma)} \leq c_{51} \left\| f_1^+(w) - \sum_{k=0}^n b_k w^k \right\|_{L^{p,\lambda}(\mathbb{T})}$$

we obtain the inequality (14) of Theorem 1.5.

Acknowledgement. The author would like to thank the referee for his/her valuable comments and suggestions.

References

- [1] F. G. Abdullayev, I. A. Shevchuk, *Uniform estimates for polynomial approximation in domains with corners*, J. Approximation Theory, 137 (2) (2005), 143-165.
- [2] F. G. Abdullayev, P. Özkartepe, V. V. Savchuk, A. I. Shidlich, *Exact constants in direct and inverse approximation theorems for functions of several variables in the spaces S^p* , Filomat 33 (5) (2019), 1471-1484.
- [3] F. G. Abdullayev, S. Chaichenko, M. I. Kyzy, A. Shidlich, *Direct and inverse approximation theorems in the weighted Orlicz-type with variable exponent*, Turkish J. of Math. 44 (1) (2020), 284-299.
- [4] F. G. Abdullayev, S. Chaichenko, M. Imashkyzy, A. Shidlich, *Jackson-type inequalities and widths of functional classes in the Musielak-Orlicz type spaces*, Rocky Mountain J. Math. 51 (4) (2021), 1143-1155.
- [5] S. Y. Alper, *Approximation in the mean of analytic functions of class E^p* , In: Investigations on the Modern Problems of the Function Theory of a Complex Variable, Gos. Izdat. Fiz.-Mat. Lit., Moscow, 1960, pp. 272-286. (In Russian.)
- [6] J. E. Andersson, *On the degree of polynomial approximation in $E^p(D)$* . J. Approximation Theory 19 (1977), 61-68.
- [7] V. V. Andrievskii, D. M. Israfilov., *Approximations of functions on quasiconformal curves by rational functions*, Izv. Akad. Nauk Azerb. SSR ser. Fiz.-Tekhn. Math. 36 (4) (1980), (in Russian).
- [8] R. Akgün, *Trigonometric approximation of functions in generalized Lebesgue spaces with variable exponent*, Ukrainian Math. J. 63 (1) (2011), 3-23.
- [9] R. Akgün, *Approximation by polynomials in rearrangement invariant quasi Banach function spaces*, Banach J. Math. Anal. 6 (2) (2012), 113-131.
- [10] A. Almeida, S. Samko, *Approximation in generalized Morrey spaces*, Georgian Math. J. 25 (2) (2018), 155-168.
- [11] A. Almeida, J. Hasanov, S. Samko, *Maximal and potential operators in variable exponent Morrey spaces*, Georgian Math. J. 15 (2) (2008), 195-2008.
- [12] A. Böttcher, Y. I. Karlovich, Carleson Curves, Muckenhoupt Weights and Teoplitz Operators, Birkhauser-Verlag, 1997.
- [13] B. T. Bilalov, N. R. Ahmedzadeh, T. Z. Garayev, *Some remarks on solvability of Diriclet problem for Laplace equation in non-standatr function space*, Mediterr. J. Math. 19: 133 (2022), 25p.
- [14] A. Cavus, D. M. Israfilov, *Approximation by Faber-Laurent retional functions in the mean of functions of the class $L_p(\Gamma)$ with $1 < p < \infty$* , Approx. Theory Appl. 11 (1) (1995), 105-118.
- [15] Z. Cakir, C. Aykol, D. Soylemez, A. Serbetci, *Approximation by trigonometric poynomials in weighted Morrey spaces*, Tbilisi Mathematical Journal, 13 (1) (2020) 123-138.
- [16] Z. Cakir, C. Aykol, V. S. Guliyev, A. Serbetci, *Approximation by trigonometric polynomials in the variable exponent weighted Morrey spaces*, Carpathian Math. Publ. 13 (3) 2021, 750-763.
- [17] G. David, *Operateurs integraux singulers sur certains courbes du plan complexe*, Ann. Sci. Ecol. Norm. Super. 4 (1984), 157-189.
- [18] L. D'Onofrio, C. Sbordone and R. Schiattarella., *Grand Sobolev spaces and their application in geometric function theory and PDEs*, Journal of Fixed Point ttheory and Appl. , 13 (2013), 309-340.
- [19] P. L. Duren, *Theory of H^p spaces*, Academic Press, 1970.
- [20] E. M. Dyn'kin and B.P. Osilenker, *Weighted estimates for singular integrals and their appllications*, In: Mathematical Analysis, Vol. 21. Akad. Nauk. SSSR, Vsesoyuz. Inst. Nauchn. i Tekhn. Inform., Moscow, 1983, pp.42-129.
- [21] X. Fan, *The regularity of Lagrangians $f(x, \xi) = \|\xi\|_{\alpha(x)}$ with Höder exponents $\alpha(x)$* , Acta Math. Sin. (N:S.) 12 (3) (1996), 254-261.
- [22] D. Fan, S. Lu, D. Yang, *Regularity in Morrey spaces of strang solitions to nondivergence elliptic wquations with VMO coefficients*, Georgian Math. J. 5 (1998), 425-440.
- [23] L. Greco, T. Iwaniec and C. Sbordone., *Inverting the p harmonic operator*, Manuscripta Math, 92 (1997), 249-258.
- [24] V. S. Guliyev, J. Hasanov, S. Samko, *Boundedness of the maximal, potential and singular operators in the generalized variable exponent Morrey spaces*, Math. Scand. 107 (2010), 285-304.
- [25] V. S. Guliyev, A. Ghorbanalizadeh, Y. Sawano, *Approximation by trigonometric polynomials in variable exponent Morrey spaces*, Anal. Math. Phys. 9 (3) (2019), 1265-1286.

- [26] G. M. Goluzin, *Geometric Theory of Functions of a Complex Variable*, Translation of Mathematical Monographs, 26, Providence, RI: AMS, 1968
- [27] T. Iwaniec and C. Sbordone., *On integrability of the Jacobian under minimal hypotheses*, Arch Rational Mechanics Anal, **119** (1992), 129-143.
- [28] D. M. Israfilov, *Approximation by p -Faber polynomials in the weighted Smirnov class $E^p(G, w)$ and the Bieberbach polynomials*, Constr. Approx. **17** (2001), 335-351.
- [29] D. M. Israfilov, R. Akgün, *Approximate by polynomials and rational functions in rearrangement invariant spaces*, J. Math. Anal. Appl. **346** (2) (2008), 489-500.
- [30] D. M. Israfilov and N. P. Tozman, *Approximation by polynomials in Morrey- Smirnov classes*, East J. Approx, **14** (3) (2008), 255-269.
- [31] D. M. Israfilov, N. P. Tozman, *Approximation in Morrey-Smirnov classes*, Azerbaijan J. Math. **1** (2) (2011), 99-113.
- [32] D. M. Israfilov, A. Testici, *Approximation in Smirnov classes with variable exponent*, Complex Var. Elliptic Equ. **60** 9 (2015), 1243-1253.
- [33] D. M. Israfilov, A. Testici, *Approximation by Faber-Laurent rational functions in Lebesgue spaces with variable exponent*, Indagationes Math. **27** (2016), 914-922.
- [34] D. M. Israfilov, A. Testici, *Approximation in weighted generalized grand Lebesgue spaces*, Colloquium Mathematicum, **143** (1) (2016), 113-126.
- [35] M. Izuki, E. Nakai, S. Sawano, *Function spaces with variable exponents- an introduction*, Sci. Math. Jpn. **77** (2) (2014), 187-315:
- [36] S. Z. Jafarov, *Approximation by rational functions in Smirnov-Orlicz classes*, J. Math. Anal. Appl. **379** (2011), 870-877.
- [37] S. Z. Jafarov, *Approximation of function by p -Faber-Laurent functions*, Complex Variables and Elliptic Equations, **60** (3) (2015), 416-428.
- [38] A. Yu. Karlovich, *Algebras of singular integral operators with piecewise continuous coefficients on relexive Orlicz spaces*, Math. Nachr. **179** (1996), 187-222.
- [39] V. M. Kokilashvili., *Boundedness criteria for singular integrals in weighted grand Lebesgue spaces*, Jour. Math. Sci., **170** (3), 2010, 20-33.
- [40] V. M. Kokilashvili, V. Paatasvili, S. Samko, *Boundary value problems for analytic functions in the class of Cauchy type integrals with density in $L^{p(\cdot)}(\Gamma)$* , Boundary Value Problems, Vol. 2005, Hindawi Publ. Cor., 2005, pp.43-71.
- [41] V. M. Kokilashvili, S. Samko, *Weighted boundedness in Lebesgue spaces with variable exponentsof classical operators on Carleson curves*, Proc. A. Razmadze Math. Inst. **138** (2005) 106-110.
- [42] V. M. Kokilashvili, *A direct theorem on mean approximation of analytic functions by polynomials*, Soviet Math. Dokl. **10** (1969), 411-414.
- [43] V. M. Kokilashvili and S. Samko, *Singular integrals and potentials in some Banach function spaces with variable exponent*, J. Funct. Spaces Appl. **1** (1) (2003), 45-59.
- [44] V. M. Kokilashvili, A. Meskhi, *Boundednes of maximal and singular operators in Morrey spaces with variable exponent*, Armen. J. Math. **1** (1) (2008), 18-28.
- [45] A. Kufner, O. John, S. Fucik, *Function spaces*, p. 454+XV. Noordhoff Internationasl Publishing, London (1977).
- [46] A. L. Mazzucato, *Decomposition of Besov- Morrey spaces*, In: *Proceerings of the Conference on Harmonic Analysis*, in : *Contemp. Math. Amer. Math. Soc. Providence, RI*, **320** (2003), 279-294.
- [47] A. Meskhi, *Maximal functions and singular integrals in Morrey spaces associated with grand Lebesgue spaces*, Proc. A. Razmadze Math. Inst. **151** (2009), 130-143.
- [48] T. Ohno, *Continuity properties for logarithmic potentials of functions in Morrey spaces of variable exponent*, Hiroshima Math. J. **38** (3) (2008), 363-383.
- [49] A. A. Pekariski, *Rational approximation of absolutely continuous functions with derivatives in an Orlicz space*, Math. Sb. **45** (1983), 121-137.
- [50] Ch. Pommerenke, *Boundary Behavior of Conformal Maps*, Berlin, SpringerVerlag, 1992.
- [51] H. Raferio, N. Samko, S. Samko, *Morrey-Campanato spaces: an overviev, (English summary)* Operator theory, pseudo-differential equations, and mathematical physics, Oper. Theory Adv. Appl. **228** (2013), 293-323.
- [52] M. Ruzicka, *Elektorrheological fluids: modeling and mathematical theory*, Vol. 1748, Lecture notes in mathematics, Berlin: Springer-Verlag; 2000.
- [53] C. Sbordine., *Grand Sobolev spaces and their applications to variational problems*, Le Matematiche, **LI** (2) (1996), 335-347.
- [54] C. Sbordine., *Nonlinear elliptic equations with right hand side in nonstandard spaces*, Rend. Sem. Math. Fis. Modena, Supplemento al XLVI (1998), 361-368.
- [55] Y. Sawano, H. Tanako, *The Fatou property of block spaces*, J. Math. Sci. Univ. Tokyo **22** (2015), 663-683.
- [56] N. Samko, *Weighted Hardy and singular operators in Morrey spaces*, J. Math. Anal. Appl. **350** (1) (2009), 56-72
- [57] P. K. Suetin, *Series of Faber polynomials*, Gordon and Breach Science Publishers, 1998.
- [58] J. L. Walsh, H.G. Russel, *Integrated continuity conditions and degree of approximation by polynomials or by bounded analytic functions*. Trans. Amer. Math. Soc. **92** (1959), 355-370.
- [59] S. E. Warschawskii, *Über das Randverhalten der Ableitung der Abbildungsfunktionen bei Konformer Abbildung*, Math. Z., **35** (1932), 321-456
- [60] H. Yurt, A. Guven, *Approximation by Faber-Laurent rational functions on doubly connected domains*, New Zealand Journal of Math. **44** (2014), 113-124.