



## On the completeness of a system of Bessel functions of index $-3/2$ in weighted $L^2$ -space

Ruslan Khats'<sup>a</sup>

<sup>a</sup>Department of Mathematics and Economics, Drohobych Ivan Franko State Pedagogical University, 3 Stryis'ka Str., 82100 Drohobych, Ukraine

**Abstract.** In this paper, we study an integral representation of some class  $E_{2,+}$  of even entire functions of exponential type  $\sigma \leq 1$ . We also obtain an analog of the Paley-Wiener theorem related to the class  $E_{2,+}$ . In addition, we find necessary and sufficient conditions for the completeness of a system  $\{s_k \sqrt{x s_k} J_{-3/2}(x s_k) : k \in \mathbb{N}\}$  in the space  $L^2((0; 1); x^2 dx)$ , where  $J_{-3/2}$  be the Bessel function of the first kind of index  $-3/2$ ,  $(s_k)_{k \in \mathbb{N}}$  be a sequence of distinct nonzero complex numbers and  $L^2((0; 1); x^2 dx)$  be the weighted Lebesgue space of all measurable functions  $f : (0; 1) \rightarrow \mathbb{C}$  satisfying  $\int_0^1 x^2 |f(x)|^2 dx < +\infty$ . Those results are formulated in terms of sequences of zeros of functions from the class  $E_{2,+}$ . We also obtain some other sufficient conditions for the completeness of the considered system of Bessel functions. Our results complement similar results on completeness of the systems of Bessel functions of index  $\nu < -1$ ,  $\nu \notin \mathbb{Z}$ .

### 1. Introduction

Let  $L^2(X)$  be the space of all measurable functions  $f : X \rightarrow \mathbb{C}$  on a measurable set  $X \subseteq \mathbb{R}$  endowed with a norm

$$\|f\|_{L^2(X)}^2 := \int_X |f(x)|^2 dx,$$

let  $\gamma \in \mathbb{R}$  and let  $L^2((0; 1); x^\gamma dx)$  be the weighted Lebesgue space of all measurable functions  $f : (0; 1) \rightarrow \mathbb{C}$ , satisfying

$$\int_0^1 x^\gamma |f(x)|^2 dx < +\infty.$$

Let (see, for example, [2, p. 4], [17, p. 345], [27, p. 40])

$$J_\nu(z) = \sum_{k=0}^{\infty} \frac{(-1)^k (z/2)^{\nu+2k}}{k! \Gamma(\nu+k+1)}, \quad z = x + iy = r e^{i\varphi},$$

2020 *Mathematics Subject Classification.* Primary 33C10, 42C30, 44A15; Secondary 30B60, 30D20.

*Keywords.* Bessel function; Paley-Wiener theorem; Phragmén-Lindelöf theorem; Hahn-Banach theorem; Jensen's formula; Entire function of exponential type; Complete system.

Received: 06 June 2022; Accepted: 24 February 2023

Communicated by Hari M. Srivastava

Email address: khats@ukr.net (Ruslan Khats')

be the Bessel function of the first kind of index  $\nu \in \mathbb{R}$ , where  $\Gamma$  is the classical Gamma function (see, for details, [15, pp. 1505-1506]). By Hurwitz's theorem (see [2, p. 59], [27, p. 483]), for  $\nu > 1$  the function  $J_{-\nu}$  has an infinity of real zeros and also  $2[\nu]$  pairwise conjugate complex zeros, among them two pure imaginary zeros when  $[\nu]$  is an odd integer. In particular, the function  $J_{-3/2}$  has an infinite set  $\{s_k : k \in \mathbb{Z} \setminus \{0\}\}$  of zeros, among them  $s_1$  and  $s_{-1} = \bar{s}_1 = -s_1$  are two pure imaginary zeros, the positive zeros  $s_k, k \in \mathbb{N} \setminus \{1\}$ , and negative zeros  $s_{-k} := -s_k, k \in \mathbb{N} \setminus \{1\}$ . Moreover (see [17, p. 350], [27, p. 55]),  $\sqrt{z}J_{-3/2}(z) = -\sqrt{2/\pi}z^{-1}(\cos z + z \sin z)$  and the function  $s\sqrt{xs}J_{-3/2}(xs)$  belongs to the space  $L^2((0; 1); x^2 dx)$  for every  $s \in \mathbb{C}$ . A system of elements  $\{e_k : k \in \mathbb{N}\}$  in a separable Hilbert space  $\mathcal{H}$  is called complete (see [10, p. 131], [11, p. 4258]) if  $\overline{\text{span}}\{e_k : k \in \mathbb{N}\} = \mathcal{H}$ .

We remark that, in survey-cum-expository review article [14], Hari M. Srivastava presented a brief introductory overview and survey of some of the recent developments on the theory and applications of the Bessel functions and other higher transcendental functions. It is in [14] in which the interested reader can find some developments involving a hybrid version of several known extensions and generalizations of the Mittag-Leffler type functions, as well as the Hurwitz-Lerch type zeta functions, together with its associated fractional integrals and fractional derivatives (see, for details, [15]). In addition, Srivastava's investigations are motivated essentially by a number of extensive developments on the familiar Laplace and Hankel transforms as well as on the extensions and generalizations of each of these integral transforms. In particular, in the article [16], Hari M. Srivastava presented several (presumably new) properties and characteristics as well as inter-relationships among each of such general families of integral transforms as Srivastava's generalized Whittaker transform, Hardy's generalized Hankel transform and Srivastava's  $\epsilon$ -generalized Hankel transform. These results can indeed be appropriately specialized to deduce a large number of known or new relationships between various simpler integral transforms.

Basis properties (completeness, minimality, basicity) of the systems of Bessel functions has been studied in many papers (see, for instance, [1–9, 12, 13, 17–27]). In this case, various approximation properties of the systems of Bessel functions  $J_\nu$  for  $\nu < -1, \nu \notin \mathbb{Z}$ , were investigated in [5, 8, 9, 12, 13, 18, 19, 25, 26]. In particular, in [25] (see also [26]) it was proven that the system  $\{s_k \sqrt{xs_k} J_{-3/2}(xs_k) : k \in \mathbb{N}\}$  is complete in the space  $L^2((0; 1); x^2 dx)$  and the system  $\{s_k \sqrt{xs_k} J_{-3/2}(xs_k) : k \in \mathbb{N} \setminus \{1\}\}$  is complete, minimal and is not a basis in this space, where  $(s_k)_{k \in \mathbb{Z} \setminus \{0\}}, s_{-k} := -s_k$ , is a sequence of zeros of the function  $J_{-3/2}$ . In addition, in [12] it was shown that the system  $\{s_k^2 \sqrt{xs_k} J_{-5/2}(xs_k) : k \in \mathbb{N} \setminus \{1, 2\}\}$  is complete and minimal in  $L^2((0; 1); x^4 dx)$ , where  $(s_k)_{k \in \mathbb{N}}$  is a sequence of zeros of  $J_{-5/2}$ . Besides, in [13] has been established that the system  $\{s_k^{\nu-1/2} \sqrt{xs_k} J_{-\nu}(xs_k) : k \in \mathbb{N} \setminus \{1, 2; \dots; l\}\}$  is complete in  $L^2((0; 1); x^{2\nu-1} dx)$  if  $\nu = l + 1/2, l \in \mathbb{N}$  and  $(s_k)_{k \in \mathbb{N}}$  is a sequence of zeros of  $J_{-\nu}$ . However, the problem on completeness of this system in  $L^2((0; 1); x^{2\nu-1} dx)$  when  $\nu = l + 1/2, l \in \mathbb{N}$  and  $(s_k)_{k \in \mathbb{N}}$  is an arbitrary sequence of distinct nonzero complex numbers remains open. In this direction, in [18] the authors obtained a criterion for the completeness and minimality of a system  $\{xs_k \sqrt{xs_k} J_{-3/2}(xs_k) : k \in \mathbb{N}\}$  in  $L^2(0; 1)$  with an arbitrary sequence of nonzero complex numbers  $(s_k)_{k \in \mathbb{N}}$ . Also, in [19] it was proven that the system  $\{x^{-2}(s_k \sqrt{xs_k} J_{-3/2}(xs_k) - s_1 \sqrt{xs_1} J_{-3/2}(xs_1)) : k \in \mathbb{N} \setminus \{1\}\}$  is complete and minimal in  $L^2((0; 1); x^2 dx)$  where  $(s_k)_{k \in \mathbb{N}}$  is a sequence of distinct nonzero complex numbers such that  $s_k^2 \neq s_m^2$  for  $k \neq m$ . In addition, using methods of [5–7, 11, 18–21], in [8] were established necessary and sufficient completeness conditions of a system  $\{s_k^2 \sqrt{xs_k} J_{-5/2}(xs_k) : k \in \mathbb{N}\}$  in the space  $L^2((0; 1); x^4 dx)$  with an arbitrary sequence of distinct nonzero complex numbers  $(s_k)_{k \in \mathbb{N}}$  in terms of entire functions.

The aim of this paper is to obtain an analog of the Paley-Wiener theorem for some class  $E_{2,+}$  of even entire functions of exponential type  $\sigma \leq 1$  (see Theorem 3.1). In addition, we will find new necessary and sufficient conditions for the completeness of a system  $\{\theta_k : k \in \mathbb{N}\}$  with  $\theta_k(x) := s_k \sqrt{xs_k} J_{-3/2}(xs_k)$  in the space  $L^2((0; 1); x^2 dx)$  in terms of entire functions from  $E_{2,+}$  with the set of zeros coinciding with the sequence of distinct nonzero complex numbers  $(s_k)_{k \in \mathbb{N}}$  (see Theorems 3.6–3.11). Also, in Theorems 3.12 and 3.13, we obtain some other sufficient conditions of the completeness of the system  $\{\theta_k : k \in \mathbb{N}\}$ . Our results complement the results of papers [3, 5, 8, 9, 12, 13, 18, 19, 25, 26].

## 2. Preliminaries

An entire function  $G$  is said to be of exponential type  $\sigma \in [0; +\infty)$  ([10, p. 4], [11, p. 4262]) if for any  $\varepsilon > 0$  there exists a constant  $c(\varepsilon)$  such that  $|G(z)| \leq c(\varepsilon) \exp((\sigma + \varepsilon)|z|)$  for all  $z \in \mathbb{C}$ .

Denote by  $PW_\sigma^2$  the set of all entire functions of exponential type  $\sigma \in (0; +\infty)$  whose narrowing on  $\mathbb{R}$  belongs to the space  $L^2(\mathbb{R})$ , and by  $PW_{\sigma,+}^2$  denote the class of even entire functions from  $PW_\sigma^2$ . By the Paley-Wiener theorem (see [10, p. 69], [11, p. 4263]), the class  $PW_\sigma^2$  coincides with the class of functions  $G$  admitting the representation

$$G(z) = \int_{-\sigma}^{\sigma} e^{itz} g(t) dt, \quad g \in L^2(-\sigma; \sigma),$$

and the class  $PW_{\sigma,+}^2$  consists of the functions  $G$  representable in the form

$$G(z) = \int_0^{\sigma} \cos(tz) g(t) dt, \quad g \in L^2(0; \sigma).$$

Moreover,  $\|g\|_{L^2(0;\sigma)} = \sqrt{2/\pi} \|G\|_{L^2(0;+\infty)}$  and

$$g(t) = \frac{2}{\pi} \int_0^{+\infty} G(z) \cos(tz) dz.$$

Let  $\log^+ x = \max(0; \log x)$  for  $x > 0$ . Here and so on by  $C_1, C_2, \dots$  we denote arbitrary positive constants. To prove our main results we need the following auxiliary lemmas.

**Lemma 2.1.** (see [18, p. 13], [25, p. 39]) *Let an entire function  $Q$  be defined by the formula*

$$Q(z) = -\sqrt{\frac{2}{\pi}} \int_0^1 (\cos(tz) + tz \sin(tz)) q(t) dt, \quad q \in L^2(0; 1). \tag{1}$$

*Then for all  $z = x + iy = re^{i\varphi} \in \mathbb{C}$ , we have*

$$|Q(z)| \leq C_1 \frac{e^{|\operatorname{Im} z|}}{\sqrt{1 + |\operatorname{Im} z|}} (1 + |z|),$$

*and  $Q$  is an even entire function of exponential type  $\sigma \leq 1$ .*

**Lemma 2.2.** (see [11, p. 4263]) *Let  $Q$  be an entire function of exponential type  $\sigma \leq 1$  for which*

$$\int_{-\infty}^{+\infty} \frac{\log^+ |Q(t)|}{1 + t^2} dt < +\infty,$$

*and let  $(s_k)_{k \in \mathbb{N}}$  be a sequence of nonzero roots of the function  $Q(z)$ . Then*

$$\sum_{k \in \mathbb{N}} \left| \operatorname{Im} \frac{1}{s_k} \right| < +\infty.$$

## 3. Main results

Our main results are the following statements.

**Theorem 3.1.** *An entire function  $Q$  has the representation*

$$Q(z) = \int_0^1 z \sqrt{tz} J_{-3/2}(tz) t^2 h(t) dt \quad (2)$$

with some function  $h \in L^2((0; 1); t^2 dt)$  if and only if it is an even entire function of exponential type  $\sigma \leq 1$  such that

$$Q(0) = -\sqrt{\frac{2}{\pi}} \int_0^1 th(t) dt, \quad (3)$$

and the function  $z^{-1}Q'(z)$  belongs to the space  $PW_{1,+}^2$ . If these conditions are fulfilled, then

$$h(t) = -\sqrt{\frac{2}{\pi}} \frac{1}{t^3} \int_0^{+\infty} \frac{Q'(z)}{z} \cos(tz) dz.$$

*Proof. Necessity.* Let  $Q$  has the representation (2) with some function  $h \in L^2((0; 1); t^2 dt)$ . Since

$$z \sqrt{tz} J_{-3/2}(tz) = -\sqrt{\frac{2}{\pi}} \frac{\cos(tz) + tz \sin(tz)}{t},$$

we have

$$Q(z) = -\sqrt{\frac{2}{\pi}} \int_0^1 (\cos(tz) + tz \sin(tz)) th(t) dt, \quad Q(0) = -\sqrt{\frac{2}{\pi}} \int_0^1 th(t) dt.$$

Therefore, by Lemma 2.1, the function  $Q$  is an even entire function of exponential type  $\sigma \leq 1$ , and

$$Q'(z) = -\sqrt{\frac{2}{\pi}} \int_0^1 t^3 z \cos(tz) h(t) dt, \quad Q'(0) = 0, \quad \frac{Q'(z)}{z} = -\sqrt{\frac{2}{\pi}} \int_0^1 \cos(tz) t^2 q(t) dt,$$

where  $q(t) := th(t)$ . Since  $h \in L^2((0; 1); t^2 dt)$ , we have  $q \in L^2(0; 1)$ , and in accordance with the Paley-Wiener theorem, the function  $z^{-1}Q'(z)$  belongs to the space  $PW_{1,+}^2$ . *Sufficiency.* If all the conditions of the theorem hold, then from the formula for the inverse Fourier cosine-transformation it follows that the function

$$q(t) = -\sqrt{\frac{2}{\pi}} \frac{1}{t^2} \int_0^{+\infty} \frac{Q'(z)}{z} \cos(tz) dz$$

belongs to the space  $L^2(0; 1)$ , and

$$Q'(z) = -\sqrt{\frac{2}{\pi}} \int_0^1 z \cos(tz) t^3 h(t) dt.$$

Using Fubini's theorem, we get

$$Q(z) - Q(0) = -\sqrt{\frac{2}{\pi}} \int_0^1 t^3 h(t) dt \int_0^z w \cos(tw) dw = -\sqrt{\frac{2}{\pi}} \int_0^1 (\cos(tz) + tz \sin(tz) - 1) th(t) dt.$$

Further, using (3), we obtain

$$Q(z) = -\sqrt{\frac{2}{\pi}} \int_0^1 (\cos(tz) + tz \sin(tz)) th(t) dt = \int_0^1 z \sqrt{tz} J_{-3/2}(tz) t^2 h(t) dt.$$

Thus, the theorem is proved.  $\square$

Let  $\widetilde{E}_{2,+}$  be the class of entire functions  $Q$  that can be presented in the form (2) with some function  $h \in L^2((0; 1); t^2 dt)$ , and let  $E_{2,+}$  be the class of even entire functions  $Q$  of exponential type  $\sigma \leq 1$  satisfying (3) with  $h \in L^2((0; 1); t^2 dt)$  and the function  $z^{-1}Q'(z)$  belongs to the space  $PW_{1,+}^2$ .

**Corollary 3.2.**  $\widetilde{E}_{2,+} = E_{2,+}$ .

**Corollary 3.3.** The class  $E_{2,+}$  coincides with a set of entire functions  $Q$  representing in the form (1).

**Example 3.4.** Evidently,  $\cos z \notin E_{2,+}$ .

**Example 3.5.** The function

$$Q(z) = -\sqrt{\frac{2}{\pi}} \frac{\cos z}{z^2 - \pi^2/4} \left( 1 - \frac{4(-\pi + 2)}{\pi^3} (z^2 - \pi^2/4) \right)$$

belongs to  $E_{2,+}$  with

$$h(t) = \frac{4}{\pi^3 t^3} \left( -2 \cos\left(\frac{\pi}{2}t\right) - \pi t \sin\left(\frac{\pi}{2}t\right) + 2 \right).$$

**Theorem 3.6.** Let  $(s_k)_{k \in \mathbb{N}}$  be a sequence of nonzero complex numbers such that  $s_k^2 \neq s_n^2$  for  $k \neq n$ . The system  $\{\theta_k : k \in \mathbb{N}\}$  is incomplete in  $L^2((0; 1); x^2 dx)$  if and only if a sequence  $(s_k)_{k \in \mathbb{Z} \setminus \{0\}}$ , where  $s_{-k} := -s_k$ ,  $k \in \mathbb{N}$ , is a subsequence of zeros of some nonzero entire function  $Q \in E_{2,+}$ .

*Proof.* According to the Hahn-Banach theorem (see [10, p. 131], [11, p. 4258]), the system  $\{\theta_k : k \in \mathbb{N}\}$  is incomplete in  $L^2((0; 1); x^2 dx)$  if and only if there exists a nonzero function  $h \in L^2((0; 1); x^2 dx)$  such that

$$\int_0^1 s_k \sqrt{ts_k} J_{-3/2}(ts_k) t^2 h(t) dt = 0$$

for all  $k \in \mathbb{N}$ . Hence, taking into account Theorem 3.1, we obtain the required proposition. Theorem 3.6 is proved.  $\square$

**Theorem 3.7.** Let  $(s_k)_{k \in \mathbb{N}}$  be a sequence of distinct nonzero complex numbers such that  $|\operatorname{Im} s_k| \geq \delta |s_k|$  for all  $k \in \mathbb{N}$  and some  $\delta > 0$ . If a system  $\{\theta_k : k \in \mathbb{N}\}$  is complete in  $L^2((0; 1); x^2 dx)$ , then

$$\sum_{k=1}^{\infty} \frac{1}{|s_k|} = +\infty. \tag{4}$$

*Proof.* Suppose, to the contrary, that the system  $\{\theta_k : k \in \mathbb{N}\}$  is not complete in the space  $L^2((0; 1); x^2 dx)$ . Then, by Theorem 3.6, there exists a nonzero entire function  $Q \in E_{2,+}$  for which the sequence  $(s_k)_{k \in \mathbb{Z} \setminus \{0\}}$  is a subsequence of zeros. By virtue of Corollary 3.3, the function  $Q$  is of the kind (1). Due to Lemma 2.1, we have  $|Q(x)| \leq C_1(1 + |x|)$  for all  $x \in \mathbb{R}$ . This implies

$$\int_{-\infty}^{+\infty} \frac{\log^+ |Q(x)|}{1 + x^2} dx < +\infty.$$

Therefore, by Lemma 2.2, we get

$$\sum_{k \in \mathbb{N}} \left| \operatorname{Im} \frac{1}{s_k} \right| < +\infty.$$

Since  $|\operatorname{Im} s_k| \geq \delta |s_k|$  for all  $k \in \mathbb{N}$  and some  $\delta > 0$ , and

$$\left| \operatorname{Im} \frac{1}{s_k} \right| = \frac{|\operatorname{Im} s_k|}{|s_k|^2} \geq \frac{\delta}{|s_k|},$$

we have

$$\sum_{k=1}^{\infty} \frac{1}{|s_k|} < +\infty.$$

This contradicts condition (4). Thus, the theorem is proved.  $\square$

**Theorem 3.8.** *Let  $(s_k)_{k \in \mathbb{N}}$  be a sequence of distinct nonzero complex numbers such that  $s_k^2 \neq s_m^2$  for  $k \neq m$ . Let a sequence  $(s_k)_{k \in \mathbb{Z} \setminus \{0\}}$ , where  $s_{-k} := -s_k$ , be a sequence of zeros of some even entire function  $G$  of exponential type  $\sigma \leq 1$  for which on the rays  $\{z : \arg z = \varphi_j\}$ ,  $j \in \{1; 2; 3; 4\}$ ,  $\varphi_1 \in [0; \pi/2)$ ,  $\varphi_2 \in [\pi/2; \pi)$ ,  $\varphi_3 \in (\pi; 3\pi/2]$ ,  $\varphi_4 \in (3\pi/2; 2\pi)$ , we have  $|G(z)| \geq C_2(1 + |z|)e^{|\operatorname{Im} z|}$ . Then the system  $\{\theta_k : k \in \mathbb{N}\}$  is complete in  $L^2((0; 1); x^2 dx)$ .*

*Proof.* Assume the converse. Then, according to Theorem 3.6, there exists a nonzero even entire function  $Q \in E_{2,+}$  for which the sequence  $(s_k)_{k \in \mathbb{Z} \setminus \{0\}}$  is a subsequence of zeros. Let  $V(z) = Q(z)/G(z)$ . Then  $V$  is an even entire function of order  $\tau \leq 1$ , for which by Corollary 3.3 and Lemma 2.1, we obtain

$$|V(z)| \leq C_3 \frac{1}{\sqrt{1 + |\operatorname{Im} z|}}, \quad \arg z = \varphi_j, \quad j \in \{1; 2; 3; 4\}.$$

Therefore, according to the Phragmén-Lindelöf theorem (see [10, p. 38], [11, p. 4263]), we get  $V(z) \equiv 0$ . Hence  $Q(z) \equiv 0$ . This contradiction proves the theorem.  $\square$

**Corollary 3.9.** *Let  $(s_k)_{k \in \mathbb{N}}$  be a sequence of zeros of the function  $J_{-3/2}$ . Then the system  $\{\theta_k : k \in \mathbb{N}\}$  is complete in  $L^2((0; 1); x^2 dx)$ .*

*Proof.* Indeed, a sequence  $(s_k)_{k \in \mathbb{Z} \setminus \{0\}}$ , where  $s_{-k} = -s_k$ , is a sequence of zeros of an entire function  $G(z) = \cos z + z \sin z$ , and this function satisfies the conditions of Theorem 3.8. Therefore, a system  $\{\theta_k : k \in \mathbb{N}\}$  is complete in  $L^2((0; 1); x^2 dx)$ . Corollary 3.9 is proved.  $\square$

**Theorem 3.10.** *Let  $(s_k)_{k \in \mathbb{N}}$  be a sequence of distinct nonzero complex numbers such that  $s_k^2 \neq s_m^2$  for  $k \neq m$ . Let a sequence  $(s_k)_{k \in \mathbb{Z} \setminus \{0\}}$ , where  $s_{-k} := -s_k$ , be a sequence of zeros of some even entire function  $G \notin E_{2,+}$  of exponential type  $\sigma \leq 1$  for which on the rays  $\{z : \arg z = \varphi_j\}$ ,  $j \in \{1; 2; 3; 4\}$ ,  $\varphi_1 \in [0; \pi/2)$ ,  $\varphi_2 \in [\pi/2; \pi)$ ,  $\varphi_3 \in (\pi; 3\pi/2]$ ,  $\varphi_4 \in (3\pi/2; 2\pi)$ , the inequality  $|G(z)| \geq C_4(1 + |z|)^{-\alpha} e^{|\operatorname{Im} z|}$  holds with  $\alpha < 3/2$ . Then the system  $\{\theta_k : k \in \mathbb{N}\}$  is complete in  $L^2((0; 1); x^2 dx)$ .*

*Proof.* Assume the converse. Then, according to Theorem 3.6, there exists a nonzero even entire function  $Q \in E_{2,+}$  for which the sequence  $(s_k)_{k \in \mathbb{Z} \setminus \{0\}}$  is a subsequence of zeros. Let  $V(z) = Q(z)/G(z)$ . Then  $V$  is an even entire function of order  $\tau \leq 1$ , for which by Corollary 3.3 and Lemma 2.1, we get

$$|V(z)| \leq C_5 \frac{(1 + |z|)^{\alpha+1}}{\sqrt{1 + |\operatorname{Im} z|}}, \quad \arg z = \varphi_j, \quad j \in \{1; 2; 3; 4\}.$$

Since  $\alpha + 1 < 5/2$ , according to the Phragmén-Lindelöf theorem, the function  $V$  is a polynomial of degree  $\zeta < 2$ . However,  $V$  is an even entire function, and therefore the function  $V$  is a constant. Hence,  $Q(z) = C_6 G(z)$  and  $Q \notin E_{2,+}$ . Thus, we have a contradiction and the proof of the theorem is completed.  $\square$

**Theorem 3.11.** Let  $(s_k)_{k \in \mathbb{N}}$  be a sequence of distinct nonzero complex numbers such that  $s_k^2 \neq s_m^2$  for  $k \neq m$ . Let a sequence  $(s_k)_{k \in \mathbb{Z} \setminus \{0\}}$ , where  $s_{-k} := -s_k$ , be a sequence of zeros of some even entire function  $F \notin E_{2,+}$  of exponential type  $\sigma \leq 1$  such that for some  $\alpha < 1$  and  $\eta \in \mathbb{R}$

$$|F(x + i\eta)| \geq \delta|x|^{-\alpha}, \quad \delta > 0, \quad |x| > 1. \tag{5}$$

Then the system  $\{\theta_k : k \in \mathbb{N}\}$  is complete in  $L^2((0; 1); x^2 dx)$ .

*Proof.* Let  $F \notin E_{2,+}$  and the inequality (5) is true. Suppose, to the contrary, that the system  $\{\theta_k : k \in \mathbb{N}\}$  is not complete in  $L^2((0; 1); x^2 dx)$ . Then, by Theorem 3.6, there exists a nonzero even entire function  $Q \in E_{2,+}$  which vanishes at the points  $s_k$ . However, the sequence  $(s_k)_{k \in \mathbb{Z} \setminus \{0\}}$  is a sequence of zeros of an even entire function  $F(z) \notin E_{2,+}$  of exponential type  $\sigma \leq 1$ . Therefore,  $T(z) = Q(z)/F(z)$  is an even entire function of order  $\tau \leq 1$ . Since  $Q \in E_{2,+}$ , taking into account Corollary 3.3 and Lemma 2.1, we obtain

$$|Q(x + i\eta)| \leq C_7 \frac{e^{|\eta|}}{\sqrt{1 + |\eta|}} \left(1 + \sqrt{x^2 + \eta^2}\right), \quad x \in \mathbb{R}.$$

Using (5), we get  $|T(x + i\eta)| \leq C_8(1 + |x|)^{1+\alpha}$ ,  $x \in \mathbb{R}$ . It is easy to see that  $T(z)$  is a polynomial of degree  $\zeta < 2$ . Further, since  $T$  is an even entire function, we have  $T(z) = C_9$ . Furthermore,  $F(z) = C_{10}Q(z)$  and  $F(z) \in E_{2,+}$ . This contradiction concludes the proof of the theorem.  $\square$

Let  $n(t)$  be the number of points of the sequence  $(s_k)_{k \in \mathbb{N}} \subset \mathbb{C}$  satisfying the inequality  $|s_k| \leq t$ , i.e.,  $n(t) := \sum_{|s_k| \leq t} 1$ , and let

$$N(r) := \int_0^r \frac{n(t)}{t} dt, \quad r > 0.$$

**Theorem 3.12.** Let  $(s_k)_{k \in \mathbb{N}}$  be an arbitrary sequence of distinct nonzero complex numbers. If

$$\limsup_{r \rightarrow +\infty} \left( N(r) - \frac{2r}{\pi} + \frac{1}{2} \log r - \log(1 + r) \right) = +\infty,$$

then the system  $\{\theta_k : k \in \mathbb{N}\}$  is complete in  $L^2((0; 1); x^2 dx)$ .

*Proof.* It suffices to assume the incompleteness of the system  $\{\theta_k : k \in \mathbb{N}\}$  and prove that

$$\limsup_{r \rightarrow +\infty} \left( N(r) - \frac{2r}{\pi} + \frac{1}{2} \log r - \log(1 + r) \right) < +\infty. \tag{6}$$

Due to Theorem 3.6, there exists a nonzero even entire function  $Q \in E_{2,+}$  of exponential type  $\sigma \leq 1$  for which the sequence  $(s_k)_{k \in \mathbb{N}}$  is a subsequence of zeros. We may consider that  $Q(0) = 1$ . Then, consecutively applying the Jensen formula (see [10, p. 10], [11, p. 4316]), Corollary 3.3 and Lemma 2.1, we obtain

$$\begin{aligned} N(r) &\leq \frac{1}{2\pi} \int_0^{2\pi} \log |Q(re^{i\varphi})| d\varphi \\ &\leq C_{11} + \frac{1}{2\pi} \int_0^{2\pi} \left( r|\sin \varphi| - \frac{1}{2} \log(1 + r|\sin \varphi|) + \log(1 + r) \right) d\varphi \\ &\leq C_{11} + \frac{1}{2\pi} \int_0^{2\pi} \left( r|\sin \varphi| - \frac{1}{2} \log r - \frac{1}{2} \log |\sin \varphi| + \log(1 + r) \right) d\varphi \\ &= \frac{2r}{\pi} - \frac{1}{2} \log r + \log(1 + r) + C_{12}, \quad r > 0, \end{aligned}$$

whence it follows (6). The theorem is proved.  $\square$

**Theorem 3.13.** Let  $(s_k)_{k \in \mathbb{N}}$  be a sequence of distinct nonzero complex numbers. Let  $|s_k| \leq \Delta k + \beta + \alpha_k$  for  $0 < \Delta < \pi/2$ ,  $-\Delta < \beta < 1 - \Delta(2 + \pi)/\pi$ , and the sequence  $(\alpha_k)_{k \in \mathbb{N}}$  such that  $\alpha_k \geq 0$ ,  $\alpha_k = O(1)$  as  $k \rightarrow +\infty$  and

$$\sum_{k=1}^{\infty} |\alpha_{k+1} - \alpha_k| < +\infty, \quad \sum_{k=1}^{\infty} \frac{\alpha_k}{k} < +\infty.$$

Then the system  $\{\theta_k : k \in \mathbb{N}\}$  is complete in  $L^2((0; 1); x^2 dx)$ .

*Proof.* Let  $\mu_k = \Delta k + \beta + \alpha_k$ ,  $k \in \mathbb{N}$ , and

$$n_1(t) = \sum_{\mu_k \leq t} 1, \quad N_1(r) = \int_0^r \frac{n_1(t)}{t} dt, \quad r > 0.$$

Then  $n(t) \geq n_1(t)$ ,  $N(r) \geq N_1(r)$  and  $n_1(t) = m$  for  $\Delta m + \beta + \alpha_m \leq t < \Delta(m + 1) + \beta + \alpha_{m+1}$  ( $n_1(t) = 0$  on  $(0; \mu_1)$ ). Let  $r \in [\mu_j; \mu_{j+1})$ . Then  $j = r/\Delta + O(1)$  as  $r \rightarrow +\infty$ . Therefore, under the assumptions of the theorem, by analogy with [7, p. 894] (see also [6, p. 9]), we obtain

$$\begin{aligned} N_1(r) &\geq \sum_{k=1}^{j-1} k \log \frac{\Delta(k+1) + \beta}{\Delta k + \beta} - \left| \sum_{k=1}^{j-1} k \left( \log \frac{\Delta(k+1) + \beta + \alpha_{k+1}}{\Delta k + \beta + \alpha_k} - \log \frac{\Delta(k+1) + \beta}{\Delta k + \beta} \right) \right| + O(1) \\ &\geq \frac{r}{\Delta} - \left( \frac{1}{2} + \frac{\beta}{\Delta} \right) \log r - C_{13} \sum_{k=1}^{\infty} \left( |\alpha_{k+1} - \alpha_k| + \frac{\alpha_k}{k} \right) + O(1) \\ &\geq \frac{r}{\Delta} - \left( \frac{1}{2} + \frac{\beta}{\Delta} \right) \log r + O(1), \quad r \rightarrow +\infty. \end{aligned}$$

In view of this, for  $0 < \Delta < \pi/2$  and  $-\Delta < \beta < 1 - \Delta(2 + \pi)/\pi$ , we get

$$\begin{aligned} &\limsup_{r \rightarrow +\infty} \left( N(r) - \frac{2r}{\pi} + \frac{1}{2} \log r - \log(1+r) \right) \\ &\geq \limsup_{r \rightarrow +\infty} \left( N_1(r) - \frac{2r}{\pi} + \frac{1}{2} \log r - \log(1+r) \right) \\ &\geq \limsup_{r \rightarrow +\infty} \left( \frac{r}{\Delta} - \left( \frac{1}{2} + \frac{\beta}{\Delta} \right) \log r - \frac{2r}{\pi} + \frac{1}{2} \log r - \log(1+r) + O(1) \right) \\ &\geq \limsup_{r \rightarrow +\infty} \left( r \left( \frac{1}{\Delta} - \frac{2}{\pi} \right) - \left( \frac{\beta}{\Delta} + 1 \right) \log(1+r) + O(1) \right) \\ &\geq \limsup_{r \rightarrow +\infty} \left( r \left( \frac{1}{\Delta} - \frac{2}{\pi} - \frac{\beta}{\Delta} - 1 \right) + O(1) \right) = +\infty. \end{aligned}$$

Finally, according to Theorem 3.12, we obtain the required proposition. The proof of theorem is completed.  $\square$

**References**

[1] L. D. Abreu, *Completeness, special functions and uncertainty principles over q-linear grids*, J. Phys. A: Math. Gen. **39**(47) (2006), 14567–14580.  
 [2] H. Bateman, A. Erdélyi, *Higher transcendental functions*, Vol. II, McGraw-Hill, New York-Toronto-London, 1953.  
 [3] R. P. Boas, H. Pollard, *Complete sets of Bessel and Legendre functions*, Ann. of Math. **48**(2) (1947), 366–384.  
 [4] H. Hochstadt, *The mean convergence of Fourier-Bessel series*, SIAM Rev. **9** (1967), 211–218.  
 [5] R. V. Khats', *Integral representation of one class of entire functions*, Armen. J. Math. **14**(1) (2022), 1–9.  
 [6] R. V. Khats', *On conditions of the completeness of some systems of Bessel functions in the space  $L^2((0; 1); x^{2p} dx)$* , Azerb. J. Math. **11**(1) (2021), 3–10.



- [7] R. V. Khats', *Completeness conditions of systems of Bessel functions in weighted  $L^2$ -spaces in terms of entire functions*, Turk. J. Math. **45**(2) (2021), 890–895.
- [8] R. V. Khats', *Completeness of the systems of Bessel functions of index  $-5/2$* , Carpathian Math. Publ. (2022), accepted.
- [9] R. V. Khats', *Generalized eigenvectors of linear operators and biorthogonal systems*, Constr. Math. Anal. **5**(2) (2022), 60–71.
- [10] B. Ya. Levin, *Lectures on entire functions*, In: Lyubarskii Yu., Sodin M., Tkachenko V. (Ed.) Transl. Math. Monogr. Amer. Math. Soc., Vol. 150, Providence, R.I., 1996.
- [11] A. M. Sedletskii, *Analytic Fourier transforms and exponential approximations. I.*, J. Math. Sci. **129**(6) (2005), 4251–4408.
- [12] O. V. Shavala, *On some approximation properties of the Bessel functions of order  $-5/2$* , Mat. Stud. **43**(2) (2015), 180–184. (in Ukrainian)
- [13] O. V. Shavala, *On completeness of systems of functions generated by the Bessel function*, Bukovinian Math. J. **5**(3-4) (2017), 168–171. (in Ukrainian)
- [14] H. M. Srivastava, *A survey of some recent developments on higher transcendental functions of analytic number theory and applied mathematics*, Symmetry **13** (2021), 2294.
- [15] H. M. Srivastava, *Some parametric and argument variations of the operators of fractional calculus and related special functions and integral transformations*, J. Nonlinear Convex Anal. **22** (2021), 1501–1520.
- [16] H. M. Srivastava, *Some general families of integral transformations and related results*, Appl. Math. Comput. Sci. **6**(1) (2022), 27–41.
- [17] V. S. Vladimirov, *Equations of mathematical physics*, Nauka, Moscow, 1981. (in Russian)
- [18] B. V. Vynnyts'kyi, V. M. Dilnyi, *On approximation properties of one trigonometric system*, Russ. Math. **58**(11) (2014), 10–21.
- [19] B. V. Vynnyts'kyi, R. V. Khats', *Some approximation properties of the systems of Bessel functions of index  $-3/2$* , Mat. Stud. **34**(2) (2010), 152–159.
- [20] B. V. Vynnyts'kyi, R. V. Khats', *Completeness and minimality of systems of Bessel functions*, Ufa Math. J. **5**(2) (2013), 131–141.
- [21] B. V. Vynnyts'kyi, R. V. Khats', *On the completeness and minimality of sets of Bessel functions in weighted  $L^2$ -spaces*, Eurasian Math. J. **6**(1) (2015), 123–131.
- [22] B. V. Vynnyts'kyi, R. V. Khats', *A remark on basis property of systems of Bessel and Mittag-Leffler type functions*, J. Contemp. Math. Anal. **50**(6) (2015), 300–305.
- [23] B. V. Vynnyts'kyi, R. V. Khats', *Complete biorthogonal systems of Bessel functions*, Mat. Stud. **48**(2) (2017), 150–155.
- [24] B. V. Vynnyts'kyi, R. V. Khats', I. B. Sheparovych, *Unconditional bases of systems of Bessel functions*, Eurasian Math. J. **11**(4) (2020), 76–86.
- [25] B. V. Vynnyts'kyi, O. V. Shavala, *Boundedness of solutions of a second-order linear differential equation and a boundary value problem for Bessel's equation*, Mat. Stud. **30**(1) (2008), 31–41. (in Ukrainian)
- [26] B. V. Vynnyts'kyi, O. V. Shavala, *Some properties of boundary value problems for Bessel's equation*, Math. Bull. Shevchenko Sci. Soc. **10** (2013), 169–172.
- [27] G. N. Watson, *A treatise on the theory of Bessel functions*, Cambridge University Press, Cambridge, 1944.