



# Judgement of two Weyl type theorems for bounded linear operators

Tengjie Zhang<sup>a,\*</sup>, Xiaohong Cao<sup>a</sup>

<sup>a</sup>*School of Mathematics and Statistics, Shaanxi Normal University, Xi'an, 710119, China*

**Abstract.** Let  $H$  be an infinite dimensional separable complex Hilbert space and  $B(H)$  the algebra of all bounded linear operators on  $H$ .  $T \in B(H)$  is said to satisfy property  $(UW_{\Pi})$  if  $\sigma_a(T) \setminus \sigma_{ea}(T) = \Pi(T)$ , where  $\sigma_a(T)$  and  $\sigma_{ea}(T)$  denote the approximate point spectrum and the essential approximate point spectrum of  $T$  respectively,  $\Pi(T)$  denotes the set of all poles of  $T$ .  $T \in B(H)$  satisfies a-Weyl's theorem if  $\sigma_a(T) \setminus \sigma_{ea}(T) = \pi_{00}^a(T)$ , where  $\pi_{00}^a(T) = \{\lambda \in \text{iso}\sigma_a(T) : 0 < n(T - \lambda I) < \infty\}$ . In this paper, we give necessary and sufficient conditions for a bounded linear operator and its function calculus to satisfy both property  $(UW_{\Pi})$  and a-Weyl's theorem by topological uniform descent. In addition, the property  $(UW_{\Pi})$  and a-Weyl's theorem under perturbations are also discussed.

## 1. Introduction and preliminaries

Throughout this paper,  $\mathbb{C}$  and  $\mathbb{N}$  denote the set of complex numbers and the set of nonnegative integers. The unit closed disk and unit circle on the complex plane  $\mathbb{C}$  are denoted by  $\mathbb{D}$  and  $\Gamma$ , respectively. Let  $H$  be a complex separable infinite dimensional Hilbert space and  $B(H)$  the algebra of all bounded linear operators on  $H$ . Let  $T \in B(H)$ . We denote by  $n(T)$  the dimension of the kernel  $N(T)$  and by  $d(T)$  the codimension of the range  $R(T)$ . If  $R(T)$  is closed and  $n(T) < \infty$ , then  $T$  is called an upper semi-Fredholm operator.  $T$  is said to be a lower semi-Fredholm operator if  $d(T) < \infty$ . An operator  $T$  is said to be Fredholm operator if it is both lower and upper semi-Fredholm. Especially, if  $T$  is an upper semi-Fredholm operator and  $n(T) = 0$ , then  $T$  is called a bounded below operator. The index of  $T$  is defined by  $\text{ind}(T) = n(T) - d(T)$ . An operator  $T$  is said to be an upper semi-Weyl operator if it is an upper semi-Fredholm operator with  $\text{ind}(T) \leq 0$ . If  $T$  is an upper semi-Fredholm operator and  $\text{ind}(T) = 0$ , then  $T$  is called Weyl operator. The spectrum of  $T$ , the approximate point spectrum  $\sigma_a(T)$ , the essential approximate point spectrum  $\sigma_{ea}(T)$ , the upper semi-Fredholm spectrum  $\sigma_{SF_+}(T)$  are defined by

$$\begin{aligned}\sigma(T) &= \{\lambda \in \mathbb{C} : T - \lambda I \text{ is not invertible}\}, \\ \sigma_a(T) &= \{\lambda \in \mathbb{C} : T - \lambda I \text{ is not a bounded below operator}\}, \\ \sigma_{ea}(T) &= \{\lambda \in \mathbb{C} : T - \lambda I \text{ is not an upper semi-Weyl operator}\}, \\ \sigma_{SF_+}(T) &= \{\lambda \in \mathbb{C} : T - \lambda I \text{ is not an upper semi-Fredholm operator}\}.\end{aligned}$$

The ascent and descent of  $T$  are defined by  $\text{asc}(T) = \inf\{n \in \mathbb{N} : N(T^n) = N(T^{n+1})\}$  and  $\text{des}(T) = \inf\{n \in \mathbb{N} : R(T^n) = R(T^{n+1})\}$ . If the infimum does not exist, then we write  $\text{asc}(T) = \infty$  (resp.  $\text{des}(T) = \infty$ ). If

2020 *Mathematics Subject Classification.* Primary 47A10; Secondary 47A53, 47A55

*Keywords.* property  $(UW_{\Pi})$ ; a-Weyl's theorem; topological uniform descent; perturbation.

Received: 22 March 2023; Accepted: 06 April 2023

Communicated by Dragan S. Djordjević

\* Corresponding author: Tengjie Zhang

*Email addresses:* zhangtengjie123@snnu.edu.cn (Tengjie Zhang), xiaohongcao@snnu.edu.cn (Xiaohong Cao)

$\text{asc}(T) = \text{des}(T) < \infty$ , then  $T$  is Drazin invertible.  $T$  is called a Browder operator if  $T$  is both Fredholm operator and Drazin invertible. The Drazin spectrum  $\sigma_D(T)$ , the left Browder spectrum  $\sigma_{ab}(T)$  and the Browder spectrum  $\sigma_b(T)$  are defined by

$$\begin{aligned} \sigma_D(T) &= \{\lambda \in \mathbb{C} : T - \lambda I \text{ is not Drazin invertible}\}, \\ \sigma_{ab}(T) &= \{\lambda \in \mathbb{C} : T - \lambda I \text{ is not upper semi-Fredholm or } \text{asc}(T - \lambda I) = \infty\}, \\ \sigma_b(T) &= \{\lambda \in \mathbb{C} : T - \lambda I \text{ is not a Browder operator}\}. \end{aligned}$$

Let  $\rho(T) = \mathbb{C} \setminus \sigma(T)$ ,  $\rho_a(T) = \mathbb{C} \setminus \sigma_a(T)$ ,  $\rho_b(T) = \mathbb{C} \setminus \sigma_b(T)$ . We denote by  $\sigma_0(T)$  the set of all normal eigenvalues of  $T$ , thus  $\sigma_0(T) = \sigma(T) \setminus \sigma_b(T)$ . For a set  $E \subseteq \mathbb{C}$ , we write  $\text{iso}E$ ,  $\text{acc}E$  and  $\partial E$  as the set of isolated points, accumulation points and boundary points of  $E$ .

For a Cauchy domain  $\Omega$ , if all the curves of  $\partial\Omega$  are regular analytic Jordan curves, we say that  $\Omega$  is an analytic Cauchy domain. For  $T \in B(H)$ , if  $\sigma$  is a clopen subset of  $\sigma(T)$ , then there exists an analytic Cauchy domain  $\Omega$  such that  $\sigma \subseteq \Omega$  and  $[\sigma(T) \setminus \sigma] \cap \bar{\Omega} = \emptyset$ , where  $\bar{\Omega}$  is the closure of  $\Omega$ . We denote by  $E(\sigma; T)$  the Riesz idempotent of  $T$  corresponding to  $\sigma$ , i.e.,

$$E(\sigma; T) = \frac{1}{2\pi i} \int_{\Gamma} (\lambda I - T)^{-1} d\lambda,$$

where  $\Gamma = \partial\Omega$  is positively oriented with respect to  $\Omega$  in the sense of complex variable theory. In this case, we have  $H(\sigma; T) = R(E(\sigma; T))$ . Clearly, if  $\lambda \in \text{iso}\sigma(T)$ , then  $\{\lambda\}$  is a clopen subset of  $\sigma(T)$ . We write  $H(\lambda; T)$  instead of  $H(\{\lambda\}; T)$ ; if in addition,  $\dim H(\lambda; T) < \infty$ , then  $\lambda \in \sigma_0(T)$ .

Spectral theory of operators is an important part of operator theory. Weyl's theorem, as an important conclusion in spectral theory, is discovered by H.Weyl in 1909 ([16]) when he studied the spectral set of self-adjoint operators on Hilbert spaces. As one of the research focuses of spectral theory in recent years, scholars have made various modifications to it.

The variation of Weyl's theorem, namely, a-Weyl's theorem ([13, 14]) were given by Rakočević. We say that the a-Weyl's theorem holds for  $T$  if

$$\sigma_a(T) \setminus \sigma_{ea}(T) = \pi_{00}^a(T),$$

where  $\pi_{00}^a(T) = \{\lambda \in \text{iso}\sigma_a(T) : 0 < n(T - \lambda I) < \infty\}$ .

Property  $(UW_{\Pi})$ , as well as a-Weyl's theorem, is also a variant of Weyl's theorem. In [6], Berkani and Kachad introduced the definition of property  $(UW_{\Pi})$ .  $T \in B(H)$  satisfies property  $(UW_{\Pi})$  and denoted by  $T \in (UW_{\Pi})$ , if

$$\sigma_a(T) \setminus \sigma_{ea}(T) = \Pi(T),$$

where  $\Pi(T) = \sigma(T) \setminus \sigma_D(T)$ . If  $\lambda \in \Pi(T)$ , then  $\lambda$  is a pole of  $T$ .

The concept of topological uniform descent was first proposed by Sandy Grabiner ([9]). The introduction of this concept provides a new tool for the study of operator theory, and many scholars have achieved corresponding research results by using topological uniform descent ([8, 11, 15]). If  $T \in B(H)$ , then for each nonnegative integer  $n$ ,  $T$  induces a linear transformation

$$\Gamma_n : R(T^n)/R(T^{n+1}) \longrightarrow R(T^{n+1})/R(T^{n+2}),$$

we will let  $k_n(T)$  be the dimension of the null space of the induced map and let  $k(T) = \sum_{n=0}^{\infty} k_n(T)$ . The operator range topology on  $R(T^n)$  is defined by the norm  $\|y\|_n = \inf\{\|x\|, x \in H, y = T^n x\}$ . If there is a nonnegative integer  $d$  for which  $k_n(T) = 0$  for  $n \geq d$  and  $R(T^n)$  is closed in the operator range topology of  $R(T^d)$  for  $n \geq d$ , then we say that  $T$  has topological uniform descent.

It can be shown that if  $T$  is semi-Fredholm, then  $T$  has topological uniform descent. If  $T - \lambda I$  has topological uniform descent and  $\lambda \in \partial\sigma(T)$ , then  $\lambda \in \Pi(T)$  ([9, Corollary 4.9]). The topological uniform descent spectrum of  $T$  is defined by

$$\sigma_{\tau}(T) = \{\lambda \in \mathbb{C} : T - \lambda I \text{ has not topological uniform descent}\},$$

and  $\rho_{\tau}(T) = \mathbb{C} \setminus \sigma_{\tau}(T)$ .

**Example 1.1.** (i) It is easy to see that if  $T \in (UW_{\Pi})$ , then  $\sigma_a(T) \setminus \sigma_{ea}(T) \subseteq \pi_{00}^a(T)$ . But property  $(UW_{\Pi})$  does not imply a-Weyl's theorem. Let  $A, B \in B(\ell^2)$  be defined by

$$A(x_1, x_2, \dots) = (0, x_1, x_2, \dots), B(x_1, x_2, \dots) = (0, 0, \frac{x_2}{2}, \frac{x_3}{3}, \dots).$$

Put  $T = \begin{pmatrix} A & 0 \\ 0 & B \end{pmatrix}$ . Then we have  $\sigma(T) = \mathbb{D}$ ,  $\Pi(T) = \emptyset$ ,  $\sigma_a(T) = \sigma_{ea}(T) = \{0\} \cup \Gamma$ ,  $\pi_{00}^a(T) = \{0\}$ . It follows that  $T \in (UW_{\Pi})$ , but the a-Weyl's theorem does not hold for  $T$ .

(ii) A-Weyl's theorem does not imply property  $(UW_{\Pi})$ . Let  $A, B \in B(\ell^2)$  be defined by

$$A(x_1, x_2, \dots) = (0, x_1, x_2, \dots), B(x_1, x_2, \dots) = (0, x_2, x_3, \dots).$$

Put  $T = \begin{pmatrix} A & 0 \\ 0 & B \end{pmatrix}$ . Then we have  $\sigma(T) = \mathbb{D}$ ,  $\Pi(T) = \emptyset$ ,  $\sigma_a(T) = \{0\} \cup \Gamma$ ,  $\sigma_{ea}(T) = \Gamma$ ,  $\pi_{00}^a(T) = \{0\}$ . It follows that  $T$  satisfies a-Weyl's theorem, but  $T \notin (UW_{\Pi})$ .

(iii) There exists  $T \in B(H)$  such that neither property  $(UW_{\Pi})$  nor a-Weyl's theorem holds for  $T$ . Let  $A, B \in B(\ell^2)$  be defined by

$$A(x_1, x_2, \dots) = (0, x_2, 0, x_4, \dots), B(x_1, x_2, \dots) = (0, 0, \frac{x_2}{2}, \frac{x_3}{3}, \dots).$$

Put  $T = \begin{pmatrix} A & 0 \\ 0 & B - I \end{pmatrix}$ . Then we have  $\sigma(T) = \{-1, 0, 1\}$ ,  $\Pi(T) = \{0, 1\}$ ,  $\sigma_a(T) = \sigma_{ea}(T) = \{-1, 0, 1\}$ ,  $\pi_{00}^a(T) = \{-1\}$ . Thus, neither property  $(UW_{\Pi})$  nor a-Weyl's theorem holds for  $T$ .

We have seen in Example 1.1 that there is no relationship between  $T \in (UW_{\Pi})$  and  $T$  satisfies a-Weyl's theorem although the forms of property  $(UW_{\Pi})$  and a-Weyl's theorem are similar.

In this paper, we will give necessary and sufficient conditions for bounded linear operators to satisfy both property  $(UW_{\Pi})$  and a-Weyl's theorem by topological uniform descent in section 2. What's more, we also discuss both property  $(UW_{\Pi})$  and a-Weyl's theorem under quasi-nilpotent perturbation for bounded linear operators. In section 3, we will talk about operator functions to satisfy both property  $(UW_{\Pi})$  and a-Weyl's theorem in terms of topological uniform descent. In addition, we also discuss the case that Drazin invertible operators satisfy both property  $(UW_{\Pi})$  and a-Weyl's theorem.

## 2. Property $(UW_{\Pi})$ and a-Weyl's theorem of bounded linear operators

In this section, we will describe both property  $(UW_{\Pi})$  and a-Weyl's theorem hold for  $T$  by means of the property of topological uniform descent.

**Theorem 2.1.** Let  $T \in B(H)$ . The following statements are equivalent:

- (1)  $T$  satisfies both the property  $(UW_{\Pi})$  and a-Weyl's theorem;
- (2)  $\sigma_b(T) = [\sigma_{\tau}(T) \cap \{\lambda \in \mathbb{C} : n(T - \lambda I) = d(T - \lambda I)\}] \cup \{\lambda \in \sigma(T) : n(T - \lambda I) = 0\} \cup [\text{acc}\sigma_a(T) \cap \sigma_{ea}(T)] \cup \{\lambda \in \mathbb{C} : n(T - \lambda I) = \infty\} \cap \text{acc}[\rho_a(T) \cap \sigma(T)]$ .

*Proof.* (1)  $\Rightarrow$  (2). The inclusion " $\supseteq$ " is obvious. For the opposite inclusion, take arbitrarily  $\lambda_0$  that does not belong to the right side of (2). Without loss of generality, suppose that  $\lambda_0 \in \sigma(T)$ . Then we have  $n(T - \lambda_0 I) > 0$ .

**Case 1** Suppose that  $\lambda_0 \notin \sigma_{ea}(T)$ . Then  $\lambda_0 \in \sigma_a(T) \setminus \sigma_{ea}(T)$ . Since  $T \in (UW_{\Pi})$ , we have  $\lambda_0 \notin \sigma_b(T)$ .

**Case 2** Suppose that  $\lambda_0 \notin \text{acc}\sigma_a(T) \cup \{\lambda \in \mathbb{C} : n(T - \lambda I) = \infty\}$ . Then  $\lambda_0 \in \pi_{00}^a(T)$ . Since  $T$  satisfies both property  $(UW_{\Pi})$  and a-Weyl's theorem, we can get that  $\lambda_0 \notin \sigma_b(T)$ .

**Case 3** Suppose that  $\lambda_0 \notin \sigma_{\tau}(T) \cup \text{acc}\sigma_a(T) \cup \text{acc}[\rho_a(T) \cap \sigma(T)]$ . Then  $\lambda_0 \in \rho_{\tau}(T) \cap \partial\sigma(T)$ , we can get  $\lambda_0 \in \Pi(T)$  ([9, Corollary 4.9]). From  $T \in (UW_{\Pi})$  we get that  $\lambda_0 \notin \sigma_b(T)$ .

**Case 4** Suppose that  $\lambda_0 \notin \{\lambda \in \mathbb{C} : n(T - \lambda I) = d(T - \lambda I)\} \cup \text{acc}\sigma_a(T) \cup \text{acc}[\rho_a(T) \cap \sigma(T)]$ . We have  $0 < n(T - \lambda_0 I) < \infty$ , thus  $\lambda_0 \in \pi_{00}^a(T)$ . Since  $T$  satisfies both property  $(UW_{\Pi})$  and a-Weyl's theorem, we get that  $\lambda_0 \notin \sigma_b(T)$ .

(2)  $\Rightarrow$  (1). It is clear that  $\{[\sigma_a(T)\setminus\sigma_{ea}(T)] \cup \Pi(T) \cup \pi_{00}^a(T)\} \cap [\sigma_\tau(T) \cap \{\lambda \in \mathbb{C} : n(T - \lambda I) = d(T - \lambda I)\}] = \emptyset$ ,  $\{[\sigma_a(T)\setminus\sigma_{ea}(T)] \cup \Pi(T) \cup \pi_{00}^a(T)\} \cap \{\lambda \in \sigma(T) : n(T - \lambda I) = 0\} = \emptyset$ ,  $\{[\sigma_a(T)\setminus\sigma_{ea}(T)] \cup \Pi(T) \cup \pi_{00}^a(T)\} \cap [\text{acc}\sigma_a(T) \cap \sigma_{ea}(T)] = \emptyset$ ,  $\{[\sigma_a(T)\setminus\sigma_{ea}(T)] \cup \Pi(T) \cup \pi_{00}^a(T)\} \cap \{\lambda \in \mathbb{C} : n(T - \lambda I) = \infty\} \cap \text{acc}[\rho_a(T) \cap \sigma(T)] = \emptyset$ . Hence  $[\sigma_a(T)\setminus\sigma_{ea}(T)] \cup \Pi(T) \cup \pi_{00}^a(T) = \sigma_0(T)$ . It follows that  $T$  satisfies both property  $(UW_\Pi)$  and a-Weyl's theorem.  $\square$

**Remark 2.2.** (i) In Theorem 2.1, suppose  $T \in B(H)$  satisfies both property  $(UW_\Pi)$  and a-Weyl's theorem, then each part of the decomposition of  $\sigma_b(T)$  can not be deleted.

(a) Let  $A, B \in B(\ell^2)$  be defined by

$$A(x_1, x_2, \dots) = (0, x_1, 0, \frac{x_3}{3}, 0, \frac{x_5}{5}, \dots), B(x_1, x_2, \dots) = (0, 0, \frac{x_2}{2}, \frac{x_3}{3}, \dots).$$

Put  $T = \begin{pmatrix} A & 0 \\ 0 & B \end{pmatrix}$ . Then we have  $\sigma(T) = \{0\}$ ,  $\Pi(T) = \emptyset$ ,  $\sigma_a(T) = \sigma_{ea}(T) = \{0\}$ ,  $\pi_{00}^a(T) = \emptyset$ . Hence  $T$  satisfies both property  $(UW_\Pi)$  and a-Weyl's theorem. But  $\{\lambda \in \sigma(T) : n(T - \lambda I) = 0\} = [\text{acc}\sigma_a(T) \cap \sigma_{ea}(T)] = \{\{\lambda \in \mathbb{C} : n(T - \lambda I) = \infty\} \cap \text{acc}[\rho_a(T) \cap \sigma(T)]\} = \emptyset$ . Thus  $\sigma_\tau(T) \cap \{\lambda \in \mathbb{C} : n(T - \lambda I) = d(T - \lambda I)\}$  cannot be deleted.

(b) Let  $T \in B(\ell^2)$  be defined by

$$T(x_1, x_2, \dots) = (0, x_1, x_2, \dots).$$

Then we have  $\sigma(T) = \mathbb{D}$ ,  $\Pi(T) = \emptyset$ ,  $\sigma_a(T) = \sigma_{ea}(T) = \Gamma$ ,  $\pi_{00}^a(T) = \emptyset$ , and so  $T$  satisfies both property  $(UW_\Pi)$  and a-Weyl's theorem. But  $\sigma_b(T) \neq [\sigma_\tau(T) \cap \{\lambda \in \mathbb{C} : n(T - \lambda I) = d(T - \lambda I)\}] \cup [\text{acc}\sigma_a(T) \cap \sigma_{ea}(T)] \cup \{\{\lambda \in \mathbb{C} : n(T - \lambda I) = \infty\} \cap \text{acc}[\rho_a(T) \cap \sigma(T)]\}$ . It follows that  $\{\lambda \in \sigma(T) : n(T - \lambda I) = 0\}$  cannot be deleted.

(c) Let  $T \in B(\ell^2)$  be defined by

$$T(x_1, x_2, \dots) = (0, \frac{x_2}{2}, \frac{x_3}{3}, \dots).$$

Then we have  $\sigma(T) = \sigma_a(T) = \{0, \frac{1}{2}, \frac{1}{3}, \frac{1}{4}, \dots\}$ ,  $\sigma_{ea}(T) = \{0\}$ ,  $\Pi(T) = \pi_{00}^a(T) = \{\frac{1}{2}, \frac{1}{3}, \frac{1}{4}, \dots\}$ . So property  $(UW_\Pi)$  and a-Weyl's theorem hold for  $T$ . But  $\sigma_b(T) \neq [\sigma_\tau(T) \cap \{\lambda \in \mathbb{C} : n(T - \lambda I) = d(T - \lambda I)\}] \cup \{\{\lambda \in \sigma(T) : n(T - \lambda I) = 0\} \cup \{\{\lambda \in \mathbb{C} : n(T - \lambda I) = \infty\} \cap \text{acc}[\rho_a(T) \cap \sigma(T)]\}\}$ . Thus  $\text{acc}\sigma_a(T) \cap \sigma_{ea}(T)$  cannot be deleted.

(d) Let  $A, B \in B(\ell^2)$  be defined by

$$A(x_1, x_2, \dots) = (0, x_1, x_2, \dots), B(x_1, x_2, \dots) = (0, x_2, 0, x_4, \dots).$$

Put  $T = \begin{pmatrix} A & 0 \\ 0 & B \end{pmatrix}$ . Then we have  $\sigma(T) = \mathbb{D}$ ,  $\Pi(T) = \emptyset$ ,  $\sigma_a(T) = \sigma_{ea}(T) = \{0\} \cup \Gamma$ ,  $\pi_{00}^a(T) = \emptyset$ . It follows that  $T$  satisfies both property  $(UW_\Pi)$  and a-Weyl's theorem. However,  $\sigma_b(T) \neq [\sigma_\tau(T) \cap \{\lambda \in \mathbb{C} : n(T - \lambda I) = d(T - \lambda I)\}] \cup \{\{\lambda \in \sigma(T) : n(T - \lambda I) = 0\} \cup [\text{acc}\sigma_a(T) \cap \sigma_{ea}(T)]\}$ , which means that  $\{\lambda \in \mathbb{C} : n(T - \lambda I) = \infty\} \cap \text{acc}[\rho_a(T) \cap \sigma(T)]$  cannot be deleted.

(ii) It is clear that  $\sigma_{ea}(T) = \sigma_{SF_+}(T) \cup \{\lambda \in \mathbb{C} : n(T - \lambda I) > d(T - \lambda I)\}$ . From Theorem 2.1, we can get that  $T$  satisfies both property  $(UW_\Pi)$  and a-Weyl's theorem if and only if  $\sigma_b(T) = [\sigma_\tau(T) \cap \{\lambda \in \mathbb{C} : n(T - \lambda I) = d(T - \lambda I)\}] \cup \{\{\lambda \in \sigma(T) : n(T - \lambda I) = 0\} \cup [\text{acc}\sigma_a(T) \cap \sigma_{SF_+}(T)] \cup \{\{\lambda \in \mathbb{C} : n(T - \lambda I) = \infty\} \cap \text{acc}[\rho_a(T) \cap \sigma(T)]\} \cup \{\lambda \in \mathbb{C} : n(T - \lambda I) > d(T - \lambda I)\}$ .

(iii) If  $\sigma_\tau(T) = \emptyset$ , we claim that  $\text{int}\sigma(T) = \emptyset$ . If not, there exists a continuous curve segment  $L \subseteq \partial\sigma(T)$ . Take  $\lambda_0 \in L$ , from  $\sigma_\tau(T) = \emptyset$  we can get that  $\lambda_0 \in \Pi(T)$ . Then  $\lambda_0 \in \text{iso}\sigma(T)$ , a contradiction. Thus,  $\sigma(T) = \partial\sigma(T)$ . Take arbitrarily  $\lambda \in \sigma(T) = \partial\sigma(T)$ . It follows from  $\lambda \notin \sigma_\tau(T)$  that  $\lambda \in \Pi(T)$  and  $\lambda \in \text{iso}\sigma(T)$ . Since  $\sigma(T)$  is a bounded set, we can get that  $\sigma(T)$  consists of finite points. Therefore, if  $\sigma_\tau(T) = \emptyset$ , then  $\sigma(T) = \Pi(T)$ .

From Theorem 2.1, we can obtain this result: If  $\sigma_\tau(T) = \emptyset$  and  $T$  satisfies both property  $(UW_\Pi)$  and a-Weyl's theorem (or only property  $(UW_\Pi)$  is required), then  $\sigma_b(T) = \emptyset$ , a contradiction with the fact that  $\sigma_b(T)$  is nonempty. Therefore, if  $T$  satisfies both property  $(UW_\Pi)$  and a-Weyl's theorem (or only property  $(UW_\Pi)$ ),  $\sigma_\tau(T) \neq \emptyset$ .

**Corollary 2.3.** Let  $T \in B(H)$ . The following statements are equivalent:

- (1)  $T$  satisfies both property  $(UW_\Pi)$  and a-Weyl's theorem;
- (2)  $\sigma_b(T) = [\sigma_\tau(T) \cap \{\lambda \in \mathbb{C} : n(T - \lambda I) = \infty\}] \cup \{\lambda \in \sigma(T) : n(T - \lambda I) = 0\} \cup [\text{acc}\sigma_a(T) \cap \sigma_{ea}(T)] \cup \{\{\lambda \in \mathbb{C} : n(T - \lambda I) = d(T - \lambda I)\} \cap \text{acc}[\rho_a(T) \cap \sigma(T)]\}$ .

*Proof.* (1)  $\Rightarrow$  (2). The inclusion “ $\supseteq$ ” is obvious. For the opposite inclusion, we know that  $\sigma_\tau(T) \cap \{\lambda \in \mathbb{C} : n(T - \lambda I) = d(T - \lambda I) < \infty\} = \emptyset$ . Hence  $[\sigma_\tau(T) \cap \{\lambda \in \mathbb{C} : n(T - \lambda I) = d(T - \lambda I)\}] = [\sigma_\tau(T) \cap \{\lambda \in \mathbb{C} : n(T - \lambda I) = d(T - \lambda I) = \infty\}] \subseteq \sigma_\tau(T) \cap \{\lambda \in \mathbb{C} : n(T - \lambda I) = \infty\}$ . From  $\{\lambda \in \mathbb{C} : n(T - \lambda I) = \infty, d(T - \lambda I) < \infty\} \cap \text{acc}[\rho_a(T) \cap \sigma(T)] = \emptyset$  we can get that  $\{\lambda \in \mathbb{C} : n(T - \lambda I) = \infty\} \cap \text{acc}[\rho_a(T) \cap \sigma(T)] = \{\{\lambda \in \mathbb{C} : n(T - \lambda I) = d(T - \lambda I) = \infty\} \cap \text{acc}[\rho_a(T) \cap \sigma(T)]\} \subseteq \{\lambda \in \mathbb{C} : n(T - \lambda I) = d(T - \lambda I)\} \cap \text{acc}[\rho_a(T) \cap \sigma(T)]$ . According to Theorem 2.1, the inclusion “ $\subseteq$ ” is obvious.

(2)  $\Rightarrow$  (1). Similar to the proof of Theorem 2.1, this result is trivial.  $\square$

It is easy to get that  $[\rho_a(T) \cap \sigma(T)] \subseteq \{\lambda \in \mathbb{C} : n(T - \lambda I) < d(T - \lambda I)\}$  and  $\{\lambda \in \mathbb{C} : n(T - \lambda I) > d(T - \lambda I)\} \subseteq \text{acc}\{\lambda \in \mathbb{C} : n(T - \lambda I) > d(T - \lambda I)\}$ . From (ii) in Remark 2.2 we obtain the following corollary.

**Corollary 2.4.** *Let  $T \in B(H)$ . The following statements are equivalent:*

- (1)  $T$  satisfies both property  $(UW_\Pi)$  and  $a$ -Weyl’s theorem;
- (2)  $\sigma_b(T) = [\sigma_\tau(T) \cap \{\lambda \in \mathbb{C} : n(T - \lambda I) = d(T - \lambda I)\}] \cup \{\lambda \in \sigma(T) : n(T - \lambda I) = 0\} \cup [\text{acc}\sigma_a(T) \cap \sigma_{SF_+}(T)] \cup \{\{\lambda \in \mathbb{C} : n(T - \lambda I) = \infty\} \cap \text{acc}\{\lambda \in \mathbb{C} : n(T - \lambda I) < d(T - \lambda I)\}\} \cup \text{acc}\{\lambda \in \mathbb{C} : n(T - \lambda I) > d(T - \lambda I)\}$ .

**Corollary 2.5.** *Let  $T \in B(H)$ . The following statements are equivalent:*

- (1)  $T$  satisfies both property  $(UW_\Pi)$  and  $a$ -Weyl’s theorem;
- (2)  $\pi_{00}^a(T) \subseteq \rho_\tau(T) \subseteq \rho_b(T) \cup \{\lambda \in \sigma(T) : n(T - \lambda I) = 0\} \cup [\text{acc}\sigma_a(T) \cap \sigma_{SF_+}(T)] \cup \{\{\lambda \in \mathbb{C} : n(T - \lambda I) = \infty\} \cap \text{acc}\{\lambda \in \mathbb{C} : n(T - \lambda I) < d(T - \lambda I)\}\} \cup \text{acc}\{\lambda \in \mathbb{C} : n(T - \lambda I) > d(T - \lambda I)\}$ .

*Proof.* (1)  $\Rightarrow$  (2). It is obvious that  $\pi_{00}^a(T) \subseteq \rho_\tau(T)$ . Suppose that  $\lambda_0 \in \rho_\tau(T)$ . Then  $\lambda_0 \notin [\sigma_\tau(T) \cap \{\lambda \in \mathbb{C} : n(T - \lambda I) = d(T - \lambda I)\}]$ . If  $\lambda_0 \notin \{\lambda \in \sigma(T) : n(T - \lambda I) = 0\} \cup [\text{acc}\sigma_a(T) \cap \sigma_{SF_+}(T)] \cup \{\{\lambda \in \mathbb{C} : n(T - \lambda I) = \infty\} \cap \text{acc}\{\lambda \in \mathbb{C} : n(T - \lambda I) < d(T - \lambda I)\}\} \cup \text{acc}\{\lambda \in \mathbb{C} : n(T - \lambda I) > d(T - \lambda I)\}$ , from Corollary 2.4 we can get that  $\lambda_0 \in \rho_b(T)$ .

(2)  $\Rightarrow$  (1). It is clear that  $\{[\sigma_a(T) \setminus \sigma_{ea}(T)] \cup \Pi(T)\} \subseteq \rho_\tau(T)$ . From Corollary 2.4 and the proof of Theorem 2.1, we have  $\{[\sigma_a(T) \setminus \sigma_{ea}(T)] \cup \Pi(T)\} \subseteq \sigma_0(T)$  and  $\pi_{00}^a(T) \subseteq \sigma_0(T)$ . Thus  $T$  satisfies both property  $(UW_\Pi)$  and  $a$ -Weyl’s theorem.  $\square$

Weyl type Theorem and its perturbation problems have attracted extensive attention in recent years ([7, 10, 17]). In the following, we will discuss quasi-nilpotent perturbation of both property  $(UW_\Pi)$  and  $a$ -Weyl’s theorem.

We call  $R \in B(H)$  is Riesz operator if  $R - \lambda I$  is Fredholm operator for every nonzero  $\lambda$ . In [3, Theorem 4.7], we have that

$$\sigma_*(T) = \sigma_*(T + R)$$

for every Riesz operator  $R$  commuting with  $T \in B(H)$ , where  $*$   $\in \{ea, ab, b\}$ . It is clear that quasi-nilpotent operators are Riesz operators.  $T \in B(H)$  is said to be  $a$ -isoloid operator if  $\text{iso}\sigma_a(T) \subseteq \sigma_p(T)$ , where  $\sigma_p(T) = \{\lambda \in \mathbb{C} : n(T - \lambda I) > 0\}$ . If  $\text{iso}\sigma_a(T) \subseteq \Pi(T)$ , then  $T$  is called  $a$ -polaroid operator.

**Example 2.6.** (1) Let  $T, Q \in B(\ell^2)$  be defined by

$$T(x_1, x_2, x_3, \dots) = (0, x_1, x_2, x_3, \dots), Q(x_1, x_2, x_3, \dots) = (0, -x_1, 0, 0, \dots).$$

We have that  $T$  satisfies both property  $(UW_\Pi)$  and  $a$ -Weyl’s theorem. However,  $\sigma(T + Q) = \sigma_a(T + Q) = \mathbb{D}$ ,  $\sigma_{ea}(T) = \Gamma$ ,  $\Pi(T) = \pi_{00}^a(T) = \emptyset$ . It follows that both property  $(UW_\Pi)$  and  $a$ -Weyl’s theorem don’t hold for  $T + Q$ .

(2) Let  $A, B \in B(\ell^2)$  be defined by

$$A(x_1, x_2, \dots) = (0, x_1, 0, \frac{x_3}{3}, 0, \frac{x_5}{5}, \dots), B(x_1, x_2, \dots) = (0, 0, \frac{x_2}{2}, \frac{x_3}{3}, \dots).$$

Put  $T = \begin{pmatrix} A & 0 \\ 0 & B \end{pmatrix}$ ,  $Q = \begin{pmatrix} 0 & 0 \\ 0 & -B \end{pmatrix}$ . Then we have  $QT = TQ$ ,  $T$  is  $a$ -isoloid operator,  $\sigma(T) = \sigma_a(T) = \sigma_{ea}(T) = \{0\}$ ,  $\pi_{00}^a(T) = \Pi(T) = \emptyset$ . It follows that  $T$  satisfies both property  $(UW_\Pi)$  and  $a$ -Weyl’s theorem, but we can see that  $T + Q \notin (UW_\Pi)$ .

From Example 2.6 we know that the commutativity of  $T$  is indispensable, and we can't also induce that  $T + Q$  satisfies both property  $(UW_{\Pi})$  and a-Weyl's theorem if  $T$  is a-isoloid operator. Now, let  $Q$  be a quasi-nilpotent operator with  $QT = TQ$ . For  $T \in B(H)$ , the quasi-nilpotent part of  $T$  is defined by

$$H_0(T) = \{x \in H : \lim_{n \rightarrow \infty} \|T^n x\|^{\frac{1}{n}} = 0\}.$$

It is known that  $T$  is a quasi-nilpotent operator if and only if  $H_0(T) = H$ . Thus we have the following lemma.

**Lemma 2.7.** [3, Theorem 4.9] *Let  $T \in B(H)$  and  $Q$  a quasi-nilpotent operator with  $QT = TQ$ , then  $\sigma(T) = \sigma(T + Q)$  and  $\sigma_a(T) = \sigma_a(T + Q)$ .*

*Proof.* Since  $-Q$  is quasi-nilpotent operator, we only need  $T + Q$  is bounded below if  $T$  is bounded below. We claim that  $H_0(T) = \{0\}$  if  $T$  is bounded below. In fact, since  $T$  is bounded below, there exists  $k > 0$  such that  $\|Tx\| \geq k\|x\|, \forall x \in H$ . Suppose that  $x_0 \in H_0(T)$ , then  $\lim_{n \rightarrow \infty} \|T^n x_0\|^{\frac{1}{n}} = 0$  and  $\|T^n x_0\| \geq k^n \|x_0\|$ . It follows that  $\|T^n x_0\|^{\frac{1}{n}} \geq k\|x_0\|^{\frac{1}{n}}$ . Thus,  $x_0 = 0$ . Since  $T + Q$  is upper semi-Weyl operator, we only need  $N(T + Q) = \{0\}$ . For all  $x \in N(T + Q)$  we have that  $Qx = -Tx$ . Thus  $Q^n x_0 = (-1)^n T^n x_0$ . From  $Q$  is quasi-nilpotent operator we can get that  $\lim_{n \rightarrow \infty} \|Q^n x\|^{\frac{1}{n}} = 0$ , so  $\lim_{n \rightarrow \infty} \|T^n x\|^{\frac{1}{n}} = 0$ . It follows that  $x \in H_0(T)$ . Since  $T$  is bounded below,  $H_0(T) = \{0\}$ . So,  $x = 0$ . Hence,  $T + Q$  is bounded below.

If  $T$  is invertible, we know that  $T + Q$  is Weyl operator. From  $T + Q$  is bounded below we can get that  $T + Q$  is invertible.  $\square$

**Theorem 2.8.** *Let  $T \in B(H)$  and  $Q$  a quasi-nilpotent operator with  $QT = TQ$ . Then the following statements are equivalent:*

- (1)  $T$  satisfies both property  $(UW_{\Pi})$  and a-Weyl's theorem, and  $T$  is a-polaroid operator;
- (2)  $T + Q$  satisfies both property  $(UW_{\Pi})$  and a-Weyl's theorem, and  $T + Q$  is a-polaroid operator.

*Proof.* Since  $-Q$  is quasi-nilpotent operator, we only need to show (1)  $\Rightarrow$  (2). Let  $\lambda \in \sigma_a(T + Q) \setminus \sigma_{ea}(T + Q)$ , from Lemma 2.7 we can get that  $\lambda \in \sigma_a(T) \setminus \sigma_{ea}(T)$ . It follows from  $T \in (UW_{\Pi})$  that  $\lambda \in \sigma_0(T)$  and so  $\lambda \in \sigma_0(T + Q)$ . Let  $\lambda_0 \in \Pi(T + Q)$ , then  $\lambda_0 \in \text{iso}\sigma(T + Q)$ . From Lemma 2.7 and  $T$  is a-polaroid operator we have  $\lambda_0 \in \Pi(T)$ . By  $T \in (UW_{\Pi})$  we can get  $\lambda_0 \in \sigma_0(T)$ . Then  $\lambda_0 \in \sigma_0(T + Q)$ . Let  $\mu_0 \in \pi_{00}^a(T + Q)$ , from Lemma 2.7 and  $T$  is a-polaroid operator, we get that  $\mu_0 \in \Pi(T)$ . By  $T \in (UW_{\Pi})$  we can get  $\mu_0 \in \sigma_0(T)$  and  $\mu_0 \in \sigma_0(T + Q)$ . Let  $\mu \in \text{iso}\sigma_a(T + Q)$ . Similar to the above proof, it is clear that  $\mu \in \Pi(T + Q)$ . Thus,  $T + Q \in (UW_{\Pi})$  and satisfies a-Weyl's theorem, and  $T + Q$  is a-polaroid operator.  $\square$

In the following, we will discuss the quasi-nilpotent perturbation of both property  $(UW_{\Pi})$  and a-Weyl's theorem according to topological uniform descent.

**Theorem 2.9.** *Let  $T \in B(H)$ . Then the following statements are equivalent:*

- (1)  $T$  satisfies both property  $(UW_{\Pi})$  and a-Weyl's theorem, and  $T$  is a-polaroid operator;
- (2)  $\sigma_b(T) = [\sigma_{\tau}(T) \cap \text{acc}\sigma_a(T)] \cup \text{acc}\sigma_{ea}(T) \cup [\rho_a(T) \cap \sigma(T)]$ .

*Proof.* (1)  $\Rightarrow$  (2). The inclusion " $\supseteq$ " is obvious. For the opposite inclusion, take arbitrarily  $\lambda_0$  that does not belong to the right side of (2). Without loss of generality, suppose that  $\lambda_0 \in \sigma(T)$ . Then we have  $\lambda_0 \in \sigma_a(T)$ .

**Case 1** Suppose  $\lambda_0 \notin \sigma_{\tau}(T) \cup \text{acc}\sigma_{ea}(T)$ . Then there exists  $\epsilon > 0$  such that  $\lambda \in \rho_a(T) \cup [\sigma_a(T) \setminus \sigma_{ea}(T)]$  when  $0 < \|\lambda - \lambda_0\| < \epsilon$ . From  $T$  is a-polaroid operator and  $T \in (UW_{\Pi})$ , we can get that  $\lambda_0 \in \Pi(T)$ . Thus  $\lambda_0 \notin \sigma_b(T)$ .

**Case 2** Suppose  $\lambda_0 \notin \text{acc}\sigma_a(T) \cup \text{acc}\sigma_{ea}(T)$ . Then  $\lambda_0 \in \text{iso}\sigma_a(T)$ . From  $T$  is a-polaroid operator and  $T \in (UW_{\Pi})$ , we get that  $\lambda_0 \notin \sigma_b(T)$ .

(2)  $\Rightarrow$  (1). It is clear that  $\{[\sigma_a(T) \setminus \sigma_{ea}(T)] \cup \Pi(T) \cup \pi_{00}^a(T)\} \cap [\sigma_{\tau}(T) \cap \text{acc}\sigma_a(T)] = \emptyset, \{[\sigma_a(T) \setminus \sigma_{ea}(T)] \cup \Pi(T) \cup \pi_{00}^a(T)\} \cap \text{acc}\sigma_{ea}(T) = \emptyset, \{[\sigma_a(T) \setminus \sigma_{ea}(T)] \cup \Pi(T) \cup \pi_{00}^a(T)\} \cap [\rho_a(T) \cap \sigma(T)] = \emptyset$ . Thus,  $[\sigma_a(T) \setminus \sigma_{ea}(T)] \cup \Pi(T) \cup \pi_{00}^a(T) = \sigma_0(T)$ . And  $\text{iso}\sigma_a(T) \cap \{[\sigma_{\tau}(T) \cap \text{acc}\sigma_a(T)] \cup \text{acc}\sigma_{ea}(T) \cup [\rho_a(T) \cap \sigma(T)]\} = \emptyset$ . It follows that both property  $(UW_{\Pi})$  and a-Weyl's theorem hold for  $T$ , and  $T$  is a-polaroid operator.  $\square$

From Theorem 2.8 and Theorem 2.9 we finally get the following result.

**Corollary 2.10.** Let  $T \in B(H)$  and  $Q$  a quasi-nilpotent operator with  $QT = TQ$ . Then the following statements are equivalent:

- (1)  $T + Q$  satisfies both property  $(UW_{\Pi})$  and  $a$ -Weyl's theorem, and  $T + Q$  is  $a$ -polaroid operator;
- (2)  $\sigma_b(T) = [\sigma_{\tau}(T) \cap acc\sigma_a(T)] \cup acc\sigma_{ea}(T) \cup [\rho_a(T) \cap \sigma(T)]$ .

### 3. Property $(UW_{\Pi})$ and $a$ -Weyl's theorem of operator functions

For  $T \in B(H)$ , we use  $Hol(\sigma(T))$  to denote the class of all complex-valued functions analytic on a neighborhood of  $\sigma(T)$  and not constant on any components of  $\sigma(T)$ .

**Remark 3.1.** (i)  $T$  satisfies both property  $(UW_{\Pi})$  and  $a$ -Weyl's theorem does not imply  $f(T)$  satisfies both property  $(UW_{\Pi})$  and  $a$ -Weyl's theorem, where  $f \in Hol(\sigma(T))$ .

Let  $A, B \in B(\ell^2)$  be defined by

$$A(x_1, x_2, \dots) = (0, x_1, x_2, \dots), B(x_1, x_2, \dots) = (x_2, x_3, x_4, \dots).$$

Put  $T = \begin{pmatrix} A+I & 0 \\ 0 & B-I \end{pmatrix}$ . Then  $T$  satisfies both property  $(UW_{\Pi})$  and  $a$ -Weyl's theorem. Let  $f(z) = (z-1)(z+1)$ ,  $z \in \mathbb{C}$ , we can get  $0 \in \sigma_a(f(T)) \setminus \sigma_{ea}(f(T))$ . But  $0 \notin \Pi(T)$ ,  $0 \notin \pi_{00}^a(T)$ . We know that both property  $(UW_{\Pi})$  and  $a$ -Weyl's theorem don't hold for  $f(T)$ .

(ii)  $f(T)$  satisfies both property  $(UW_{\Pi})$  and  $a$ -Weyl's theorem for some  $f \in Hol(\sigma(T))$  does not imply  $T$  satisfies both property  $(UW_{\Pi})$  and  $a$ -Weyl's theorem. Let  $A, B, C \in B(\ell^2)$  be defined by

$$A(x_1, x_2, \dots) = (0, x_2, 0, x_4, \dots), B(x_1, x_2, \dots) = (0, 0, \frac{x_2}{2}, \frac{x_3}{3}, \dots), C(x_1, x_2, \dots) = (0, x_1, x_2, \dots).$$

Put  $T = \begin{pmatrix} A+I & 0 & 0 \\ 0 & B-I & 0 \\ 0 & 0 & C+I \end{pmatrix}$ . We know that  $\sigma_a(T^2) = \sigma_{ea}(T^2) = \{re^{i\theta} : r = 2(1 + \cos \theta)\} \cup \{1, \frac{1}{3}\}$ ,  $\Pi(T^2) = \emptyset$ ,

$\pi_{00}^a(T^2) = \emptyset$ . So  $T^2 \in (UW_{\Pi})$  and satisfies  $a$ -Weyl's theorem. But  $\Pi(T) = \{\frac{1}{3}\}$ ,  $\pi_{00}^a(T) = \{-1\}$ ,  $\sigma_a(T) = \sigma_{ea}(T) = \{-1, -\frac{1}{3}\} \cup \{\lambda \in \mathbb{C} : \|\lambda - 1\| = 1\}$ . Thus both property  $(UW_{\Pi})$  and  $a$ -Weyl's theorem don't hold for  $T$ .

From the above Remark,  $T$  and  $f(T)$  satisfy both property  $(UW_{\Pi})$  and  $a$ -Weyl's theorem are not directly connected. In the following, we will discuss the property  $(UW_{\Pi})$  and  $a$ -Weyl's theorem for operator functions through the relation between  $\sigma_b(T)$  and  $\sigma_{\tau}(T)$ .

First we have this fact: For any  $f \in Hol(\sigma(T))$ ,  $f(\sigma_{ea}(T)) = \sigma_{ea}(f(T))$  if and only if for any  $\lambda, \mu \in \rho_{SF_+}(T)$ ,  $\text{ind}(T - \lambda I) \cdot \text{ind}(T - \mu I) \geq 0$ . Next, we will use topological uniform descent to describe the properties of Fredholm index.

**Lemma 3.2.** Let  $T \in B(H)$  and  $f \in Hol(\sigma(T))$ . If  $f(T) \in (UW_{\Pi})$ , then for any  $\lambda, \mu \in \rho_{SF_+}(T)$ ,  $\text{ind}(T - \lambda I) \cdot \text{ind}(T - \mu I) \geq 0$ .

*Proof.* If not, then there exist  $\lambda_0, \mu_0 \in \rho_{SF_+}(T)$  such that  $\text{ind}(T - \lambda_0 I) = m > 0$ ,  $\text{ind}(T - \mu_0 I) = -n < 0$ . Suppose that  $f(z) = (z - \lambda_0)^n (z - \mu_0)^m$  when  $n < \infty$  and  $f(z) = (z - \lambda_0)(z - \mu_0)$  when  $n = \infty$ . In both instances, we can get  $0 \in \sigma_a(f(T)) \setminus \sigma_{ea}(f(T))$ . From  $f(T) \in (UW_{\Pi})$ , we know that  $f(T)$  is Browder operator. Thus  $\lambda_0 \notin \sigma_b(T)$ , a contradiction.  $\square$

**Lemma 3.3.** Let  $T \in B(H)$  and  $T \in (UW_{\Pi})$ . Then the following statements hold:

- (1)  $\rho_{\tau}(T) \subseteq \rho_b(T) \cup acc\sigma_{ea}(T)$  if and only if for any  $\lambda \in \rho_{SF_+}(T)$ ,  $\text{ind}(T - \lambda I) \geq 0$ ;
- (2)  $\rho_{\tau}(T) \subseteq \rho_b(T) \cup acc\sigma_{SF_+}(T) \cup acc\{\lambda \in \mathbb{C} : n(T - \lambda I) = 0\}$  if and only if for any  $\lambda \in \rho_{SF_+}(T)$ ,  $\text{ind}(T - \lambda I) \leq 0$ .

*Proof.* (1). " $\Rightarrow$ ". If not, there exist  $\lambda_0 \in \rho_{SF_+}(T)$  such that  $\text{ind}(T - \lambda_0 I) < 0$ . We can get that  $\lambda_0 \in \rho_{\tau}(T)$  and  $\lambda_0 \notin acc\sigma_{ea}(T)$ , then  $\lambda_0 \in \rho_b(T)$ , a contradiction.

“ $\Leftarrow$ ”. Suppose that for any  $\lambda \in \rho_{SF_+}(T)$ ,  $\text{ind}(T - \lambda I) \geq 0$ . Take  $\lambda_0 \in \rho_\tau(T)$  but  $\lambda_0 \notin \text{acc}\sigma_{ea}(T)$ , then there exists  $\epsilon > 0$  such that  $T - \lambda I$  is an upper semi-Weyl operator when  $0 < \|\lambda - \lambda_0\| < \epsilon$ . By  $\text{ind}(T - \lambda I) \geq 0$  and  $T \in (UW_\Pi)$  we get that  $T - \lambda I$  is a Browder operator. It follows that  $\lambda_0 \in \partial\sigma(T) \cap \rho_\tau(T)$  and  $\lambda_0 \in \Pi(T)$ . Therefore,  $\lambda_0 \in \rho_b(T)$ .

(2). Similar to the proof of (1), this result is obvious.  $\square$

**Theorem 3.4.** Let  $T \in B(H)$ . Then for any  $f \in \text{Hol}(\sigma(T))$ ,  $f(T)$  satisfies both property  $(UW_\Pi)$  and a-Weyl’s theorem if and only if:

- (1)  $T$  satisfies both property  $(UW_\Pi)$  and a-Weyl’s theorem;
- (2)  $\rho_\tau(T) \subseteq \rho_b(T) \cup \text{acc}\sigma_{ea}(T)$  or  $\rho_\tau(T) \subseteq \rho_b(T) \cup \text{acc}\sigma_{SF_+}(T) \cup \text{acc}\{\lambda \in \mathbb{C} : n(T - \lambda I) = 0\}$ ;
- (3) If  $\sigma_0(T) \neq \emptyset$ , then  $\sigma_b(T) = [\sigma_\tau(T) \cap \{\lambda \in \mathbb{C} : n(T - \lambda I) = \infty\}] \cup [\text{acc}\sigma_a(T) \cap \sigma_{ea}(T)]$ .

*Proof.* “ $\Rightarrow$ ”. From Lemma 3.2 and Lemma 3.3, we only need to prove (3) holds. The inclusion “ $\supseteq$ ” is clear. For the converse, we first claim that  $\{\lambda \in \text{iso}\sigma_a(T) : n(T - \lambda I) < \infty\} = \sigma_0(T)$ . In fact, take  $\lambda_1 \in \sigma_0(T)$ ,  $\lambda_2 \in \{\lambda \in \text{iso}\sigma_a(T) : n(T - \lambda I) < \infty\}$ . Set  $\sigma_1 = \{\lambda_1\}$ ,  $\sigma_2 = \{\lambda_2\}$  and  $\sigma_3 = \sigma(T) \setminus [\sigma_1 \cup \sigma_2]$ . Then by [12, Theorem

2.10]  $T$  can be represented as  $T = \begin{pmatrix} T_1 & 0 & 0 \\ 0 & T_2 & 0 \\ 0 & 0 & T_3 \end{pmatrix}$ , where  $\sigma(T_i) = \sigma_i$ ,  $i = 1, 2, 3$ . Put  $f_0(z) = (z - \lambda_1)(z - \lambda_2)$ .

Then  $f_0(T) = \begin{pmatrix} f_0(T_1) & 0 & 0 \\ 0 & f_0(T_2) & 0 \\ 0 & 0 & f_0(T_3) \end{pmatrix}$ . Therefore  $0 \in \text{iso}\sigma_a(f_0(T))$  and  $0 < n(f_0(T)) < \infty$ . It follows that

$0 \in \pi_{00}^a(f_0(T))$ . From  $f_0(T)$  satisfies both property  $(UW_\Pi)$  and a-Weyl’s theorem, we obtain that  $f_0(T)$  is a Browder operator, and so is  $T - \lambda_2 I$ . The inclusion “ $\supseteq$ ” is clear. So  $\{\lambda \in \text{iso}\sigma_a(T) : n(T - \lambda I) < \infty\} = \sigma_0(T)$ .

Then we prove  $\sigma(T) = \sigma_a(T)$ . If not, put  $\lambda_1 \in \sigma(T) \setminus \sigma_a(T)$ . Let  $\lambda_2 \in \sigma_0(T)$  and  $f_1(T) = (T - \lambda_1 I)(T - \lambda_2 I)$ , then  $0 \in \sigma_a(f_1(T)) \setminus \sigma_{ea}(f_1(T))$ . Since  $f_1(T) \in (UW_\Pi)$ , we can get that  $f_1(T)$  is a Browder operator. It implies that  $\lambda_1 \in \rho(T)$ , a contradiction.

Take arbitrarily  $\lambda_0$  that does not belong to the right side of (3). Without loss of generality, suppose that  $\lambda_0 \in \sigma(T)$ .

**Case 1** Suppose that  $\lambda_0 \notin \sigma_\tau(T) \cup \text{acc}\sigma_a(T)$ . From  $\sigma(T) = \sigma_a(T)$  we can get that  $\lambda_0 \in \rho_\tau(T) \cap \text{iso}\sigma(T)$ , then  $\lambda_0 \in \Pi(T)$ . It follows that  $\lambda_0 \notin \sigma_b(T)$ .

**Case 2** Suppose that  $\lambda_0 \notin \{\lambda \in \mathbb{C} : n(T - \lambda I) = \infty\} \cup \text{acc}\sigma_a(T)$ . It follows that  $\lambda_0 \in \{\lambda \in \text{iso}\sigma_a(T) : n(T - \lambda I) < \infty\}$ . Thus  $\lambda_0 \in \sigma_0(T)$ .

**Case 3** Suppose that  $\lambda_0 \notin \sigma_{ea}(T)$ . Then  $\lambda_0 \in \rho_a(T) \cup [\sigma_a(T) \setminus \sigma_{ea}(T)]$ . Since  $\sigma(T) = \sigma_a(T)$  and  $T \in (UW_\Pi)$ , we can get  $\lambda_0 \notin \sigma_b(T)$ .

“ $\Leftarrow$ ”. **Case 1** Suppose that  $\sigma_0(T) = \emptyset$ . Since  $T$  satisfies both property  $(UW_\Pi)$  and a-Weyl’s theorem, we know that  $\sigma_a(T) = \sigma_{ea}(T)$ ,  $\Pi(T) = \pi_{00}^a(T) = \emptyset$ . From the condition (2) and Lemma 3.3 we can get that  $\sigma_a(f(T)) = f(\sigma_a(T)) = f(\sigma_{ea}(T)) = \sigma_{ea}(f(T))$ . Thus  $\sigma_a(f(T)) \setminus \sigma_{ea}(f(T)) = \emptyset$ . Meanwhile,  $\Pi(f(T)) \subseteq f(\Pi(T)) = \emptyset$ ,  $\pi_{00}^a(f(T)) \subseteq f(\pi_{00}^a(T)) = \emptyset$ . So  $f(T)$  satisfies both property  $(UW_\Pi)$  and a-Weyl’s theorem.

**Case 2** Suppose that  $\sigma_0(T) \neq \emptyset$ . The fact  $\sigma_b(T) = [\sigma_\tau(T) \cap \{\lambda \in \mathbb{C} : n(T - \lambda I) = \infty\}] \cup [\text{acc}\sigma_a(T) \cap \sigma_{ea}(T)]$  implies that  $\sigma_{ea}(T) = \sigma_b(T)$ . Take  $\mu_0 \in \sigma_a(f(T)) \setminus \sigma_{ea}(f(T))$  and suppose that

$$f(T) - \mu_0 I = a(T - \lambda_1 I)^{n_1} (T - \lambda_2 I)^{n_2} \cdots (T - \lambda_t I)^{n_t} g(T),$$

where  $\lambda_i \neq \lambda_j$  if  $i \neq j$  and  $g(T)$  is invertible. From the condition (2) and Lemma 3.3 we have  $\lambda_i \in \rho_a(T) \cup [\sigma_a(T) \setminus \sigma_{ea}(T)]$  for  $1 \leq i \leq t$ . Since  $\sigma_{ea}(T) = \sigma_b(T)$  and  $T \in (UW_\Pi)$ , we know that  $\lambda_i \in \rho_b(T)$  for  $1 \leq i \leq t$ . It follows that  $\mu_0 \in \sigma_0(f(T))$ . Take arbitrarily  $\mu_0 \in \Pi(f(T))$  and suppose that  $f(T) - \mu_0 I$  has the same decomposition as above. Then  $T - \lambda_i I$  is Drazin invertible for  $1 \leq i \leq t$ . Since  $T \in (UW_\Pi)$ , we can get that  $\mu_0 \in \sigma_0(f(T))$ . Take arbitrarily  $\mu_0 \in \pi_{00}^a(f(T))$  and suppose that  $f(T) - \mu_0 I$  has the same decomposition as above. Then  $\lambda_i \in \rho_a(T) \cup \text{iso}\sigma_a(T)$  and  $n(T - \lambda_i I) < \infty$ . From the condition (3) we have  $\mu_0 \in \sigma_0(f(T))$ . Hence for any  $f \in \text{Hol}(\sigma(T))$ ,  $f(T)$  satisfies both property  $(UW_\Pi)$  and a-Weyl’s theorem.  $\square$

From (3) in Theorem 3.4 we can get that  $T$  satisfies both property  $(UW_\Pi)$  and a-Weyl’s theorem, and for any  $\lambda \in \rho_{SF_+}(T)$ ,  $\text{ind}(T - \lambda I) \geq 0$ . Hence we have the following fact:



**Corollary 3.5.** Let  $T \in B(H)$  and  $\sigma_0(T) \neq \emptyset$ . Then for any  $f \in \text{Hol}(\sigma(T))$ ,  $f(T)$  satisfies both property  $(UW_{\Pi})$  and  $a$ -Weyl's theorem if and only if  $\sigma_b(T) = [\sigma_{\tau}(T) \cap \{\lambda \in \mathbb{C} : n(T - \lambda I) = \infty\}] \cup [\text{acc}\sigma_a(T) \cap \sigma_{ea}(T)]$ .

**Corollary 3.6.** Let  $T \in B(H)$ . Then  $\sigma_0(T) = \emptyset$  and for any  $f \in \text{Hol}(\sigma(T))$ ,  $f(T)$  satisfies both property  $(UW_{\Pi})$  and  $a$ -Weyl's theorem if and only if one of the following conditions holds:

- (1)  $\sigma(T) = [\sigma_{\tau}(T) \cap \{\lambda \in \mathbb{C} : n(T - \lambda I) = \infty\}] \cup \{\lambda \in \sigma_a(T) : n(T - \lambda I) = 0\} \cup [\text{acc}\sigma_a(T) \cap \sigma_{ea}(T)]$ ;
- (2)  $\sigma(T) = [\sigma_{\tau}(T) \cap \{\lambda \in \mathbb{C} : n(T - \lambda I) = d(T - \lambda I)\}] \cup \{\lambda \in \sigma(T) : n(T - \lambda I) = 0\} \cup [\text{acc}\sigma_a(T) \cap \sigma_{SF_+}(T)] \cup \{\lambda \in \mathbb{C} : n(T - \lambda I) = \infty\} \cap \text{acc}[\rho_a(T) \cap \sigma(T)]$ .

*Proof.* “ $\Rightarrow$ ”. The inclusion “ $\supseteq$ ” is clear. For the converse, by Lemma 3.2 we know that for any  $\lambda, \mu \in \rho_{SF_+}(T)$ ,  $\text{ind}(T - \lambda I) \cdot \text{ind}(T - \mu I) \geq 0$ .

**Case 1** Suppose that  $\lambda \in \rho_{SF_+}(T)$ ,  $\text{ind}(T - \lambda I) \geq 0$ . Take arbitrarily  $\lambda_0$  that does not belong to the right side of (1). We claim that  $\lambda_0 \notin \sigma_a(T)$ . In fact, if  $\lambda_0 \in \sigma_a(T)$ , then  $n(T - \lambda_0 I) > 0$ . If  $\lambda_0 \notin \sigma_{\tau}(T) \cup \text{acc}\sigma_a(T)$ , from the proof of Theorem 3.4 we can get  $\sigma(T) = \sigma_a(T)$ . Then  $\lambda_0 \in \text{iso}\sigma(T)$  and  $\lambda_0 \in \Pi(T)$ . It follows from  $T \in (UW_{\Pi})$  that  $\lambda_0 \in \sigma_0(T)$ . If  $\lambda_0 \notin \{\lambda \in \mathbb{C} : n(T - \lambda I) = \infty\} \cup \text{acc}\sigma_a(T)$ , then  $\lambda_0 \in \pi_{00}^a(T)$  and hence  $\lambda_0 \in \sigma_0(T)$ . If  $\lambda_0 \notin \sigma_{ea}(T)$ , then  $\lambda_0 \in \sigma_a(T) \setminus \sigma_{ea}(T)$ . It follows that  $\lambda_0 \in \sigma_0(T)$ . But  $\sigma_0(T) = \emptyset$ . This contradiction shows that  $\lambda_0 \notin \sigma_a(T)$ . From  $\text{ind}(T - \lambda_0 I) \geq 0$  we get that  $\lambda_0 \notin \sigma(T)$ .

**Case 2** Suppose that  $\lambda \in \rho_{SF_+}(T)$ ,  $\text{ind}(T - \lambda I) \leq 0$ . Take arbitrarily  $\lambda_0$  that does not belong to the right side of (2). We claim that  $\lambda_0 \notin \sigma(T)$ . Similar to the proof of case 1, this claim is clear.

“ $\Leftarrow$ ”. **Case 1** If condition (1) holds, we obtain that for any  $\lambda \in \rho_{SF_+}(T)$ ,  $\text{ind}(T - \lambda I) \geq 0$ . If not, there exist  $\lambda_0 \in \rho_{SF_+}(T)$ ,  $\text{ind}(T - \lambda_0 I) < 0$ . It follows that  $\lambda_0 \notin [\sigma_{\tau}(T) \cap \{\lambda \in \mathbb{C} : n(T - \lambda I) = \infty\}] \cup \{\lambda \in \sigma_a(T) : n(T - \lambda I) = 0\} \cup [\text{acc}\sigma_a(T) \cap \sigma_{ea}(T)]$ , then  $\lambda_0 \notin \sigma(T)$ , a contradiction. If there exist  $\mu \in \sigma_0(T)$ , then  $\mu \notin [\sigma_{\tau}(T) \cap \{\lambda \in \mathbb{C} : n(T - \lambda I) = \infty\}] \cup \{\lambda \in \sigma_a(T) : n(T - \lambda I) = 0\} \cup [\text{acc}\sigma_a(T) \cap \sigma_{ea}(T)]$ . By condition (1) we can get  $\mu \notin \sigma(T)$ , a contradiction. Hence  $\sigma_0(T) = \emptyset$ . It is clear that  $\{[\sigma_a(T) \setminus \sigma_{ea}(T)] \cup \Pi(T) \cup \pi_{00}^a(T)\} \cap [\sigma_{\tau}(T) \cap \{\lambda \in \mathbb{C} : n(T - \lambda I) = \infty\}] = \emptyset$ ,  $\{[\sigma_a(T) \setminus \sigma_{ea}(T)] \cup \Pi(T) \cup \pi_{00}^a(T)\} \cap \{\lambda \in \sigma_a(T) : n(T - \lambda I) = 0\} = \emptyset$ ,  $\{[\sigma_a(T) \setminus \sigma_{ea}(T)] \cup \Pi(T) \cup \pi_{00}^a(T)\} \cap [\text{acc}\sigma_a(T) \cap \sigma_{ea}(T)] = \emptyset$ . Thus  $\{[\sigma_a(T) \setminus \sigma_{ea}(T)] \cup \Pi(T) \cup \pi_{00}^a(T)\} \subseteq \rho(T)$ , a contradiction. So we can get  $\sigma_a(T) \setminus \sigma_{ea}(T) = \Pi(T) = \pi_{00}^a(T) = \emptyset$ , then  $T$  satisfies both property  $(UW_{\Pi})$  and  $a$ -Weyl's theorem. From Theorem 3.4 we know that for any  $f \in \text{Hol}(\sigma(T))$ ,  $f(T)$  satisfies both property  $(UW_{\Pi})$  and  $a$ -Weyl's theorem.

**Case 2** If condition (2) holds, we get that for any  $\lambda \in \rho_{SF_+}(T)$ ,  $\text{ind}(T - \lambda I) \leq 0$ . Similar to the proof of case 1, the result is trivial.  $\square$

From Corollary 3.5 and Corollary 3.6 we can describe the property  $(UW_{\Pi})$  and  $a$ -Weyl's theorem for operator functions through the relation between  $\sigma_b(T)$  and  $\sigma_{\tau}(T)$ .

**Theorem 3.7.** Let  $T \in B(H)$ . Then for any  $f \in \text{Hol}(\sigma(T))$ ,  $f(T)$  satisfies both property  $(UW_{\Pi})$  and  $a$ -Weyl's theorem if and only if one of the following statements holds:

- (1)  $\sigma(T) = [\sigma_{\tau}(T) \cap \{\lambda \in \mathbb{C} : n(T - \lambda I) = \infty\}] \cup \{\lambda \in \sigma_a(T) : n(T - \lambda I) = 0\} \cup [\text{acc}\sigma_a(T) \cap \sigma_{ea}(T)]$ ;
- (2)  $\sigma(T) = [\sigma_{\tau}(T) \cap \{\lambda \in \mathbb{C} : n(T - \lambda I) = d(T - \lambda I)\}] \cup \{\lambda \in \sigma(T) : n(T - \lambda I) = 0\} \cup [\text{acc}\sigma_a(T) \cap \sigma_{SF_+}(T)] \cup \{\lambda \in \mathbb{C} : n(T - \lambda I) = \infty\} \cap \text{acc}[\rho_a(T) \cap \sigma(T)]$ ;
- (3)  $\sigma_b(T) = [\sigma_{\tau}(T) \cap \{\lambda \in \mathbb{C} : n(T - \lambda I) = \infty\}] \cup [\text{acc}\sigma_a(T) \cap \sigma_{ea}(T)]$ .

We can get that property  $(UW_{\Pi})$  is transmitted from Drazin invertible operator to its Drazin inverse in [2]. Now, we will discuss  $S$  satisfies both property  $(UW_{\Pi})$  and  $a$ -Weyl's theorem by topological uniform descent.

If  $T \in B(H)$  is Drazin invertible with inverse  $S$ , then  $\text{asc}(T) = \text{des}(T) = p$  for any  $p \in \mathbb{N}$ . We know that  $R(T^p)$  is closed and  $H = N(T^p) \oplus R(T^p)$ . Under this space decomposition,  $T = T_1 \oplus T_2$ , where  $T_1$  is nilpotent operator and  $T_2$  is invertible. Thus  $S = 0 \oplus T_2^{-1}$ . In [1, 4, 5], we get that

$$\sigma(S) \setminus \{0\} = \{\frac{1}{\lambda} : \lambda \in \sigma(T) \setminus \{0\}\}, \sigma_*(S) \setminus \{0\} = \{\frac{1}{\lambda} : \lambda \in \sigma_*(T) \setminus \{0\}\}, * \in \{b, ea, a, \tau, D\}.$$

Besides, one can verify that for any  $\lambda \neq 0$ ,  $n(S - \lambda I) = n(T - \frac{1}{\lambda} I)$ ,  $d(S - \lambda I) = d(T - \frac{1}{\lambda} I)$  and  $\text{acc}\sigma_a(S) \setminus \{0\} = \{\frac{1}{\lambda} : \lambda \in \text{acc}\sigma_a(T) \setminus \{0\}\}$ .

**Theorem 3.8.** Let  $T \in B(H)$  be Drazin invertible with inverse  $S$ . Then

(1)  $T$  satisfies both property  $(UW_{\Pi})$  and  $a$ -Weyl's theorem if and only if  $S$  satisfies both property  $(UW_{\Pi})$  and  $a$ -Weyl's theorem;

(2) For any  $f \in \text{Hol}(\sigma(T)) \cap \text{Hol}(\sigma(S))$ ,  $f(T)$  satisfies both property  $(UW_{\Pi})$  and  $a$ -Weyl's theorem if and only if  $f(S)$  satisfies both property  $(UW_{\Pi})$  and  $a$ -Weyl's theorem.

*Proof.* (1). “ $\Rightarrow$ ”. From Theorem 2.1, we only need  $\sigma_b(S) = [\sigma_{\tau}(S) \cap \{\lambda \in \mathbb{C} : n(S - \lambda I) = d(S - \lambda I)\}] \cup \{\lambda \in \sigma(S) : n(S - \lambda I) = 0\} \cup [\text{acc}\sigma_a(S) \cap \sigma_{ea}(S)] \cup \{\{\lambda \in \mathbb{C} : n(S - \lambda I) = \infty\} \cap \text{acc}[\rho_a(S) \cap \sigma(S)]\}$ . If  $T$  is invertible, then  $S = T^{-1}$ . The conclusion is clear. In the following, we assume  $T$  is not invertible but Drazin invertible.

Let  $\lambda$  does not belong to the right side. If  $\lambda = 0$ . Since  $0 \in \Pi(T)$  and  $T \in (UW_{\Pi})$ , we get that  $0 \notin \sigma_b(T)$ . Thus  $0 \notin \sigma_b(S)$  ([2, Lemma 4.9]). If  $\lambda \neq 0$ . Now,  $S - \lambda I = \begin{pmatrix} -\lambda I & 0 \\ 0 & \lambda T_2^{-1}(\frac{1}{\lambda}I - T_2) \end{pmatrix}$ , then  $\frac{1}{\lambda} \notin [\sigma_{\tau}(T_2) \cap \{\lambda \in \mathbb{C} : n(T_2 - \lambda I) = d(T_2 - \lambda I)\}] \cup \{\lambda \in \sigma(T_2) : n(T_2 - \lambda I) = 0\} \cup [\text{acc}\sigma_a(T_2) \cap \sigma_{ea}(T_2)] \cup \{\{\lambda \in \mathbb{C} : n(T_2 - \lambda I) = \infty\} \cap \text{acc}[\rho_a(T_2) \cap \sigma(T_2)]\}$ . Under above space decomposition, we know that  $T - \frac{1}{\lambda}I = \begin{pmatrix} T_1 - \frac{1}{\lambda}I & 0 \\ 0 & T_2 - \frac{1}{\lambda}I \end{pmatrix}$ , where  $T_1 - \lambda I$  is invertible. So  $\frac{1}{\lambda} \notin [\sigma_{\tau}(T) \cap \{\lambda \in \mathbb{C} : n(T - \lambda I) = d(T - \lambda I)\}] \cup \{\lambda \in \sigma(T) : n(T - \lambda I) = 0\} \cup [\text{acc}\sigma_a(T) \cap \sigma_{ea}(T)] \cup \{\{\lambda \in \mathbb{C} : n(T - \lambda I) = \infty\} \cap \text{acc}[\rho_a(T) \cap \sigma(T)]\}$ . By Theorem 2.1 we can get  $\frac{1}{\lambda} \notin \sigma_b(T)$  and so  $\frac{1}{\lambda} \notin \sigma_b(S)$ .

“ $\Leftarrow$ ”. Suppose that  $S$  satisfies both property  $(UW_{\Pi})$  and  $a$ -Weyl's theorem. The Drazin inverse of  $S$  is  $U := T^2S = TST$  and Drazin inverse of  $U$  is  $T$  ([1, Chapter 1]). Thus,  $T$  satisfies both property  $(UW_{\Pi})$  and  $a$ -Weyl's theorem.

(2). “ $\Rightarrow$ ”. It is obvious that  $T$  satisfies both property  $(UW_{\Pi})$  and  $a$ -Weyl's theorem. Then from (1), we get that  $S$  satisfies both property  $(UW_{\Pi})$  and  $a$ -Weyl's theorem. Suppose that  $\sigma_0(S) \neq \emptyset$ , we claim that  $\sigma_0(T) \neq \emptyset$ . In fact, let  $\lambda \in \sigma_0(S)$ . If  $\lambda = 0$ , then  $0 < \dim N(T^p) < \infty$ . It follows that  $n(T_1) > 0$  and  $T$  is not invertible. By  $T \in (UW_{\Pi})$  we have that  $T_1$  is Browder operator. So  $T$  is Browder operator and  $0 \in \sigma_0(T)$ . If  $\lambda \neq 0$ . Now,  $S - \lambda I = \begin{pmatrix} -\lambda I & 0 \\ 0 & \lambda T_2^{-1}(\frac{1}{\lambda}I - T_2) \end{pmatrix}$ . It follows that  $\frac{1}{\lambda}I - T_2$  is Browder operator but not invertible. We know that  $T - \frac{1}{\lambda}I = \begin{pmatrix} T_1 - \frac{1}{\lambda}I & 0 \\ 0 & T_2 - \frac{1}{\lambda}I \end{pmatrix}$ , then  $\frac{1}{\lambda} \in \sigma_0(T)$ . From Corollary 3.5 we get  $\sigma_b(T) = [\sigma_{\tau}(T) \cap \{\lambda \in \mathbb{C} : n(T - \lambda I) = \infty\}] \cup [\text{acc}\sigma_a(T) \cap \sigma_{ea}(T)]$ . By using the similar way of (1) we get that  $\sigma_b(S) = [\sigma_{\tau}(S) \cap \{\lambda \in \mathbb{C} : n(S - \lambda I) = \infty\}] \cup [\text{acc}\sigma_a(S) \cap \sigma_{ea}(S)]$ . Moreover,  $\sigma_{ea}(f(S)) = f(\sigma_{ea}(S))$  for any  $f \in \text{Hol}(\sigma(T)) \cap \text{Hol}(\sigma(S))$ . From Lemma 3.3 and Theorem 3.4  $f(S)$  satisfies both property  $(UW_{\Pi})$  and  $a$ -Weyl's theorem.

“ $\Leftarrow$ ”. The same as the above proof.  $\square$

## References

- [1] P. Aiena, *Fredholm and local spectral theory II, with application to Weyl-type theorems*, Springer Lecture Notes of Math 2235, 2018.
- [2] P. Aiena, M. Kachad, *Property  $(UW_{\Pi})$  under perturbations*, Ann. Func. Anal. **11** (2020), 29–46.
- [3] P. Aiena, *Semi-Fredholm operators, perturbation theory and localized SVEP*, Mérida, Venezuela, 2007.
- [4] P. Aiena, S. Triolo, *Fredholm spectra and Weyl type theorems for Drazin invertible operators*, Mediterr. J. Math. **13** (2016), 4385–4400.
- [5] P. Aiena, S. Triolo, *Local spectral theory for Drazin invertible operators*, J. Math. Anal. Appl. **435** (2016), 414–424.
- [6] M. Berkani, M. Kachad, *New Browder and Weyl type theorems*, Bull. Korean. Math. Soc. **52** (2015), 439–452.
- [7] X. Cao, J. Dong, J. Liu, *Weyl's theorem and its perturbations for the functions of operators*, Oper. Matrices. **12** (2018), 1145–1157.
- [8] X. Cao, *Topological uniform descent and Weyl type theorem*, Linear. Algebra. Appl. **420** (2007), 175–182.
- [9] S. Grabiner, *Uniform ascent and descent of bounded operators*, J. Math. Soc. Japan. **34** (1982), 317–337.
- [10] C. Li, S. Zhu, Y. Feng, *Weyl's theorem for functions of operators and approximation*, Intege. Equ. Oper. Theory. **67** (2010), 481–497.
- [11] S. Qiu, X. Cao, *Property  $(UWE)$  for operators and operator matrices*, J. Math. Anal. Appl. **509** (2022), 125951.
- [12] H. Radjavi, P. Rosenthal, *Invariant Subspaces*, 2nd edn, Dover Publications, Mineola, 2003.
- [13] V. Rakočević, *On a class of operators*, Mat. Vesnik. **37** (1985), 423–425.
- [14] V. Rakočević, *Operators obeying  $a$ -Weyl's theorem*, Rev. Roum. Math. Pures. Appl. **34** (1989), 915–919.
- [15] W. Shi, *Topological uniform descent and compact perturbations*, RACSAM. **113** (2019), 2221–2233.
- [16] H. Weyl, *Über beschränkte quadratische Formen, deren Differenz vollsteig ist*, Rend. Circ. Mat. Di. Palermo. **27** (1909), 373–392.
- [17] L. Yang, X. Cao, *Single-Valued Extension Property and Property  $(w)$* , Funct. Anal. its. Appl. **55** (2021), 316–325.