



Continuity of pseudodifferential operators with nonsmooth symbols on mixed-norm Lebesgue spaces

Ivan Ivec^a

^aFaculty of Metallurgy, University of Zagreb, Aleja narodnih heroja 3, Sisak, Croatia

Abstract. Mixed-norm Lebesgue spaces found their place in the study of some questions in the theory of partial differential equations, as can be seen from recent interest in the continuity of certain classes of pseudodifferential operators on these spaces. In this paper, we use some recent advances in the pseudodifferential calculus for nonsmooth symbols to prove the boundedness of pseudodifferential operators with such symbols on mixed-norm Lebesgue spaces.

1. Introduction

Continuity of pseudodifferential operators with symbols in the nonsmooth Hörmander class $S_{\rho,\delta,N,N'}^m$ (the definition is recalled at the beginning of Section 3) on mixed-norm Lebesgue spaces is our main interest in this paper. Regarding the continuity on classical Lebesgue spaces, there is a famous result by Coifman and Meyer [7, Theorem 7], namely that for $0 \leq \delta \leq \rho \leq 1$, $\delta < 1$ and $m = 0$ it is enough to have $N, N' > \frac{d}{2}$ to obtain the continuity on $L^2(\mathbb{R}^d)$. Moreover, in Hörmander [10] it was shown that we have the following necessary condition for continuity on $L^p(\mathbb{R}^d)$ spaces:

$$m \leq -d(1 - \rho) \left| \frac{1}{2} - \frac{1}{p} \right|. \quad (1)$$

From this condition, it is clear that for the zeroth-order operators ($m = 0$) and $p \neq 2$ we can obtain the continuity result only in the case $\rho = 1$. An interesting result for nonsmooth symbols in the L^p setting is obtained in [1, Theorem 2], where the norm of an operator is estimated in terms of the norm of its symbol. That was useful for the construction of certain variants of H-distributions (introduced in [4]), both in [1] and [12]. The condition (1) proved to be relevant also for continuity on Lebesgue spaces on compact and graded Lie groups [6, 9]. We investigate this further for an arbitrary mixed-norm Lebesgue space $L^{\mathbf{p}}(\mathbb{R}^d)$, $\mathbf{p} \in \langle 1, \infty \rangle^d$. Mixed-norm spaces are suitable to describe properties of functions that have different growth with respect to different variables. They are naturally connected with problems for estimating solutions of partial differential equations modeling physical processes depending both on space and time. For instance, they were used in the famous Strichartz estimates for solutions of the Schrödinger equation

2020 *Mathematics Subject Classification.* Primary 35S05; Secondary 46E30, 47G30

Keywords. pseudodifferential operators, nonsmooth symbols, mixed-norm Lebesgue spaces

Received: 29 September 2022; Accepted: 10 April 2023

Communicated by Marko Nedeljkov

Research supported by the Croatian Science Foundation under the project 2449 MiTPDE.

Email address: iivec@simet.unizg.hr (Ivan Ivec)

[8]. The definition and some properties of mixed-norm Lebesgue spaces are given in [5] and recalled in the Appendix of [3]. They are also partially recalled under the notation below.

In [2, 3] the boundedness of pseudodifferential operators on mixed-norm Lebesgue spaces have already been obtained, but these results cover only the case of infinitely smooth symbols together with $m = 0$. Also, a detailed discussion on similar results from the literature was given there.

In the next section we recall some already known results that we need as ingredients in our proof: sufficient conditions for boundedness of a linear operator $A : L_c^\infty(\mathbb{R}^d) \rightarrow L_{loc}^1(\mathbb{R}^d)$ on mixed-norm Lebesgue spaces, obtained in [3], and pseudodifferential calculus for the adjoint of the pseudodifferential operator with a nonsmooth symbol, obtained in [11]. In the last section, we use these results, together with an extension of [13, VI.4, Proposition 1], to obtain the continuity of pseudodifferential operators with symbols in $S_{\rho,\delta,N,N'}^m$ on mixed-norm Lebesgue spaces.

Notation. We use the following notation and definition for the Fourier transform of the function $u \in L^1(\mathbb{R}^d)$:

$$\hat{u}(\xi) = \mathcal{F}u(\xi) = \int_{\mathbb{R}^d} e^{-ix \cdot \xi} u(x) dx.$$

When the area of integration is not specified, the whole space is assumed. We denote by $\mathcal{S}(\mathbb{R}^d)$ the Schwartz space of smooth rapidly decreasing functions on \mathbb{R}^d and by $\mathcal{S}'(\mathbb{R}^d)$ its dual, the space of tempered distributions. For $N, N' \in \mathbb{N}_0$ we use the following family of semi-norms on $\mathcal{S}(\mathbb{R}^d)$:

$$|\varphi|_{N,N'} = \sup_{|\alpha| \leq N, |\beta| \leq N'} \sup_{x \in \mathbb{R}^d} |x^\alpha \partial^\beta \varphi(x)|,$$

and by $\mathcal{S}_{N,N'}(\mathbb{R}^d)$ we denote the Banach space of all functions $\varphi \in C^{N'}(\mathbb{R}^d)$ for which $|\varphi|_{N,N'} < \infty$. We denote Lebesgue spaces by $L^p(\mathbb{R}^d)$, locally Lebesgue spaces by $L_{loc}^p(\mathbb{R}^d)$ and Lebesgue spaces with compact support by $L_c^p(\mathbb{R}^d)$. By C we always denote a constant, even if it changes during calculation, while C_p is a constant depending on parameter p . By $\lfloor x \rfloor$ we denote the largest integer not greater than x , while $\lfloor x \rfloor_2$ is the largest even integer not greater than x . We also use the standard notation $m^+ = \max\{m, 0\}$ and write m^+ as $(m)^+$ when m is a larger expression, while $\|\cdot\|_p$ is the standard p -norm on \mathbb{R}^d :

$$\|x\|_p = \sqrt[p]{|x_1|^p + \dots + |x_d|^p}, \quad \|x\|_\infty = \max\{|x_1|, \dots, |x_d|\},$$

and we denote $\|x\|_2$ simply by $|x|$.

Finally, we recall the definition and some basic properties of mixed-norm Lebesgue spaces. For any $\mathbf{p} \in \langle 0, \infty \rangle^d$, $L^{\mathbf{p}}(\mathbb{R}^d)$ denotes (with identification of almost everywhere equal functions) the space of all measurable complex functions f on \mathbb{R}^d for which we have

$$\|f\|_{\mathbf{p}} = \left(\int_{\mathbb{R}} \dots \left(\int_{\mathbb{R}} \left(\int_{\mathbb{R}} |f(x_1, \dots, x_d)|^{p_1} dx_1 \right)^{p_2/p_1} dx_2 \right)^{p_3/p_2} \dots dx_d \right)^{1/p_d} < \infty.$$

In other words, for $i = 1, \dots, d$ we compute (in that order) the (quasi)norms $\|\cdot\|_{L^{p_i}}$ in variable x_i . This extends analogously in the case when some $p_i = \infty$, with obvious changes.

For $\mathbf{p} \in [1, \infty]^d$ thus defined $\|\cdot\|_{\mathbf{p}}$ is a norm on $L^{\mathbf{p}}(\mathbb{R}^d)$, which becomes a Banach space. Furthermore, if we define the conjugate exponent $\mathbf{p}' = (p'_1, \dots, p'_d)$ to $\mathbf{p} = (p_1, \dots, p_d)$ as the one satisfying $\frac{1}{p_i} + \frac{1}{p'_i} = 1$, for $i \in \{1, 2, \dots, d\}$, which we shall conveniently write as $1/\mathbf{p} + 1/\mathbf{p}' = \mathbf{1}$, then $L^{\mathbf{p}'}(\mathbb{R}^d)$ is the dual of $L^{\mathbf{p}}(\mathbb{R}^d)$, for $\mathbf{p} \in [1, \infty)^d$, and one has the Hölder inequality:

Lemma 1.1. For any $\mathbf{p} \in [1, \infty]^d$, $f \in L^{\mathbf{p}}(\mathbb{R}^d)$ and $g \in L^{\mathbf{p}'}(\mathbb{R}^d)$ one has

$$\left| \int_{\mathbb{R}^d} f(x)g(x) dx \right| \leq \|f\|_{\mathbf{p}} \|g\|_{\mathbf{p}'}$$

In the proof of Theorem 3.2 we also need the generalized Minkowski inequality:

Lemma 1.2. For any $\mathbf{p} \in [1, \infty)^{d_1}$ and a function $f \in L^{(\mathbf{p}, 1, \dots, 1)}(\mathbb{R}^{d_1+d_2})$ one has

$$\left\| \int_{\mathbb{R}^{d_2}} f(\cdot, y) dy \right\|_{\mathbf{p}} \leq \int_{\mathbb{R}^{d_2}} \|f(\cdot, y)\|_{\mathbf{p}} dy.$$

An important fact in proving many results are the inclusions $\mathcal{S}(\mathbb{R}^d) \subseteq L^{\mathbf{p}}(\mathbb{R}^d) \subseteq \mathcal{S}'(\mathbb{R}^d)$, where for $\mathbf{p} \in [1, \infty)^d$, they are dense and continuous [2].

2. The general framework

Our main tool is the following theorem, proved in [3, Theorem 1]. To state the theorem the following notation is convenient: Take $l \in \{0, \dots, (d-1)\}$ and split $x = (\bar{x}, x') = (x_1, \dots, x_l; x_{l+1}, \dots, x_d)$. Next define (with a slight abuse of notation, for simplicity) for $\bar{\mathbf{p}} = (p_1, \dots, p_l)$

$$L^{\bar{\mathbf{p}}, p}(\mathbb{R}^d) = L^{(\bar{\mathbf{p}}, p, \dots, p)}(\mathbb{R}^d) \quad \text{and} \quad \|f\|_{\bar{\mathbf{p}}, p} = \|f\|_{(\bar{\mathbf{p}}, p, \dots, p)}.$$

Of course, for $l = 0$ we take $\|f(\cdot, x')\|_{\bar{\mathbf{p}}} = |f(x')|$ and $\|f\|_{\bar{\mathbf{p}}, p} = \|f\|_{L^p}$. We also define (for each $l \in \{0, \dots, (d-1)\}$, $t > 0$ and $y' \in \mathbb{R}^{d-l}$):

$$\mathcal{F}_{l,t}^{y'} := \left\{ f \in L^1_{loc}(\mathbb{R}^d) : \text{supp } f \subseteq \mathbb{R}^l \times \{x' : |x' - y'|_{\infty} \leq t\} \ \& \ \int_{\mathbb{R}^{d-l}} f(\bar{x}, x') dx' = 0 \ (\text{ae } \bar{x} \in \mathbb{R}^l) \right\}.$$

Theorem 2.1. Assume that $A, A^* : L^{\infty}_c(\mathbb{R}^d) \rightarrow L^1_{loc}(\mathbb{R}^d)$ are formally adjoint linear operators, i.e. such that

$$(\forall \varphi, \psi \in C^{\infty}_c(\mathbb{R}^d)) \quad \int_{\mathbb{R}^d} (A\varphi)\bar{\psi} = \int_{\mathbb{R}^d} \varphi \overline{A^*\psi}.$$

Furthermore, let us assume that (both for $T = A$ and $T = A^*$) there exist constants $N > 1$ and $c_1 > 0$ satisfying

$$(\forall l \in \{0, \dots, (d-1)\})(\forall x'_0 \in \mathbb{R}^{d-l})(\forall t > 0) \quad \int_{|x' - x'_0|_{\infty} > Nt} \|Tf(\cdot, x')\|_{\bar{\mathbf{p}}} dx' \leq c_1 \|f\|_{\bar{\mathbf{p}}, 1}, \tag{2}$$

for any function $f \in L^{\infty}_c(\mathbb{R}^d) \cap \mathcal{F}_{l,t}^{x'_0}$ and any $\bar{\mathbf{p}} \in \langle 1, \infty \rangle^l$.

If for some $q \in \langle 1, \infty \rangle$ operator A has a continuous extension to an operator from $L^q(\mathbb{R}^d)$ to itself with norm c_q , then A can be extended by the continuity to an operator from $L^{\mathbf{p}}(\mathbb{R}^d)$ to itself for any $\mathbf{p} \in \langle 1, \infty \rangle^d$, with the norm

$$\begin{aligned} \|A\|_{L^{\mathbf{p}} \rightarrow L^{\mathbf{p}}} &\leq \sum_{k=1}^d c^k \prod_{j=0}^{k-1} \max(p_{d-j}, (p_{d-j} - 1)^{-1/p_{d-j}})(c_1 + c_q) \\ &\leq c' \prod_{j=0}^{d-1} \max(p_{d-j}, (p_{d-j} - 1)^{-1/p_{d-j}})(c_1 + c_q), \end{aligned}$$

where c and c' are constants depending only on N and d .

We also use [11, Theorem 5.5] to conclude that for a pseudodifferential operator with symbol $\sigma \in S^m_{\rho, \delta, N, N'}$ where $\delta \leq \rho$, $m \in [-d, 0]$ and $N, N' \in 2\mathbb{N}_0$ are such that

$$N > \frac{(3 - \delta)d + (5 - \delta)(1 - \delta)}{(1 - \delta)^2}, \quad N' > 6d + 12, \tag{3}$$

its formally adjoint operator exists and has symbol $\sigma^* \in S^m_{\rho, \delta, M, M'}$, where $M, M' \in 2\mathbb{N}_0$ satisfy

$$N - M > \frac{d + ([d]_2 + 2)\delta}{1 - \delta}, \quad N - M \geq \frac{-m + (1 - \delta)d + ([d]_2 + 2)\delta}{1 - \delta}, \quad N' - M' \geq [d]_2 + 2. \tag{4}$$

The claim is valid for all $m \leq 0$ because it can easily be checked that the condition $m \geq -d$ used in [11, Theorem 5.5] could be omitted.

3. Boundedness of ΨDO

$S_{\rho,\delta,N,N'}^m$ is a nonsmooth variant of Hörmander class $S_{\rho,\delta'}^m$, $m \in \mathbb{R}$, $0 \leq \rho \leq 1$, $0 \leq \delta < 1$. It consists of all $\sigma : \mathbb{R}^d \times \mathbb{R}^d \rightarrow \mathbb{C}$ such that for all multi-indices $|\alpha| \leq N, |\beta| \leq N'$ it holds

$$(\forall x \in \mathbb{R}^d)(\forall \xi \in \mathbb{R}^d) \quad |\partial_x^\alpha \partial_\xi^\beta \sigma(x, \xi)| \leq C_{\alpha,\beta} \langle \xi \rangle^{m-\rho|\beta|+\delta|\alpha|}, \tag{5}$$

where $\langle \xi \rangle = (1 + |\xi|^2)^{\frac{1}{2}}$, $C_{\alpha,\beta}$ is a constant depending only on α and β and where all partial derivatives are understood to be continuous.

For such symbols we denote the corresponding pseudodifferential operator T_σ by

$$T_\sigma \varphi(x) = \int_{\mathbb{R}^d} e^{ix \cdot \xi} \sigma(x, \xi) \hat{\varphi}(\xi) d\xi, \quad \varphi \in \mathcal{S}(\mathbb{R}^d), \tag{6}$$

where $d\xi = (2\pi)^{-d} d\xi$. As T_σ doesn't map $\mathcal{S}(\mathbb{R}^d)$ to $\mathcal{S}(\mathbb{R}^d)$ when symbols are not infinitely smooth, we cannot extend this definition to the space of tempered distributions, but we can extend it to mixed-norm Lebesgue spaces $L^{\mathbf{p}}(\mathbb{R}^d)$, $\mathbf{p} \in \langle 1, \infty \rangle^d$ using a formula $\langle T_\sigma u, \varphi \rangle = \langle u, T_{\sigma^*} \varphi \rangle$, where $\langle \cdot, \cdot \rangle$ is the dual product, $u \in L^{\mathbf{p}}(\mathbb{R}^d)$, $\varphi \in \mathcal{S}(\mathbb{R}^d)$. The only requirement for this to work is $T_{\sigma^*} \varphi \in L^{\mathbf{p}'}(\mathbb{R}^d)$. This requirement also guarantees that the continuous extension of T_σ on $L^{\mathbf{p}}(\mathbb{R}^d)$, if it exists, should be given by the above duality formula. As T_σ actually maps $\mathcal{S}(\mathbb{R}^d)$ to $\mathcal{S}_{M',M}(\mathbb{R}^d)$ [11, Theorem 2.2], a sufficient condition for the above requirement to be fulfilled is $M' \geq d$. This can be seen easily by writing

$$\|T_{\sigma^*} \varphi\|_{L^{\mathbf{p}'}} = \left(\int \langle x_d \rangle^{-1-\epsilon} \dots \left(\int \langle x_1 \rangle^{-1-\epsilon} \left(\langle x_1 \rangle^{\frac{1+\epsilon}{p'_1}} \langle x_2 \rangle^{\frac{1+\epsilon}{p'_2}} \dots \langle x_d \rangle^{\frac{1+\epsilon}{p'_d}} |T_{\sigma^*} \varphi| \right)^{p'_1} dx_1 \right)^{\frac{p'_2}{p'_1}} \dots dx_d \right)^{\frac{1}{p'_d}}$$

and by noticing that we can take $\epsilon > 0$ so small that $\frac{1+\epsilon}{p'_i} \leq 1$ for every $i \in \{1, 2, \dots, d\}$. This condition will be satisfied in our main result – Theorem 3.2.

In this paper we prove that the continuous extension of T_σ on $L^{\mathbf{p}}(\mathbb{R}^d)$ exists for $\rho > 0$, $\delta \leq \rho$,

$$m \leq -(1 - \rho)(d + 1 + \rho), \tag{7}$$

and for sufficiently smooth symbols. As the estimate (7) is more crude than (1) we expect that even better results are possible.

From the estimate (5), it follows easily that for a fixed $x \in \mathbb{R}^d$ we have $\sigma(x, \cdot) \in \mathcal{S}'(\mathbb{R}^d)$ and so there is a $k(x, \cdot) \in \mathcal{S}'(\mathbb{R}^d)$ such that $\widehat{k(x, \cdot)} = \sigma(x, \cdot)$. We call the tempered distribution $k(x, \cdot)$ a kernel of the operator T_σ and using properties of the convolution and Fourier transform we can write (6) in the form

$$T_\sigma \varphi(x) = k(x, \cdot) * \varphi. \tag{8}$$

The kernel $k(x, \cdot)$ is a function away from the origin with the following estimates on its derivatives.

Lemma 3.1. *Let $\sigma \in S_{\rho,\delta,N,N'}^m$, $\rho > 0$. Then the kernel $k(x, z)$ satisfies*

$$|\partial_x^\alpha \partial_z^\beta k(x, z)| \leq C_{\alpha,\beta,L} \cdot |z|^{-d-m-\delta|\alpha|-|\beta|-L}, \quad z \neq 0, \tag{9}$$

for all $|\alpha| \leq N, |\beta| \geq 0$ and

$$L \geq (1 - \rho) \left(\left\lfloor \frac{d + m + \delta|\alpha| + |\beta|}{\rho} \right\rfloor + 1 \right)^+ \tag{10}$$

such that $N' \geq d + m + \delta|\alpha| + |\beta| + L > 0$ and $N' > \frac{d+m+\delta|\alpha|+|\beta|}{\rho}$, and where $C_{\alpha,\beta,L}$ is a constant depending only on α, β and L .

Proof. We take a fixed nonnegative $\eta \in C_c^\infty(\mathbb{R}^d)$ such that $\eta(\xi) = 1$ for $|\xi| \leq 1$ and $\eta(\xi) = 0$ for $|\xi| \geq 2$. We also define $\zeta(\xi) = \eta(\xi) - \eta(2\xi)$. Then we have the following decomposition (pointwise convergence in the sense of tempered distributions):

$$T_\sigma = \sum_{j=0}^\infty T_{\sigma_j},$$

where $\sigma_0(x, \xi) = \sigma(x, \xi)\eta(\xi)$ and $\sigma_j(x, \xi) = \sigma(x, \xi)\zeta(2^{-j}\xi)$ for $j \geq 1$. Details of this so-called dyadic decomposition are described in [13, 4.1–4.2]. We denote kernels of the operators T_{σ_j} ($j \geq 0$) by $k_j(x, \xi)$ and these are all smooth functions. Namely, as $\sigma_j(x, \xi)$ are functions with compact support they actually belong to $\mathcal{S}_{\infty, N'}(\mathbb{R}^d)$ and thus, by [11, Lemma 2.1], we have $k_j(x, \xi) \in \mathcal{S}_{N', \infty}(\mathbb{R}^d) \subseteq C^\infty(\mathbb{R}^d)$.

Obviously, $\sigma(x, \xi) = \sum_{j=0}^\infty \sigma_j(x, \xi)$ pointwise and so, using (5) and Lebesgue dominated convergence theorem, we easily obtain that $k(x, \cdot) = \sum_{j=0}^\infty k_j(x, \cdot)$, with the sum converging for each fixed x in the sense of tempered distributions. This, continuity of derivatives on the space of distributions and the uniqueness of the limit in the space of distributions ensures that it is enough to prove that

$$\sum_{j=0}^\infty |\partial_x^\alpha \partial_z^\beta k_j(x, z)| \tag{11}$$

satisfies the estimate given by the right-hand side of (9).

So, we first find certain estimates on $|\partial_x^\alpha \partial_z^\beta k_j(x, z)|$. As $\sigma_j(x, \xi)$ have compact supports, we have

$$k_j(x, z) = \int_{\mathbb{R}^d} \sigma_j(x, \xi) e^{i\xi \cdot z} d\xi,$$

and now for all multi-indices $|\alpha| \leq N$, $|\beta| \geq 0$ and $|\gamma| \leq N'$ it follows

$$(-iz)^\gamma \partial_x^\alpha \partial_z^\beta k_j(x, z) = \int_{\mathbb{R}^d} \partial_\xi^\gamma ((i\xi)^\beta \partial_x^\alpha \sigma_j(x, \xi)) e^{i\xi \cdot z} d\xi, \tag{12}$$

and, using the fact that σ_j is supported in $|\xi| \leq 2^{j+1}$ and for $j \neq 0$ also in $|\xi| \geq 2^{j-1}$, we can estimate the integrand above:

$$\begin{aligned} \left| \partial_\xi^\gamma ((i\xi)^\beta \partial_x^\alpha \sigma_j(x, \xi)) \right| &= \left| \sum_{\gamma' \leq \gamma} \binom{\gamma}{\gamma'} \partial_\xi^{\gamma'} ((i\xi)^\beta \partial_x^\alpha \partial_\xi^{\gamma-\gamma'} \sigma_j(x, \xi)) \right| \\ &\leq C_\gamma \sum_{\gamma' \leq \gamma} |\xi|^{(|\beta|-|\gamma'|)_+} \left| \sum_{\gamma'' \leq \gamma-\gamma'} \binom{\gamma-\gamma'}{\gamma''} \partial_x^\alpha \partial_\xi^{\gamma''} \sigma(x, \xi) \partial_\xi^{\gamma-\gamma'-\gamma''} (\zeta(2^{-j}\xi)) \right| \\ &\leq C_{\alpha, \gamma} \sum_{\gamma' \leq \gamma} \sum_{\gamma'' \leq \gamma-\gamma'} |\xi|^{(|\beta|-|\gamma'|)_+} \cdot \langle \xi \rangle^{m-\rho|\gamma''|+\delta|\alpha|} \cdot 2^{-j|\gamma-\gamma'-\gamma''|} \\ &\leq C_{\alpha, \beta, \gamma} \sum_{\gamma' \leq \gamma} \sum_{\gamma'' \leq \gamma-\gamma'} 2^{j(|\beta|-|\gamma'|)} \cdot 2^{j(m-\rho|\gamma''|+\delta|\alpha|)} \cdot 2^{-j|\gamma-\gamma'-\gamma''|} \\ &= C_{\alpha, \beta, \gamma} \sum_{\gamma' \leq \gamma} \sum_{\gamma'' \leq \gamma-\gamma'} 2^{j(m+\delta|\alpha|+|\beta|-\rho|\gamma|+(\rho-1)(|\gamma|-|\gamma''|))} \\ &\leq C_{\alpha, \beta, \gamma} \cdot 2^{j(m+\delta|\alpha|+|\beta|-\rho|\gamma|)}, \end{aligned}$$

where we have used $\rho \leq 1$ in the last estimate. From (12) we now obtain

$$\begin{aligned} |z^\gamma \partial_x^\alpha \partial_z^\beta k_j(x, z)| &\leq C_{\alpha, \beta, \gamma} \cdot 2^{j(m+\delta|\alpha|+|\beta|-\rho|\gamma|)} \cdot 2^{jd} \\ &= C_{\alpha, \beta, \gamma} \cdot 2^{j(d+m+\delta|\alpha|+|\beta|-\rho|\gamma|)}, \end{aligned}$$

and by taking supremum over $|\gamma| = M$ (for instance, we can take γ such that $|z'| = (\max_{1 \leq i \leq d} |z_i|)^M$) we finally get (for $|\alpha| \leq N$, $|\beta| \geq 0$ and $\mathbb{N}_0 \ni M \leq N'$)

$$|\partial_x^\alpha \partial_z^\beta k_j(x, z)| \leq C_{\alpha, \beta, M} \cdot |z|^{-M} \cdot 2^{j(d+m+\delta|\alpha|+|\beta|-\rho M)}. \tag{13}$$

In order to estimate (11) we first consider the case $0 < |z| \leq 1$. We split (11) into two parts:

$$S_1 = \sum_{2^j \leq |z|^{-1}} |\partial_x^\alpha \partial_z^\beta k_j(x, z)| \quad \text{and} \quad S_2 = \sum_{2^j > |z|^{-1}} |\partial_x^\alpha \partial_z^\beta k_j(x, z)|.$$

Because of (13) (for $M = 0$) and $j \leq \log_2 |z|^{-1}$ we have

$$S_1 \leq C_{\alpha, \beta} \sum_{2^j \leq |z|^{-1}} 2^{j(d+m+\delta|\alpha|+|\beta|)} \leq C_{\alpha, \beta} \cdot \begin{cases} |z|^{-d-m-\delta|\alpha|-|\beta|}, & \text{if } d+m+\delta|\alpha|+|\beta| > 0 \\ 1 + \ln(|z|^{-1}), & \text{if } d+m+\delta|\alpha|+|\beta| \leq 0. \end{cases}$$

Using an elementary inequality $\ln(1+x) \leq \frac{1}{\alpha} x^\alpha$, valid for $\alpha \in (0, 1]$ and $x \geq 0$, we finally obtain

$$S_1 \leq C_{\alpha, \beta, L} \cdot |z|^{-d-m-\delta|\alpha|-|\beta|-L},$$

for all $L \geq 0$ such that $d+m+\delta|\alpha|+|\beta|+L > 0$. To estimate S_2 we use (13) with $M > \frac{d+m+\delta|\alpha|+|\beta|}{\rho}$:

$$\begin{aligned} S_2 &\leq C_{\alpha, \beta, M} \cdot |z|^{-M} \cdot \sum_{2^j > |z|^{-1}} 2^{j(d+m+\delta|\alpha|+|\beta|-\rho M)} \\ &\leq C_{\alpha, \beta, M} \cdot |z|^{-M-d-m-\delta|\alpha|-|\beta|+\rho M} \\ &\leq C_{\alpha, \beta, L} \cdot |z|^{-d-m-\delta|\alpha|-|\beta|-L}, \end{aligned}$$

for $L \geq (1-\rho)M$, which is exactly the requirement (10) and the proof is complete in the case $|z| \leq 1$.

To estimate (11) in the case $|z| > 1$ we again use (13) with $M > \frac{d+m+\delta|\alpha|+|\beta|}{\rho}$ to get (if we also take $M \geq d+m+\delta|\alpha|+|\beta|+L$, which is possible under requirements of the lemma)

$$\sum_{j=0}^{\infty} |\partial_x^\alpha \partial_z^\beta k_j(x, z)| \leq C_{\alpha, \beta, M} \cdot |z|^{-M} \leq C_{\alpha, \beta, L} \cdot |z|^{-d-m-\delta|\alpha|-|\beta|-L}.$$

□

If $N' > \left(\frac{d+m}{\rho}\right)^+$, the bound from the previous lemma provides us with the following integral representation

$$T_\sigma f(x) = \int_{\mathbb{R}^d} k(x, x-y) f(y) dy, \tag{14}$$

for any $f \in C_c^\infty(\mathbb{R}^d)$ and $x \notin \text{supp} f$. Obviously, we can also take $f \in L_c^\infty(\mathbb{R}^d)$ and the representation is then valid for a.e. $x \notin \text{supp} f$.

Moreover, because of the L^2 continuity result by Coifman and Meyer mentioned in the Introduction and density of $C_c^\infty(\mathbb{R}^d)$ in $L^2(\mathbb{R}^d)$, we can easily show that, for operators of order zero and sufficiently large N' , this representation remains valid for any $f \in L^2(\mathbb{R}^d)$ and a.e. $x \notin \text{supp} f$.

We are now in a position to prove the main theorem of this paper.

Theorem 3.2. *Let $\sigma \in S_{\rho, \delta, N, N'}^m$, $[0, 1) \ni \delta \leq \rho \in (0, 1]$ and $m \leq -(1-\rho)(d+1+\rho)$. If*

$$N > \frac{(3-\delta)d + (5-\delta)(1-\delta)}{(1-\delta)^2}, \quad N' > 6d + 12,$$

then T_σ is bounded on $L^p(\mathbb{R}^d)$, $\mathbf{p} \in (1, \infty)^d$.

Proof. Because of Theorem 2.1 and known L^2 continuity result, we only need to show that T_σ and T_{σ^*} satisfy the estimate (2). Moreover, because of (3)-(4) it is enough to show that T_σ satisfies the estimate (2) – the result for T_{σ^*} is then an easy consequence.

We now follow the proof of [3, Lemma 3]. For an arbitrary $N > 1$, using (14), two linear changes of variables, and the Minkowski inequality, we get

$$\begin{aligned} & \int_{|x'-x'_0|_\infty > Nt} \|T_\sigma f(\cdot, x')\|_{\mathbb{P}} dx' \\ &= \int_{|x'-x'_0|_\infty > Nt} \left\| \int \int k(\cdot, x', \cdot - \bar{y}, x' - y') f(\bar{y}, y') d\bar{y} dy' \right\|_{\mathbb{P}} dx' \\ &= \int_{|x'|_\infty > Nt} \left\| \int \int (k(\cdot, x' + x'_0, \cdot - \bar{y}, x' - y') - k(\cdot, x' + x'_0, \cdot - \bar{y}, x')) f(\bar{y}, y' + x'_0) d\bar{y} dy' \right\|_{\mathbb{P}} dx' \\ &= \int_{|x'|_\infty > Nt} \left\| \int \int (k(\cdot, x' + x'_0, \bar{y}, x' - y') - k(\cdot, x' + x'_0, \bar{y}, x')) f(\cdot - \bar{y}, y' + x'_0) d\bar{y} dy' \right\|_{\mathbb{P}} dx' \\ &\leq \int_{|x'|_\infty > Nt} \int \int \left\| (k(\cdot, x' + x'_0, \bar{y}, x' - y') - k(\cdot, x' + x'_0, \bar{y}, x')) f(\cdot - \bar{y}, y' + x'_0) \right\|_{\mathbb{P}} d\bar{y} dy' dx' = I. \end{aligned}$$

Furthermore, using (9) and the Mean value theorem, for $|x'|_\infty > Nt$ and $|y'|_\infty \leq t$ we have an estimate

$$\begin{aligned} |k(\bar{x}, x' + x'_0, \bar{y}, x' - y') - k(\bar{x}, x' + x'_0, \bar{y}, x')| &= |\nabla_{y'} k(\bar{x}, x' + x'_0, \bar{y}, x' - \vartheta y') \cdot y'| \\ &\leq C |(\bar{y}, x' - \vartheta y')|^{-d-m-1-L} |y'|_\infty \\ &\leq C |(\bar{y}, x' - \vartheta y')|^{-d-m-1-L} t, \end{aligned}$$

for some constants $\vartheta \in (0, 1)$, $C > 0$ and L satisfying assumptions of Lemma 3.1 (the optimal value of L will be determined later). Now, using the assumption on the support of function f we continue the estimate

$$\begin{aligned} I &\leq Ct \int_{|x'|_\infty > Nt} \int_{|y'|_\infty \leq t} \int |(\bar{y}, x' - \vartheta y')|^{-d-m-1-L} \|f(\cdot - \bar{y}, y' + x'_0)\|_{\mathbb{P}} d\bar{y} dy' dx' \\ &= Ct \int_{|x'|_\infty > Nt} \int_{|y'|_\infty \leq t} \int |(\bar{y}, x' - \vartheta y')|^{-d-m-1-L} \|f(\cdot, y' + x'_0)\|_{\mathbb{P}} d\bar{y} dy' dx' \\ &= Ct \int_{|y'|_\infty \leq t} \|f(\cdot, y' + x'_0)\|_{\mathbb{P}} \int_{|x'|_\infty > Nt} \int |(\bar{y}, x' - \vartheta y')|^{-d-m-1-L} d\bar{y} dx' dy'. \end{aligned}$$

To conclude the proof we need to check that

$$II = t \int_{|x'|_\infty > Nt} \int |(\bar{y}, x' - \vartheta y')|^{-d-m-1-L} d\bar{y} dx'$$

is bounded on $|y'|_\infty \leq t$, for arbitrary $N > d$. Indeed, we have

$$\begin{aligned} d^{-d-1} \mathbb{I} &\leq t \int_{|x'| > Nt} \int (\|\bar{y}\|_1 + |x' - \mathfrak{D}y'|_1)^{-d-m-1-L} d\bar{y} dx' \\ &\leq t \int_{|x'| > Nt} \int (\|\bar{y}\|_1 + |x'|_1 - |y'|_1)^{-d-m-1-L} d\bar{y} dx' \\ &\leq t \int_{|x'| > Nt} \int (\|\bar{y}\|_1 + |x'|_1 - dt)^{-d-m-1-L} d\bar{y} dx' \\ &= t^{-m-L} \int_{|x'| > N} \int (\|\bar{y}\|_1 + |x'|_1 - d)^{-d-m-1-L} d\bar{y} dx', \end{aligned}$$

where in the last step we used a linear change of variables. It is now obvious that this can be bounded (uniformly for $t > 0$) only in the case $L = -m$ and in that case, the above integral is indeed finite.

Thus, according to Lemma 3.1 we need $N' \geq d + 1$, $N' > \frac{d+m+1}{\rho}$ and

$$-m \geq (1 - \rho) \left(\left\lfloor \frac{d+m+1}{\rho} \right\rfloor + 1 \right)^+.$$

The last requirement is obviously valid in the case $m < -d - 1$. Otherwise, it is sufficient to have

$$-m \geq (1 - \rho) \left(\frac{d+m+1}{\rho} + 1 \right),$$

which is equivalent to $m \leq -(1 - \rho)(d + 1 + \rho)$ – the assumption stated in the theorem.

Of course, the same estimate should be valid also for T_σ . Because of (3)-(4) this means that we also need $M' \geq d + 1$ and $M' > \frac{d+m+1}{\rho}$, where M, M' are defined there. In the end, we notice that it is enough to take $M = 0$ and that the assumptions of this theorem are sufficient in that case. \square

References

- [1] J. Aleksić, S. Pilipović, I. Vojnović, *H-distributions with unbounded multipliers*, J. Pseudo-Differ. Oper. Appl. 9, 615–641 (2018)
- [2] N. AntoniĆ, I. Ivec, *On the Hörmander-Mihlin theorem for mixed-norm Lebesgue spaces*, J. Math. Anal. Appl. 433, 176–199 (2016)
- [3] N. AntoniĆ, I. Ivec, I. Vojnović, *Continuity of pseudodifferential operators on mixed-norm Lebesgue spaces*, Monatshefte für Mathematik 190(4), 657–674 (2019)
- [4] N. AntoniĆ, D. Mitrović, *H-distributions: an extension of H-measures to an $L^p - L^q$ setting*, Abs. Appl. Analysis 2011 Article ID 901084 (2011)
- [5] A. I. Benedek, R. Panzone, *The spaces L^p , with mixed norm*, Duke Math. J. 28, 301–324 (1961)
- [6] D. Cardona, J. Delgado, M. Ruzhansky, *L^p -Bounds for Pseudo-differential Operators on Graded Lie Groups*, The Journal of Geometric Analysis 31:11603–11647 (2021)
- [7] R. R. Coifman, Y. Meyer, *Àu-delà des opérateurs pseudo-différentiels*, Astérisque 57, (1978)
- [8] E. Cordero, D. Zucco, *Strichartz Estimates for the Schrödinger Equation*, CUBO A Math. J. 12(3), 213–239 (2010)
- [9] J. Delgado, M. Ruzhansky, *L^p -Bounds for Pseudo-differential Operators on Compact Lie Groups*, J. Inst. Math. Jussieu 18(3), 531–559 (2019)
- [10] L. Hörmander, *Pseudo-differential operators and hypoelliptic equations*, In Proc. Symposium on Singular Integrals, Amer. Math. Soc., Providence RI 10, 138–183 (1967)
- [11] I. Ivec, I. Vojnović, *Basic calculus of pseudodifferential operators with nonsmooth symbols*, J. Pseudo-Differ. Oper. Appl. 13 (2022), <https://doi.org/10.1007/s11868-022-00443-0>
- [12] I. Ivec, I. Vojnović, *H-distributions on Hörmander spaces*, J. Math. Anal. Appl. 495(1), 124717 (2021)
- [13] E. M. Stein, *Harmonic analysis*, Princeton University Press, (1993)