



SEP elements and solutions of related equations in a ring with involution

Anqi Li^a, Mengge Guan^a, Junchao Wei^a

^aCollege of mathematical science, Yangzhou University, Yangzhou, Jiangsu 225002, P. R. China

Abstract. In this paper, we study many new characterizations of *SEP* elements in a rings with involution, mainly, we first discuss some properties of *SEP* elements by means of regular elements, $\{1,3\}$ -inverses and the equality of certain set. Next, we give some necessity and sufficiency conditions of *SEP* elements by discussing the solutions of related equations. Finally, we characterize *SEP* elements by constructing the group inverses and *MP* inverses.

1. Introduction

Let R be a ring and $a \in R$. If there exists $b \in R$ such that $a = aba$, then a is called a regular element, and b is called the inner inverse element of a . Clearly, bab is also an inner inverse of a . We denote the set of all inner inverses of a by $a\{1\}$, and a^{-} is an inner inverse of a , and denote the set of all regular element of R by R^{reg} .

If there exists $a^\# \in R$ such that

$$aa^\#a = a, a^\#aa^\# = a^\#, aa^\# = a^\#a,$$

the element a is called group invertible element and $a^\#$ is called the group inverse of a [4, 8, 9], and it is uniquely determined by these equations. We write $R^\#$ to denote the set of all group invertible elements of R .

If a map $*$: $R \rightarrow R$ satisfies

$$(a^*)^* = a, (a + b)^* = a^* + b^*, (ab)^* = b^*a^*,$$

then R is said to be an involution ring or a $*$ -ring.

Let R be a $*$ -ring and $a \in R$. If the following equations:

$$a = axa, ax = (ax)^*$$

have a common solution, then a is called $\{1,3\}$ invertible element of R , and such solution is called the $\{1,3\}$ -inverse of a . If the following equations:

$$a = axa, xa = (xa)^*$$

2020 Mathematics Subject Classification. 16B99, 16W10, 46L05

Keywords. EP element, SEP element, solutions of equation, regular, ring with involution.

Received: 11 August 2022; Revised: 20 April 2023; Accepted: 23 April 2023

Communicated by Dijana Mosić

Email addresses: angel1ee_yz@126.com (Anqi Li), 2530374647@qq.com (Mengge Guan), jcwei@yzu.edu.cn (Junchao Wei)

have a common solution, then a is called $\{1,4\}$ invertible element of R , and such solution is called the $\{1,4\}$ -inverse of a .

Let R be a $*$ -ring and $a \in R$. If there exists $a^+ \in R$ such that

$$a = aa^+a, a^+ = a^+aa^+, (aa^+)^* = aa^+, (a^+a)^* = a^+a,$$

then a is called Moore Penrose invertible element, and a^+ is called the Moore Penrose inverse of a [3, 6]. Let R^+ denote the set of all Moore Penrose invertible elements of R .

If $a \in R^\# \cap R^+$ and $a^\# = a^+$, then a is called EP element. On the studies of EP , the readers can refer to [2, 3, 5, 7, 10–14, 20, 21].

If $a = aa^*a$, then a is called partial isometry [15, 18]. It is known that $a \in R$ is a partial isometry if and only if $a \in R^+$ and $a^* = a^+$ [11].

$a \in R^\# \cap R^+$ is called SEP element if $a^\# = a^+ = a^*$. Clearly, $a \in R^\# \cap R^+$ is SEP if and only if $a \in R^{EP}$ and $a \in R^{PI}$. Where R^{EP} , R^{PI} and R^{SEP} are denoted the set of all EP elements, all PI elements and all SEP elements of R respectively.

In [11], many characterizations of SEP elements are given. In [19], it is shown that $a \in R^\# \cap R^+$ is SEP if and only if the equation $xa^+a = xaa^*$ has at least one solution in $\chi_a =: \{a, a^\#, a^+, a^*, (a^+)^*, (a^\#)^*\}$. In [16], it is proved that $a \in R^\# \cap R^+$ is SEP if and only if the equation $yx a^\# = yx a^*$ has at least one solution in $\chi_a^2 =: \{(x, y) | x, y \in \chi_a\}$. In [18], it is shown that $a \in R^\# \cap R^+$ is SEP if and only if the equation $a^*xa = aa^+x$ has at least one solution in χ_a .

Motivated by these results, this paper mainly study the ways to characterize SEP elements.

2. Some characterizations of SEP elements

Let $a \in R^\# \cap R^+$. Then, clearly, $a \in R^{SEP}$ if and only if $a^2a^+ = a = (a^+)^*$, this induces us to give the following characterization of SEP elements.

Theorem 2.1. *Let $a \in R^\# \cap R^+$. Then $a \in R^{SEP}$ if and only if $(a^2a^+ - (a^+)^*)^2 = a(a^2a^+ - (a^+)^*)$.*

Proof. (\Rightarrow) It is evident because $a^2a^+ - (a^+)^* = 0$.

(\Leftarrow) Assume that $(a^2a^+ - (a^+)^*)^2 = a(a^2a^+ - (a^+)^*)$. Then

$$a^3a^+ - a(a^+)^* - (a^+)^*a^2a^+ + (a^+)^*(a^+)^* = a^3a^+ - a(a^+)^*,$$

this gives $(a^+)^*a^2a^+ = (a^+)^*(a^+)^*$. Multiplying the equality on the left by a^* , one has $a^+a^3a^+ = a^+a(a^+)^*$. Again multiply the last equality on the left by $aa^\#$, one gets

$$a^2a^+ = (a^+)^*,$$

it follows that $(a^+)^* = (a^+)^*aa^\# = (a^2a^+)aa^\# = a$. Hence $a \in R^{PI}$ and $a^2a^+ = (a^+)^* = a$, this infers $a \in R^{EP}$ by [11, Theorem 1.2.1]. Thus $a \in R^{SEP}$. \square

The following corollary is an immediate result of Theorem 2.1.

Corollary 2.2. *Let $a \in R^\# \cap R^+$. Then $a \in R^{SEP}$ if and only if $a^2a^+ = (a^+)^*$.*

Noting that $(a^2a^+)^+ = aa^\#a^+$. Hence Corollary 2.2 leads to the following theorem.

Theorem 2.3. *Let $a \in R^\# \cap R^+$. Then $a \in R^{SEP}$ if and only if $aa^\#a^+ = a^*$.*

Since $a \in R^{SEP}$ if and only if $a^* \in R^{SEP}$, replacing a in Corollary 2.2 by a^* , one obtains

Corollary 2.4. [11, Theorem 1.5.3] *Let $a \in R^\# \cap R^+$. Then $a \in R^{SEP}$ if and only if $a^*a^+a = a^+$.*

Noting that $a^* = a^+aa^*$. Then Corollary 2.4 induces the following result.

Corollary 2.5. Let $a \in R^\# \cap R^+$. Then $a \in R^{SEP}$ if and only if $aa^*a^+ = aa^+$.

Theorem 2.6. Let $a \in R^\# \cap R^+$. Then $a \in R^{SEP}$ if and only if $(a^\#)^*a^+ = aa^+$.

Proof. Noting that $(aa^*a^+)^+ = (a^\#)^*a^+$ and $(aa^+)^+ = aa^+$. Hence we are done by Corollary 2.5. \square

Theorem 2.7. Let $a \in R^\# \cap R^+$. Then $a \in R^{SEP}$ if and only if $a^\#a(a^\#)^*a^+aa^\# = aa^+$.

Proof. Since $(aa^*a^+)^{\#} = a^\#a(a^\#)^*a^+aa^\#$ and $(aa^+)^{\#} = aa^+$, we are done by Corollary 2.5. \square

3. Construct regular elements to characterize SEP elements

From Theorem 2.3, we have the following lemma.

Lemma 3.1. Let $a \in R^\# \cap R^+$. Then $a \in R^{SEP}$ if and only if $a = aa^+(a^\#)^*$.

Theorem 3.2. Let $a \in R^\# \cap R^+$. Then $a \in R^{EP}$ if and only if $\begin{pmatrix} aa^+(a^\#)^* \\ a \end{pmatrix}$ is a regular element and $(aa^+a^* \ 1 - aa^+) \in \begin{pmatrix} aa^+(a^\#)^* \\ a \end{pmatrix}\{1\}$.

Proof. (\Rightarrow) If $a \in R^{EP}$, then $aaa^+ = aaa^\# = a$. It follows that

$$\begin{pmatrix} aa^+(a^\#)^* \\ a \end{pmatrix} (aa^+a^* \ 1 - aa^+) \begin{pmatrix} aa^+(a^\#)^* \\ a \end{pmatrix} = \begin{pmatrix} aa^+(a^\#)^* \\ aaa^+ \end{pmatrix} = \begin{pmatrix} aa^+(a^\#)^* \\ a \end{pmatrix}.$$

Hence $\begin{pmatrix} aa^+(a^\#)^* \\ a \end{pmatrix}$ is a regular element and $(aa^+a^* \ 1 - aa^+) \in \begin{pmatrix} aa^+(a^\#)^* \\ a \end{pmatrix}\{1\}$.

(\Leftarrow) If $\begin{pmatrix} aa^+(a^\#)^* \\ a \end{pmatrix}$ is a regular element and $(aa^+a^* \ 1 - aa^+) \in \begin{pmatrix} aa^+(a^\#)^* \\ a \end{pmatrix}\{1\}$, then

$$\begin{pmatrix} aa^+(a^\#)^* \\ a \end{pmatrix} = \begin{pmatrix} aa^+(a^\#)^* \\ a \end{pmatrix} (aa^+a^* \ 1 - aa^+) \begin{pmatrix} aa^+(a^\#)^* \\ a \end{pmatrix} = \begin{pmatrix} aa^+(a^\#)^* \\ aaa^+ \end{pmatrix},$$

it follows that $a^2a^+ = a$. Hence $a \in R^{EP}$. \square

Theorem 3.3. Let $a \in R^\# \cap R^+$. Then $a \in R^{SEP}$ if and only if $(a^+ \ 1 - aa^+) \in \begin{pmatrix} aa^+(a^\#)^* \\ a \end{pmatrix}\{1\}$.

Proof. (\Rightarrow) If $a \in R^{SEP}$, then $a \in R^{EP}$ and $a = aa^+(a^\#)^*$ by Lemma 3.1.

This gives $(aa^+a^* \ 1 - aa^+) \in \begin{pmatrix} aa^+(a^\#)^* \\ a \end{pmatrix}\{1\}$ by Theorem 3.2.

Noting that $a^+ = (aa^+(a^\#)^*)^+ = aa^+a^*$. Then $(a^+ \ 1 - aa^+) \in \begin{pmatrix} aa^+(a^\#)^* \\ a \end{pmatrix}\{1\}$.

(\Leftarrow) From the assumption, one has

$$\begin{pmatrix} aa^+(a^\#)^* \\ a \end{pmatrix} = \begin{pmatrix} aa^+(a^\#)^* \\ a \end{pmatrix} (a^+ \ 1 - aa^+) \begin{pmatrix} aa^+(a^\#)^* \\ a \end{pmatrix} = \begin{pmatrix} aa^+(a^\#)^*a^+(a^\#)^* \\ aa^+(a^\#)^* \end{pmatrix}.$$

This infers $a = aa^+(a^\#)^*$. By Lemma 3.1, $a \in R^{SEP}$. \square

Theorem 3.4. Let $a \in R^\# \cap R^+$. Then $a \in R^{SEP}$ if and only if $(aa^+a^* \ 1 - (a^+)^*a^\#) \in \begin{pmatrix} aa^+(a^\#)^* \\ a \end{pmatrix}\{1\}$.

Proof. (\Rightarrow) If $a \in R^{SEP}$, then $a = (a^+)^*$ and $a^\# = a^+$. It follows that $aa^+ = (a^+)^*a^\#$. Hence $\begin{pmatrix} aa^+a^* & 1 - (a^+)^*a^\# \\ aa^+(a^\#)^* & a \end{pmatrix} \in \left\{ \begin{pmatrix} aa^+(a^\#)^* & \\ a & \end{pmatrix} \{1\} \right\}$ by Theorem 3.2.

(\Leftarrow) From the assumption, one has

$$\begin{aligned} \begin{pmatrix} aa^+(a^\#)^* & \\ a & \end{pmatrix} &= \begin{pmatrix} aa^+(a^\#)^* & \\ a & \end{pmatrix} \begin{pmatrix} aa^+a^* & 1 - (a^+)^*a^\# \\ & \end{pmatrix} \begin{pmatrix} aa^+(a^\#)^* & \\ a & \end{pmatrix} \\ &= \begin{pmatrix} aa^+(a^\#)^* + aa^+(a^\#)^*(a - (a^+)^*) & \\ a^2a^+ + a(a - (a^+)^*) & \end{pmatrix}. \end{aligned}$$

This gives

$$aa^+(a^\#)^* = aa^+(a^\#)^* + aa^+(a^\#)^*(a - (a^+)^*). \tag{3.1}$$

$$a = a^2a^+ + a(a - (a^+)^*). \tag{3.2}$$

Multiplying the Eq.(3.2) on the right by a^+a , one obtains $a^2a^+ = a^2a^+a^+a$, this gives

$$a^* = a^*aa^+ = a^*a^\#a^2a^+ = a^*a^\#a^2a^+a^+a = a^*a^+a,$$

and

$$(aa^\#)^* = (a^\#)^*a^* = (a^\#)^*(a^*a^+a) = a^+a.$$

Hence $a \in R^{EP}$ by [11, Theorem 1.1.3], this implies $aa^+(a^\#)^* = a^+a(a^\#)^* = (a^\#)^*$. From the Eq.(3.1), one gets $(a^\#)^*(a - (a^+)^*) = 0$, that is, $a^*a^\# = a^+a^\#$. Hence $a \in R^{PI}$ by [11, Theorem 1.5.2]. Thus $a \in R^{SEP}$. \square

4. Using the equality of sets to characterize SEP elements

Let $a, b \in R$. We set $a \vee b = \{a^2, ab, ba, b^2\}$, and use this to characterize SEP elements as follows.

Theorem 4.1. *Let $a \in R^\# \cap R^+$. Then $a \in R^{SEP}$ if and only if $a^+a^2 \vee a = a^+a^2 \vee aa^+(a^\#)^*$.*

Proof. (\Rightarrow) Assume that $a \in R^{SEP}$. Then $a = aa^+(a^\#)^*$ by Lemma 3.1. Hence $a^+a^2 \vee a = a^+a^2 \vee aa^+(a^\#)^*$.

(\Leftarrow) Noting that $a^+a^2 \vee a = \{a^+a^3, a^2\}$,

$$a^+a^2 \vee aa^+(a^\#)^* = \{a^+a^3, a^+a^3a^+(a^\#)^*, aa^+(a^\#)^*a^+a^2, aa^+(a^\#)^*(a^\#)^*\}.$$

By hypothesis, we have $\{a^+a^3, a^2\} = \{a^+a^3, a^+a^3a^+(a^\#)^*, aa^+(a^\#)^*a^+a^2, aa^+(a^\#)^*(a^\#)^*\}$.

(1) If $a^+a^3 = a^2$, then we have $a^+a^3 = a^+a^3a^+(a^\#)^* = aa^+(a^\#)^*a^+a^2 = aa^+(a^\#)^*(a^\#)^*$. Since $a^+a^3 = a^2$, $a \in R^{EP}$ by [11], this infers

$$a^2 = a^+a^3 = a^+a^3a^+(a^\#)^* = a^2a^+(a^\#)^* = a(a^\#)^*,$$

and

$$a = a^+a^2 = a^+a(a^\#)^* = (a^\#)^*.$$

Hence $a \in R^{SEP}$.

(2) If $a^+a^3 \neq a^2$, then we can discuss the following cases.

① $a^2 = aa^+(a^\#)^*a^+a^2$. In this case, we have

$$a^+a^3 = a^+a^3a^+(a^\#)^* \text{ and } aa^+(a^\#)^*a^+a^2 = aa^+(a^\#)^*(a^\#)^*,$$

or

$$a^+a^3 = a^+a^3a^+(a^\#)^* = aa^+(a^\#)^*(a^\#)^*,$$

or

$$a^+a^3 = aa^+(a^\#)^*(a^\#)^* \text{ and } a^+a^3a^+(a^\#)^* = aa^+(a^\#)^*a^+a^2,$$

or

$$a^+ a^3 a^+ (a^\#)^* = aa^+ (a^\#)^* (a^\#)^* = aa^+ (a^\#)^* a^+ a^2.$$

In the first case and the second case, we have

$$a^+ a^3 = a^+ a^3 a^+ (a^\#)^* = (a^+ a^3 a^+ (a^\#)^*) aa^+ = a^+ a^4 a^+,$$

and

$$a = a^\# a^+ a^3 = a^\# (a^+ a^4 a^+) = a^2 a^+.$$

So $a \in R^{EP}$, one obtains $a^+ a^3 = a^2$, which is a contradiction.

In the third case, we also have $a^+ a^3 = aa^+ (a^\#)^* (a^\#)^* = aa^+ (a^\#)^* (a^\#)^* aa^+ = a^+ a^4 a^+$. So $a \in R^{EP}$, and $a^+ a^3 = a^2$, which is also a contradiction.

In the fourth case, we have $a^2 = a^+ a^3 a^+ (a^\#)^*$. Multiplying the equality on the left by $a^\#$, one has $a = aa^+ (a^\#)^*$. Hence $a \in R^{SEP}$ by Lemma 3.1.

② $a^2 = aa^+ (a^\#)^* (a^\#)^*$. Then $a^3 a^+ = aa^+ (a^\#)^* (a^\#)^* aa^+ = aa^+ (a^\#)^* (a^\#)^* = a^2$. It follows that $a \in R^{EP}$, this gives $a^+ a^3 = a^2$, which is a contradiction.

③ $a^2 = a^+ a^3 a^+ (a^\#)^*$. From the proof of the fourth case in ① we get $a \in R^{SEP}$. Therefore, in any case, we have $a \in R^{SEP}$. \square

Similarly, we have the following theorems.

Theorem 4.2. Let $a \in R^\# \cap R^+$. Then $a \in R^{SEP}$ if and only if $a^2 a^+ \vee a = a^2 a^+ \vee aa^+ (a^\#)^*$.

Theorem 4.3. Let $a \in R^\# \cap R^+$. Then $a \in R^{SEP}$ if and only if $a^* a^+ a \vee a^* = a^* a^+ a \vee a^\# aa^+$.

5. Using the (1, 3)-inverse and (1, 4)-inverse to characterize SEP elements

Theorem 5.1. Let $a \in R^\# \cap R^+$. Then $a \in R^{SEP}$ if and only if $aa^+ (a^\#)^* \in a^+ \{1, 3\}$.

Proof. (\Rightarrow) Assume that $a \in R^{SEP}$. Then $a = aa^+ (a^\#)^*$ by Lemma 3.1. Noting that $a \in a^+ \{1, 3\}$. Then $aa^+ (a^\#)^* \in a^+ \{1, 3\}$.

(\Leftarrow) If $aa^+ (a^\#)^* \in a^+ \{1, 3\}$, then

$$a^+ (aa^+ (a^\#)^*) a^+ = a^+. \tag{5.1}$$

$$a^+ (aa^+ (a^\#)^*) = (a^+ (aa^+ (a^\#)^*))^*. \tag{5.2}$$

By the Eq.(5.2), one has $a^+ (a^\#)^* = a^\# (a^+)^*$. Multiplying the equality on the right by a^* , one obtains $a^+ = a^\# aa^+$. Hence $a \in R^{EP}$ by [11, Theorem 1.2.1]. By the Eq.(5.1), one gets $a^+ = a^+ (a^\#)^* a^+$, this gives $a = aa^+ a = aa^+ (a^\#)^* a^+ a = (a^+)^*$. Hence $a \in R^{PI}$ and so $a \in R^{SEP}$. \square

Corollary 5.2. Let $a \in R^\# \cap R^+$. Then $a \in R^{SEP}$ if and only if $a^+ = a^\# (a^+)^* a^+$.

Proof. (\Rightarrow) If $a \in R^{SEP}$, then $aa^+ (a^\#)^* \in a^+ \{1, 3\}$ by Theorem 5.1. It follows that $a^+ = (aa^+ (a^\#)^*)^* (a^+)^* a^+ = a^\# aa^+ (a^+)^* a^+ = a^\# (a^+)^* a^+$.

(\Leftarrow) Assume that $a^+ = a^\# (a^+)^* a^+$. Then

$$aa^+ = aa^\# (a^+)^* a^+ = (a^+)^* a^+ \text{ and } a^* = a^* aa^+ = a^* (a^+)^* a^+ = a^+.$$

It follows that $a^+ a = (a^\# (a^+)^* a^+) a = a^\# (a^+)^* = a^\# a$. Hence $a \in R^{EP}$ and so $a \in R^{SEP}$. \square

Theorem 5.3. Let $a \in R^\# \cap R^+$. Then $a \in R^{SEP}$ if and only if $a^+ \in aa^+ (a^\#)^* \{1, 4\}$.

Proof. (⇒) If $a \in R^{SEP}$. Then $a = aa^+(a^\#)^*$ by Lemma 3.1. Noting that $a^+ \in a\{1, 4\}$. Then $a^+ \in aa^+(a^\#)^*\{1, 4\}$.

(⇐) If $a^+ \in aa^+(a^\#)^*\{1, 4\}$, then

$$aa^+(a^\#)^* = aa^+(a^\#)^*a^+aa^+(a^\#)^*. \tag{5.3}$$

$$(a^+aa^+(a^\#)^*)^* = a^+aa^+(a^\#)^*. \tag{5.4}$$

By the Eq.(5.3), one yields $aa^+(a^\#)^* = (a^+)^*a^+(a^\#)^*$. Multiplying the equality on the right by a^* , one has $aa^+ = (a^+)^*a^+$. Hence $a \in R^{PI}$. By the Eq.(5.4), one obtains

$$a^\#(a^+)^* = a^+(a^\#)^*.$$

By Theorem 5.1, we have $a \in R^{EP}$. Hence $a \in R^{SEP}$. □

Theorem 5.4. Let $a \in R^\# \cap R^+$. Then $a \in R^{SEP}$ if and only if $(a^\#)^*a^+ - aa^+ \in l(a^+) = \{x|x \in R, xa^+ = 0\}$.

Proof. (⇒) If $a \in R^{SEP}$, then we have $a = aa^+(a^\#)^*$ by Lemma 3.1. Noting that

$$aa^+(a^\#)^*a^+ = (a^+)^*a^+ \text{ and } a^+ = a^\#.$$

Then $(a^\#)^*a^+ - aa^+ = aa^+(a^\#)^*a^+ - aa^+ = aa^+ - aa^+ = 0 \in l(a^+)$.

(⇐) If $(a^\#)^*a^+ - aa^+ \in l(a^+)$, then $(a^\#)^*a^+a^+ = aa^+a^+$. By [2, Lemma 2.10], we have $(a^\#)^*a^+ = aa^+$. It follows that $a^+ = a^*(a^\#)^*a^+ = a^*aa^+ = a^*$, and so

$$(aa^\#)^* = (a^\#)^*a^* = (a^\#)^*a^+ = aa^+.$$

Hence $a \in R^{SEP}$ by [1, Theorem 1.3.1]. □

6. Characterizing SEP elements by the solution of univariate equations in a given set

When $a \in R^{SEP}$, we have $aa^+(a^\#)^* = a = aa^+a$ by Lemma 3.1. Hence we can construct the following equation:

$$xa^+(a^\#)^* = aa^+x. \tag{6.1}$$

Theorem 6.1. Let $a \in R^\# \cap R^+$. Then $a \in R^{SEP}$ if and only if Eq.(6.1) has at least one solution in $\rho_a = \{a, a^\#, a^+, a^*, (a^+)^*, (a^\#)^*, (a^+)^\#, (a^\#)^+\}$.

Proof. (⇒) If $a \in R^{SEP}$, then $x = a$ is a solution by Lemma 3.1.

(⇐) (1) If $x = a$ is a solution, then $aa^+(a^\#)^* = aa^+a = a$. Hence $a \in R^{SEP}$ by Lemma 3.1;

(2) If $x = a^\#$, then $a^\#a^+(a^\#)^* = aa^+a^\# = a^\#$. Multiplying the equality by a^2 on the left, we have $a = aa^+(a^\#)^*$. Hence $a \in R^{SEP}$ by Lemma 3.1;

(3) If $x = a^+$, then $a^+a^+(a^\#)^* = aa^+a^+$. Multiplying the equality on the right by a^* , one has $a^+a^+ = aa^+a^+a^*$. Again multiply the last equality by a^+ on the left, one gets

$$a^+a^+a^+ = a^+a^+a^*.$$

Hence $a^+a^+ = a^+a^* = a^+a^+aa^*$ and $a^+ = a^+aa^* = a^*$ by [2, Lemma 2.10]. It follows that $a^+ = a^+a^*(a^\#)^* = a^+a^+(a^\#)^* = aa^+a^+$, and $a^* = a^+aa^* = (aa^+a^+)aa^* = aa^+a^*$. Taking involution of the above equality, we obtains that $a = a^2a^+$. Hence $a \in R^{EP}$ by [11, Theorem 1.2.1]. Thus $a \in R^{SEP}$;

(4) If $x = a^*$, then $a^*a^+(a^\#)^* = aa^+a^*$. Multiplying the equality on the left by $(aa^\#)^*$, one yields $a^*a^+(a^\#)^* = a^*$. It follows that $a^* = aa^+a^*$. Hence $a \in R^{EP}$. Now we have $aa^+ = (a^+)^*a^* = (a^+)^*a^*a^+(a^\#)^* = aa^+a^+(a^\#)^* = a^+(a^\#)^*$, and $a = a^2a^+ = aa^+(a^\#)^*$. Thus $a \in R^{SEP}$ by Lemma 3.1;

(5) If $x = (a^+)^*$, then $(a^+)^*a^+(a^\#)^* = aa^+(a^+)^* = (a^+)^*$. Multiplying the equality on the left by aa^* , one gets $aa^+(a^\#)^* = a$. Hence $a \in R^{SEP}$ by Lemma 3.1;

(6) If $x = (a^\#)^*$, then $(a^\#)^*a^+(a^\#)^* = aa^+(a^\#)^*$. Multiplying the equality on the left by a^*a^* , one yields $a^*a^+(a^\#)^* = a^*$. Hence $a \in R^{SEEP}$ by (4);

(7) If $x = (a^+)^{\#} = (aa^\#)^*a(aa^\#)^*$, then $(aa^\#)^*a(aa^\#)^*a^+(a^\#)^* = aa^+(aa^\#)^*a(aa^\#)^*$, e.g., $(a^\#)^* = a(aa^\#)^*$. Multiplying the equality on the left by a^*a^+ , one obtains $a^*a^+(a^\#)^* = a^*$. Hence $a \in R^{SEEP}$ by (4);

(8) If $x = (a^\#)^+ = a^+a^3a^+$, then $a^+a^3a^+a^+(a^\#)^* = aa^+a^+a^3a^+$. Multiplying the equality on the left by a^+a , one has $aa^+a^+a^3a^+ = a^+a^2a^+a^+a^3a^+$. Again multiply the last equality on the right by $a^\#$, one gets $aa^+a^+a = a^+a^2a^+a^+a$. It follows that

$$aa^+ = aa^+(a^+aa^*(a^\#)^*) = a^+a^2a^+a^+a(a^*(a^\#)^*) = a^+a^2a^+.$$

Hence $a \in R^{EP}$, this infers $x = (a^\#)^+ = a^+a^3a^+ = a$. Thus $a \in R^{SEEP}$ by (1). \square

Noting that $a \in R^{SEEP}$ if and only if $a^* \in R^{SEEP}$. Then replace a by a^* in Eq.(6.1), one gets

$$x(a^+)^*a^\# = a^+ax. \tag{6.2}$$

Theorem 6.2. Let $a \in R^\# \cap R^+$. Then $a \in R^{SEEP}$ if and only if Eq.(6.2) has at least one solution in ρ_a .

Since $a \in R^{SEEP}$ if and only if $a \in R^{EP}$ and $(a^\#)^* = (a^\#)^*(a^\#)^*a^* = a^2a^* = a^3a^+a^*$. Hence we can revise Eq.(6.1) as follows.

$$xa^+a^3a^+a^* = aa^+x. \tag{6.3}$$

Theorem 6.3. Let $a \in R^\# \cap R^+$. Then $a \in R^{SEEP}$ if and only if Eq.(6.3) has at least one solution in ρ_a .

Noting that when $a \in R^{EP}$, we have $a^+a^3a^+ = (a^+a)^*a(aa^+)^* = (aa^\#)^*a(aa^\#)^*$. Then Eq.(6.3) can be changed as follows.

$$x(aa^\#)^*aa^* = aa^+x. \tag{6.4}$$

Theorem 6.4. Let $a \in R^\# \cap R^+$. Then $a \in R^{SEEP}$ if and only if Eq.(6.4) has at least one solution in ρ_a .

7. The general solution of bivariate equations

Now we generalize Eq.(6.1) as follows

$$xa^+(a^\#)^* = aa^+y. \tag{7.1}$$

Theorem 7.1. Let $a \in R^\# \cap R^+$. Then the general solution of Eq.(7.1) is given by

$$\begin{cases} x = aa^+p + u - ua^+a \\ y = pa^+(a^\#)^* + v - aa^+v \end{cases}, p, u, v \in R. \tag{7.2}$$

Proof. First $(aa^+p + u - ua^+a)a^+(a^\#)^* = aa^+pa^+(a^\#)^* = aa^+(pa^+(a^\#)^* + v - aa^+v)$. It follows that the formula (7.2) is the solution of Eq.(7.1).

Next, let $\begin{cases} x = x_0 \\ y = y_0 \end{cases}$ be any solution of Eq.(7.1). Then $x_0a^+(a^\#)^* = aa^+y_0$.

Choose $p = x_0, v = y_0$ and $u = x_0 - aa^+p$. Then

$$\begin{aligned} ua^+a &= (x_0 - aa^+p)a^+a = x_0a^+a - aa^+x_0a^+a = x_0a^+a - aa^+x_0(a^+(a^\#)^*a^*)a \\ &= x_0a^+a - aa^+(x_0a^+(a^\#)^*)a^*a = x_0a^+a - aa^+(aa^+y_0)a^*a = x_0a^+a - aa^+y_0a^*a \\ &= x_0a^+a - (x_0a^+(a^\#)^*)a^*a = x_0a^+a - x_0a^+a = 0. \end{aligned}$$

It follows that $x_0 = aa^+p + (x_0 - aa^+p) = aa^+p + u = aa^+p + u - ua^+a$. Noting that

$$aa^+v = aa^+y_0 = x_0a^+(a^\#)^* = pa^+(a^\#)^*.$$

Then $y_0 = pa^+(a^\#)^* + y_0 - aa^+v = pa^+(a^\#)^* + v - aa^+v$. Hence the general solution of Eq.(7.1) is given by the formula (7.2). \square

Theorem 7.2. Let $a \in R^\# \cap R^+$. Then $a \in R^{SEP}$ if and only if the general solution of Eq.(7.1) is given by

$$\begin{cases} x = aa^+p + u - ua^+a \\ y = pa^+a + v - aa^+v \end{cases}, p, u, v \in R. \tag{7.3}$$

Proof. (\Rightarrow) If $a \in R^{SEP}$, then $(a^\#)^* = a$. By Theorem 7.1, one gets the general solution of Eq.(7.1) is given by the formula (7.3).

(\Leftarrow) From the assumption, we get $(aa^+p + u - ua^+a)a^+(a^\#)^* = aa^+(pa^+a + v - aa^+v)$, e.g.,

$$aa^+pa^+(a^\#)^* = aa^+pa^+a \text{ for all } p \in R.$$

Choose $p = (a^+)^*$, then $(a^+)^*a^+(a^\#)^* = (a^+)^*$. Taking involution of the above equality, one obtains $a^\#(a^+)^*a^+ = a^+$. Multiplying the equality on the right by aa^+ , one has $aa^\#a^+ = a^+$. Again multiply the last equality on the right by a , one gets $aa^\# = a^*a$. Hence $a \in R^{SEP}$ by [11, Theorem 1.5.3]. \square

Theorem 7.3. Let $a \in R^\# \cap R^+$. Then $a \in R^{SEP}$ if and only if the general solution of Eq.(7.1) is given by

$$\begin{cases} x = a^*ap + u - ua^+a \\ y = pa^+(a^\#)^* + v - aa^+v \end{cases}, p, u, v \in R. \tag{7.4}$$

Proof. (\Rightarrow) If $a \in R^{SEP}$, then $a^*a = a^\#a = aa^\# = aa^+$. It follows from Theorem 7.1 that the general solution of Eq.(7.1) is given by the formula (7.4).

(\Leftarrow) From the assumption, we get $(a^*ap + u - ua^+a)a^+(a^\#)^* = aa^+(pa^+(a^\#)^* + v - aa^+v)$, e.g.,

$$a^*apa^+(a^\#)^* = aa^+pa^+(a^\#)^* \text{ for all } p \in R.$$

Choose $p = a$, then $a^*a^2a^+(a^\#)^* = aa^+(a^\#)^*$. Multiplying the equality on the right by $a^*aa^\#a^+$, one has $a^* = aa^\#a^+$. Hence $a \in R^{SEP}$. \square

We don't know the general solution of which equation is given by formula (7.4), for this we can construct the equation as follows.

$$xa^+(a^\#)^* = a^*a(aa^\#)^*y. \tag{7.5}$$

Theorem 7.4. Let $a \in R^\# \cap R^+$. Then the general solution of Eq.(7.5) is given by

$$\begin{cases} x = a^*ap + u - ua^+a \\ y = pa^+(a^\#)^* + v - aa^+v \end{cases}, \text{ where } p, u, v \in R \text{ with } a^+p = a^+a^+ap. \tag{7.6}$$

Proof. When $a^+p = a^+a^+ap$, we have $a^*a(a^\#)^*a^*p = a^*ap$. So

$$(a^*ap + u - ua^+a)a^+(a^\#)^* = a^*a(a^\#)^*a^*pa^+(a^\#)^* = a^*a(aa^\#)^*(pa^+(a^\#)^* + v - aa^+v).$$

It follows that the formula (7.6) is the solution of Eq.(7.5).

Next, let $\begin{cases} x = x_0 \\ y = y_0 \end{cases}$ be any solution of Eq.(7.5). Then $x_0a^+(a^\#)^* = a^*a(aa^\#)^*y_0$.

Choose $p = (aa^\#)^*y_0a^*a$, $v = y_0 - (aa^\#)^*y_0$ and $u = x_0$. Then

$$a^+p = a^+(aa^\#)^*y_0a^*a = a^+(a^+a(aa^\#)^*y_0a^*a) = a^+a^+ap,$$

$$ua^+a = x_0a^+a = x_0a^+(a^\#)^*a^*a = a^*a(aa^\#)^*y_0a^*a = a^*ap.$$

It follows that $x_0 = u = a^*ap + u - ua^+a$. Noting that

$$aa^+v = aa^+(y_0 - (aa^\#)^*y_0) = aa^+y_0 - aa^+(aa^\#)^*y_0 = aa^+y_0 - aa^+y_0 = 0,$$

and

$$\begin{aligned} pa^+(a^\#)^* &= (aa^\#)^*y_0a^*aa^+(a^\#)^* = (aa^\#)^*y_0(aa^\#)^* = a^+a(aa^\#)^*y_0(aa^\#)^* \\ &= a^+(a^\#)^*a^*a(aa^\#)^*y_0(aa^\#)^* = a^+(a^\#)^*x_0a^+(a^\#)^*(aa^\#)^* = a^+(a^\#)^*x_0a^+(a^\#)^* \\ &= a^+(a^\#)^*a^*a(aa^\#)^*y_0 = a^+a(aa^\#)^*y_0 = (aa^\#)^*y_0. \end{aligned}$$

Then $y_0 = pa^+(a^\#)^* + y_0 - (aa^\#)^*y_0 = pa^+(a^\#)^* + v - aa^+v$. Hence the general solution of Eq.(7.5) is given by formula (7.6). \square

Theorem 7.5. Let $a \in R^\# \cap R^+$. Then $a \in R^{SEP}$ if and only if Eq.(7.1) and Eq.(7.5) have the same solution.

Proof. (\Rightarrow) If $a \in R^{SEP}$. Then by Theorem 7.3, one has the general solution of Eq.(7.1) is given by formula (7.4). Since $a \in R^{EP}$, then $a^+a^+a = a^+$. Hence formula (7.6) is consistent with formula (7.4). By Theorem 7.4, one obtains the general solution of Eq.(7.5) is given by formula (7.4). Hence Eq.(7.1) and Eq.(7.5) have the same solution.

(\Leftarrow) If Eq.(7.1) and Eq.(7.5) have the same solution. By Theorem 7.1, one gets the general solution of Eq.(7.5) is given by formula (7.2). Then

$$(aa^+p + u - ua^+a)a^+(a^\#)^* = a^*a(aa^\#)^*(pa^+(a^\#)^* + v - aa^+v),$$

e.g.,

$$aa^+pa^+(a^\#)^* = a^*a(aa^\#)^*pa^+(a^\#)^* \text{ for all } p \in R.$$

Choose $p = a$, then $aa^+(a^\#)^* = a^*a(aa^\#)^*$. Multiplying the equality on the right by a^*a^+ , one has $aa^+a^+ = a^*$. Again multiply the last equality on the left by a^+ , one gets

$$a^+a^+ = a^+a^* = a^+a^+aa^*.$$

By [2, Lemma 2.10], we have $a^+ = a^+aa^* = a^*$. Hence $a \in R^{PI}$. By Theorem 7.4, one obtains the general solution of Eq.(7.1) is given by formula (7.6). Then

$$(a^*ap + u - ua^+a)a^+(a^\#)^* = aa^+(pa^+(a^\#)^* + v - aa^+v),$$

e.g.,

$$a^*apa^+(a^\#)^* = aa^+pa^+(a^\#)^* \text{ for } p \in R \text{ with } a^+a^+ap = a^+p.$$

Choose $p = a^+$, one yields $a^*a^+(a^\#)^* = aa^+a^+(a^\#)^*$. Noting that $a^+ = a^*$, then we have $a^* = aa^*a^*$, so $a = a^2a^*$. Hence $a \in R^{SEP}$ by [11, Theorem 1.5.3].

\square

8. The solution of bivariate equations in a fixed set

Lemma 8.1. Let $a \in R^\# \cap R^+$, $y \in R$, $x \in \rho_a$. If $x(a^\#)^*y = 0$, then $(a^\#)^*y = 0$.

Proof. Noting that $x^\#x = \begin{cases} aa^\#, x \in \tau_a =: \{a, a^\#, (a^+)^*\} \\ (aa^\#)^*, x \in \theta_a =: \{a^+, a^*, (a^\#)^*, (a^+)^{\#}, (a^\#)^+\} \end{cases}$. Then we have

(1) if $x \in \tau_a$, then $(a^\#)^*y = a^+a(a^\#)^*y = a^+aaa^\#(a^\#)^*y = a^+ax^\#x(a^\#)^*y = 0$.

(2) if $x \in \theta_a$, then $(a^\#)^*y = (aa^\#)^*(a^\#)^*y = x^\#x(a^\#)^*y = 0$.

Thus, in any case, we have $(a^\#)^*y = 0$. \square

Since $a \in R^{SEP}$, $aa^+(a^\#)^* = a = a(aa^\#)^*a^+a$. Hence we can give the following equation.

$$xy(a^\#)^* = x(aa^\#)^*ya. \tag{8.1}$$

Theorem 8.2. Let $a \in R^\# \cap R^+$. Then $a \in R^{SEP}$ if and only if Eq.(8.1) has at least one solution in $\rho_a^2 \triangleq \{(x, y) | x, y \in \rho_a\}$.

Proof. (\Rightarrow) Clearly $(x, y) = (a, a^+)$ is a solution.

(\Leftarrow) (1) If $y = a$, then we have $xa(a^\#)^* = x(aa^\#)^*a^2$.

① If $x = a$, then $a^2(a^\#)^* = a(aa^\#)^*a^2$. Multiplying the equality on the right by a^+a , one yields $a^2(a^\#)^* = a^2(a^\#)^*a^+a$. Again multiply the last equality on the left by $a^+a^+a^\#$, one gets

$$a^+ = a^+a^+a.$$

Hence $a \in R^{SEP}$, it follows that $a^2(a^\#)^* = a(aa^\#)^*a^2 = a(aa^+)^*a^2 = a^3$. Thus $a^2 = a^\#a^3 = a^\#a^2(a^\#)^* = a(a^\#)^*$. Hence $a \in R^{SEP}$;

② If $x = a^\#$, then $a^\#a(a^\#)^* = a^\#(aa^\#)^*a^2$. Multiplying the equality on the left by a^2 , one has $a^2(a^\#)^* = a(aa^\#)^*a^2$. Hence $a \in R^{SEP}$ by ①;

③ If $x = a^+$, then $a^+a(a^\#)^* = a^+(aa^\#)^*a^2$, e.g., $(a^\#)^* = a^+a^2$. This gives $a(a^\#)^* = a^2$. Hence $a \in R^{SEP}$;

④ If $x = a^*$, then $a^*a(a^\#)^* = a^*(aa^\#)^*a^2 = a^*a^2$. Multiplying the equality on the left by $(a^+)^*$, one obtains $a(a^\#)^* = a^2$. Hence $a \in R^{SEP}$;

⑤ If $x = (a^+)^*$, then $(a^+)^*a(a^\#)^* = (a^+)^*(aa^\#)^*a^2$. Multiplying the equality on the left by aa^* , one has $a^2(a^\#)^* = a(aa^\#)^*a^2$. Hence $a \in R^{SEP}$ by ①;

⑥ If $x = (a^\#)^*$, then $(a^\#)^*a(a^\#)^* = (a^\#)^*(aa^\#)^*a^2$. Multiplying the equality on the left by a^*a^* , one gets $a^*a(a^\#)^* = a^*(aa^\#)^*a^2$. Hence $a \in R^{SEP}$ by ④;

⑦ If $x = (a^+)^{\#} = (aa^\#)^*a(a^\#)^*$, then $(aa^\#)^*a(a^\#)^*a(a^\#)^* = (aa^\#)^*a(a^\#)^*(aa^\#)^*a^2$. Multiplying the equality on the left by a^+a^+ , one obtains $a^+a(a^\#)^* = a^+(aa^\#)^*a^2$. Hence $a \in R^{SEP}$ by ④;

⑧ If $x = (a^\#)^+ = a^+a^3a^+$, then $a^+a^3a^+a(a^\#)^* = a^+a^3a^+(aa^\#)^*a^2$, e.g., $a^+a^3(a^\#)^* = a^+a^4$. This induces $a(a^\#)^* = a^\#a^+a^3(a^\#)^* = a^\#a^+a^4 = a^2$. Hence $a \in R^{SEP}$;

(2) If $y = a^\#$, then we have $xa^\#(a^\#)^* = x(aa^\#)^*a^\#a$.

⑨ If $x = a$, then $aa^\#(a^\#)^* = a(aa^\#)^*a^\#a$. Multiplying the equality on the right by aa^+ , one obtains

$$a(aa^\#)^*a^\#a = a(aa^\#)^*aa^+ = a(aa^\#)^*.$$

Again multiply the last equality on the left by aa^+a^+ , one has $aa^\# = aa^+$. Hence $a \in R^{EP}$, it follows that $(a^\#)^* = a^+a(a^\#)^* = aa^\#(a^\#)^* = a(aa^\#)^*a^\#a = aaa^+a^\#a = a$. Hence $a \in R^{SEP}$;

⑩ If $x = a^\#$, then $a^\#a^\#(a^\#)^* = a^\#(aa^\#)^*a^\#a$. Multiplying the equality on the left by a^2 , one has $aa^\#(a^\#)^* = a(aa^\#)^*a^\#a$. Hence $a \in R^{SEP}$ by ⑨;

⑪ If $x = a^+$, then $a^+a^\#(a^\#)^* = a^+(aa^\#)^*a^\#a = a^+a^\#a$. Multiplying the equality on the left by a^3 , one gets $a(a^\#)^* = a^2$. Hence $a \in R^{SEP}$;

⑫ If $x = a^*$, then $a^*a^\#(a^\#)^* = a^*(aa^\#)^*a^\#a = a^*a^\#a$. Multiplying the equality on the left by $a^2(a^+)^*$, one has $a(a^\#)^* = a^2$. Hence $a \in R^{SEP}$;

⑬ If $x = (a^+)^*$, then $(a^+)^*a^\#(a^\#)^* = (a^+)^*(aa^\#)^*a^\#a$. Multiplying the equality on the left by aa^* , one obtains $aa^\#(a^\#)^* = a(aa^\#)^*a^\#a$. Hence $a \in R^{SEP}$ by ⑨;

⑭ If $x = (a^\#)^*$, then $(a^\#)^*a^\#(a^\#)^* = (a^\#)^*(aa^\#)^*a^\#a$. Multiplying the equality on the left by a^*a^* , one gets $a^*a^\#(a^\#)^* = a^*(aa^\#)^*a^\#a = a^*a^\#a$. Hence $a \in R^{SEP}$ by ⑫;

⑮ If $x = (a^+)^{\#}$, then $(aa^\#)^*a(a^\#)^*a^\#(a^\#)^* = (aa^\#)^*a(a^\#)^*(aa^\#)^*a^\#a$. Multiplying the equality on the left by a^+a^+ , one has $a^+a^\#(a^\#)^* = a^+a^\#a$. Hence $a \in R^{SEP}$ by ⑫;

⑯ If $x = (a^\#)^+$, then $a^+a^3a^+a^\#(a^\#)^* = a^+a^3a^+(aa^\#)^*a^\#a$, e.g., $(a^\#)^* = a^+a^2$. This gives $a(a^\#)^* = a^2$. Hence $a \in R^{SEP}$;

(3) If $y = a^+$, then we have $xa^+(a^\#)^* = x(aa^\#)^*a^+a$, that is, $xa^+(a^\#)^* = xa^+a$.

⑰ If $x = a$, then $aa^+(a^\#)^* = aa^+a = a$. Hence $a \in R^{SEP}$ by Lemma 3.1;

⑱ If $x = a^\#$, then $a^\#a^+(a^\#)^* = a^\#a^+a = a^\#$. Multiplying the equality on the left by a^2 , one gets $aa^+(a^\#)^* = a$. Hence $a \in R^{SEP}$ by Lemma 3.1;

⑲ If $x = a^+$, then $a^+a^+(a^\#)^* = a^+a^+a$. By [17, Lemma 2.10], we get $a^+(a^\#)^* = a^+a$. Multiplying the equality on the left by a , one obtains $aa^+(a^\#)^* = a$. Hence $a \in R^{SEP}$ by Lemma 3.1;

⑳ If $x = a^*$, then $a^*a^+(a^\#)^* = a^*a^+a$. Multiplying the equality on the left by $(a^\#)^*$, one has $a^+(a^\#)^* = a^+a$. Hence $a \in R^{SEP}$ by ⑱;

㉑ If $x = (a^+)^*$, then $(a^+)^*a^+(a^\#)^* = (a^+)^*a^+a = (a^+)^*$. Multiplying the equality on the left by aa^* , one gets $aa^+(a^\#)^* = a$. Hence $a \in R^{SEP}$ by Lemma 3.1;

⑳ If $x = (a^\#)^*$, then $(a^\#)^*a^+(a^\#)^* = (a^\#)^*a^+a$. Multiplying the equality on the left by aa^* , one obtains $aa^+(a^\#)^* = a$. Hence $a \in R^{SEP}$ by Lemma 3.1;

㉑ If $x = (a^+)^{\#}$, then $(aa^\#)^*a(aa^\#)^*a^+(a^\#)^* = (aa^\#)^*a(aa^\#)^*a^+a$. Multiplying the equality on the left by a^*a^+ , one gets $a^*a^+(a^\#)^* = a^*a^+a$. Hence $a \in R^{SEP}$ by ㉐;

㉒ If $x = (a^\#)^+$, then $a^+a^3a^+(a^\#)^* = a^+a^3a^+a$. Multiplying the equality on the left by $a^+a^\#a$, one has $a^+a^+(a^\#)^* = a^+a^+a$. Hence $a \in R^{SEP}$ by ㉑;

(4) If $y = a^*$, then we have $xa^*(a^\#)^* = x(aa^\#)^*a^*a$, e.g., $xa^*(a^\#)^* = xa^*a$. Multiplying the equation on the right by a^+ , one obtains $xa^+ = xa^*$. Hence $a \in R^{PI}$ by [18, Lemma 2.2]. It follows that $y = a^* = a^+$. Hence $a \in R^{SEP}$ by (3).

(5) If $y = (a^+)^*$, then we have $x(a^+)^*(a^\#)^* = x(aa^\#)^*(a^+)^*a$.

㉓ If $x = a$, then $a(a^+)^*(a^\#)^* = a(aa^\#)^*(a^+)^*a$. Multiplying the equality on the right by a^+a , one yields

$$a(a^+)^*(a^\#)^* = a(a^+)^*(a^\#)^*a^+a.$$

Again multiply the last equality on the left by $a^*a^*a^\#$, one has

$$a^*(a^\#)^* = a^+a.$$

Applying the involution to the last equality, we get $a^\#a = a^+a$. Hence $a \in R^{EP}$, it follows that $a(a^+)^*(a^\#)^* = a(a^+)^*a$. Taking involution of the above equality, we obtain

$$a^\#a^+a^* = a^*a^+a^*.$$

Multiplying the equality on the right by $(a^\#)^*a$, one gets $a^*a^+a = a^\# = a^+$. Hence $a \in R^{SEP}$ by [11, Theorem 1.5.3];

㉔ If $x = a^\#$, then $a^\#(a^+)^*(a^\#)^* = a^\#(aa^\#)^*(a^+)^*a$. Multiplying the equality on the left by a^2 , one has $a(a^+)^*(a^\#)^* = a(aa^\#)^*(a^+)^*a$. Hence $a \in R^{SEP}$ by ㉓;

㉕ If $x = a^+$, then $a^+(a^+)^*(a^\#)^* = a^+(aa^\#)^*(a^+)^*a$. Multiplying the equality on the right by a^+a , one yields

$$a^+(a^+)^*(a^\#)^* = a^+(a^+)^*(a^\#)^*a^+a.$$

Again multiply the last equality on the left by a , one has

$$(a^+)^*(a^\#)^* = (a^+)^*(a^\#)^*a^+a.$$

Applying the involution to the last equality, we get $a^\#a^+ = a^+aa^\#a^+$. Multiplying the equality on the right by a^2 , one gets $a^\#a = a^+a$. Hence $a \in R^{EP}$. It follows that $x = a^+ = a^\#$. Hence $a \in R^{SEP}$ by ㉔;

㉖ If $x = a^*$, then $a^*(a^+)^*(a^\#)^* = a^*(aa^\#)^*(a^+)^*a$, e.g., $(a^\#)^* = a^+aa$. Multiplying the equality on the left by a , one yields $a(a^\#)^* = a^2$. Hence $a \in R^{SEP}$;

㉗ If $x = (a^+)^*$, then $(a^+)^*(a^+)^*(a^\#)^* = (a^+)^*(aa^\#)^*(a^+)^*a$. Multiplying the equality on the left by aa^* , one has $a(a^+)^*(a^\#)^* = a(aa^\#)^*(a^+)^*a$. Hence $a \in R^{SEP}$ by ㉓;

㉘ If $x = (a^\#)^*$, then $(a^\#)^*(a^+)^*(a^\#)^* = (a^\#)^*(aa^\#)^*(a^+)^*a$. Multiplying the equality on the left by a^+a^* , one obtains $a^+(a^+)^*(a^\#)^* = a^+(aa^\#)^*(a^+)^*a$. Hence $a \in R^{SEP}$ by ㉕;

㉙ If $x = (a^+)^{\#}$, then $(aa^\#)^*a(aa^\#)^*(a^+)^*(a^\#)^* = (aa^\#)^*a(aa^\#)^*(aa^\#)^*(a^+)^*a$. Multiplying the equality on the left by a^*a^+ , one gets $(a^\#)^* = a^+aa$. Hence $a \in R^{SEP}$ by ㉖;

㉚ If $x = (a^\#)^+$, then $a^+a^3a^+(a^+)^*(a^\#)^* = a^+a^3a^+(aa^\#)^*(a^+)^*a$, e.g.,

$$a^+a^2(a^+)^*(a^\#)^* = a^+a^2(a^+)^*a.$$

Multiplying the equality on the left by $a^+a^\#a$, one has $a^+(a^+)^*(a^\#)^* = a^+(aa^\#)^*(a^+)^*a$. Hence $a \in R^{SEP}$ by ㉕;

(6) If $y = (a^\#)^*$, then we have $x(a^\#)^*(a^\#)^* = x(aa^\#)^*(a^\#)^*a$. Multiplying the equality on the right by $1 - aa^+$, one yields

$$x(a^\#)^*a(1 - aa^+) = 0.$$

By Lemma 8.1, we get

$$(a^\#)^*a(1 - aa^+) = 0.$$

Again multiply the last equality on the left by aa^+a^* , one has $a = a^2a^+$. Hence $a \in R^{EP}$. It follows that $y = (a^\#)^* = (a^+)^*$. Hence $a \in R^{SEP}$ by (5).

(7) If $y = (a^+)^\#$, then we have $x(a^+)^\#(a^\#)^* = x(aa^\#)^*(a^+)^\#a$, e.g.,

$$x(aa^\#)^*a(aa^\#)^*(a^\#)^* = x(aa^\#)^*(aa^\#)^*a(aa^\#)^*a.$$

Multiplying the equality on the right by $(1 - a^+a)$, one obtains

$$x(aa^\#)^*a(aa^\#)^*(1 - a^+a) = 0.$$

By Lemma 8.1, one has $(aa^\#)^*a(aa^\#)^*(1 - a^+a) = 0$. Again multiply the last equality on the left by $a^+a^*a^+$, one yields $a^+ = a^+a^+a$. Hence $a \in R^{EP}$. This infers $y = (a^+)^\# = (a^\#)^\# = a$. Hence $a \in R^{SEP}$ by (1).

(8) If $y = (a^\#)^+$, then we have $xa^+a^3a^+(a^\#)^* = x(aa^\#)^*a^+a^3a^+a$, that is

$$xa^+a^3a^+(a^\#)^* = xa^+a^3.$$

Multiplying the equality on the right by $1 - aa^+$, one obtains

$$xa^+a^3(1 - aa^+) = 0.$$

Noting that $a^+ = (a^\#)^*a^*a^+$. By Lemma 8.1, one gets

$$a^+a^3(1 - aa^+) = 0.$$

Again multiply the last equality on the left by $a^\#$, one has $a = a^2a^+$. Hence $a \in R^{EP}$. This infers $y = (a^\#)^+ = (a^+)^+ = a$. Hence $a \in R^{SEP}$ by (1). \square

It is well known that $a \in R^{SEP}$ if and only if $a^* \in R^{SEP}$. Hence instead a in Eq.(8.1) by a^* , one obtains the following equation and theorem.

$$xya^\# = xaa^\#ya^*. \tag{8.2}$$

Theorem 8.3. Let $a \in R^\# \cap R^+$. Then $a \in R^{SEP}$ if and only if Eq.(8.2) has at least one solution in ρ_a^2 .

Now we establish the following equation.

$$xya^+ + a^\# = xaa^\#ya^* + a^+. \tag{8.3}$$

Theorem 8.4. Let $a \in R^\# \cap R^+$. Then $a \in R^{SEP}$ if and only if Eq.(8.3) has at least one solution in ρ_a^2 .

Proof. (\Rightarrow) Assume that $a \in R^{SEP}$. Then $a^\# = a^+$, this infers Eq.(8.1) has the same solution as Eq.(8.3). By Theorem 8.1, we are done.

(\Leftarrow) Assume that Eq.(8.3) has at least one solution in ρ_a^2 . Let $(x, y) = (x_0, y_0)$ be any solution of Eq.(8.3). Then we have

$$x_0y_0a^+ + a^\# = x_0aa^\#y_0a^* + a^+.$$

Multiplying the equality on the right by $1 - aa^+$, one gets $a^\#(1 - aa^+) = 0$. This gives $a^\# = a^\#aa^+$. Hence $a \in R^{EP}$ by [11, Theorem 1.2.1]. Thus Eq.(8.1) has the same solution as Eq.(8.3). By Theorem 8.1, $a \in R^{SEP}$. \square

9. The consistency of equations and SEP elements

Consider the following equation

$$ax(a^\#)^* = a. \tag{9.1}$$

Theorem 9.1. *Let $a \in R^\# \cap R^+$. Then Eq.(9.1) is consistent if and only if $a \in R^{EP}$. In this case, the general solution of Eq.(9.1) is given by*

$$x = a^* + u - a^+ aua^+ a, \text{ where } u \in R. \tag{9.2}$$

Proof. (\Rightarrow) Assume that Eq.(9.1) is consistent. Then $a = a^2 a^+$, it follows that $a \in R^{EP}$.

(\Leftarrow) If $a \in R^{EP}$, then $aa^*(a^\#)^* = a(aa^\#)^* = aaa^\# = a$ by [1, Theorem 1.3.1]. Hence $x = a^*$ is a solution, so Eq.(9.1) is consistent. Now if Eq.(9.1) is consistent, then $x = a^*$ is a solution and $a \in R^{EP}$. Hence the formula (9.2) is the solution of Eq.(9.1). Let $x = x_0$ be any solution of Eq.(9.1). Then $ax_0(a^\#)^* = a$. Clearly

$$a^+ ax_0aa^+ = a^+ (ax_0(a^\#)^*)a^+ a^+ a = a^+ aa^* a^+ a = a^* a^+ a = a^*.$$

Hence $x_0 = a^* + x_0 - a^+ ax_0aa^+ a$. Thus the general solution of Eq.(9.1) is given by formula (9.2). \square

Theorem 9.2. *Let $a \in R^\# \cap R^+$. Then $a \in R^{SEP}$ if and only if Eq.(9.1) is consistent and the general solution is given by*

$$x = a^+ + u - a^+ aua^+ a, \text{ where } u \in R. \tag{9.3}$$

Proof. (\Rightarrow) If $a \in R^{SEP}$, then $a \in R^{EP}$ and $a^+ = a^*$. It follows that the formula (9.2) is equal to the formula (9.3). By Theorem 9.1, we get Eq.(9.1) is consistent and the general solution is given by the formula (9.3).

(\Leftarrow) From the assumption, we get $a(a^+ + u - a^+ aua^+ a)(a^\#)^* = a$, e.g.,

$$aa^+(a^\#)^* + au(a^\#)^* - aua^+ a(a^\#)^* = a \text{ for all } a \in R.$$

This gives $aa^+(a^\#)^* = a$. Hence $a \in R^{SEP}$ by Lemma 3.1. \square

It is easy to show that the general solution of the following equation is given by (9.3).

$$a(aa^\#)^* axa^+ a = a. \tag{9.4}$$

Hence we have the following corollary.

Corollary 9.3. *Let $a \in R^\# \cap R^+$. Then $a \in R^{SEP}$ if and only if Eq.(9.1) has the same solution as Eq.(9.4).*

Now we change Eq.(9.4) as follows.

$$a(aa^\#)^* axa^+ a = (a^\#)^*. \tag{9.5}$$

Lemma 9.4. *Let $a \in R^\# \cap R^+$. Then Eq.(9.5) is consistent if and only if $a \in R^{EP}$.*

In this case, the general solution of Eq.(9.5) is given by

$$x = a^+ a^+(a^\#)^* + u - a^+ aua^+ a, \text{ where } u \in R. \tag{9.6}$$

Theorem 9.5. *Let $a \in R^\# \cap R^+$. Then $a \in R^{SEP}$ if and only if Eq.(9.5) is consistent and the general solution of Eq.(9.5) is given by $x = a^+ + u - a^+ aua^+ a$, where $u \in R$.*

Proof. (\Rightarrow) If $a \in R^{SEP}$, then $a^\# = a^+ = a^*$, this gives $a^+ a^+(a^\#)^* = a^+$. By Lemma 9.4, we get Eq.(9.5) is consistent and the general solution of Eq.(9.5) is given by $x = a^+ + u - a^+ aua^+ a$, where $u \in R$.

(\Leftarrow) From the assumption, we get $a(aa^\#)^* a(a^+ + u - a^+ aua^+ a)a^+ a = (a^\#)^*$, this gives $a = (a^\#)^*$. Hence $a \in R^{SEP}$ by [11, Theorem 1.5.3]. \square

10. Constructions of group invertible elements and Moore Penrose invertible elements

We need the following lemma which proof is routine.

Lemma 10.1. *Let $a \in R^\# \cap R^+$. Then*

- (1) $x(aa^\#)^*x^+ = aa^+$, where $x \in \tau_a$;
- (2) $x(aa^\#)^*x^+ = a^+a$, where $x \in \theta_a$.

Noting that $a \in R^{EP}$ if and only if $a \in R^+$ and $aa^+ = a^+a$ and $a \in R^{SEP}$ if and only if $a \in R^\# \cap R^+$ and $aa^+ = a^*a$ or $a^+a = aa^*$. Hence the following theorem is an immediate corollary of Lemma 10.1.

Theorem 10.2. *Let $a \in R^\# \cap R^+$. Then the following conditions are equivalent:*

- (1) $a \in R^{EP}$;
- (2) $x(aa^\#)^*x^+ = a^+a$ for some $x \in \tau_a$;
- (3) $x(aa^\#)^*x^+ = aa^+$ for some $x \in \theta_a$;
- (4) $x(aa^\#)^*x^+ = a^*a$ for some $x \in \tau_a$;
- (5) $x(aa^\#)^*x^+ = aa^*$ for some $x \in \theta_a$.

Theorem 10.3. *Let $a \in R^\# \cap R^+$. Then*

- (1) $(xa^+(a^\#)^*)^\# = \begin{cases} aa^+a^*a(aa^\#)^*x^+, x \in \tau_a; \\ a^*ax^\#, x \in \theta_a; \end{cases}$
- (2) $((a^\#)^+)^\# = (aa^\#)^*a^\#(aa^\#)^*$;
- (3) $(xa^+(a^\#)^*)(xa^+(a^\#)^*)^\# = \begin{cases} aa^+, x \in \tau_a; \\ (aa^\#)^*, x \in \theta_a; \end{cases}$
- (4) $a \in R^{SEP}$ if and only if $(xa^+(a^\#)^*)(xa^+(a^\#)^*)^\# = a^*a$ for some $x \in \tau_a$;
- (5) $a \in R^{SEP}$ if and only if $(xa^+(a^\#)^*)(xa^+(a^\#)^*)^\# = aa^*$ for some $x \in \theta_a$.

Proof. (1) If $x \in \tau_a$, then we have $aa^+x = x$, this gives

$$\begin{aligned} (xa^+(a^\#)^*)(aa^+a^*a(aa^\#)^*x^+) &= x(aa^\#)^*x^+ = aa^+; \\ (aa^+a^*a(aa^\#)^*x^+)(xa^+(a^\#)^*) &= aa^+a^*a(aa^\#)^*a^+aa^+(a^\#)^* = aa^+; \\ (xa^+(a^\#)^*)(aa^+a^*a(aa^\#)^*x^+)(xa^+(a^\#)^*) &= aa^+xa^+(a^\#)^* = xa^+(a^\#)^*; \end{aligned}$$

and

$$(aa^+a^*a(aa^\#)^*x^+)(xa^+(a^\#)^*)(aa^+a^*a(aa^\#)^*x^+) = aa^+(aa^+a^*a(aa^\#)^*x^+) = aa^+a^*a(aa^\#)^*x^+.$$

Hence $(xa^+(a^\#)^*)^\# = aa^+a^*a(aa^\#)^*x^+$;

If $x \in \theta_a$, then we have $(aa^\#)^*x = x$, this gives

$$\begin{aligned} (xa^+(a^\#)^*)(a^*ax^\#) &= xa^+ax^\# = (aa^\#)^*; \\ (a^*ax^\#)(xa^+(a^\#)^*) &= a^*a(aa^\#)^*a^+(a^\#)^* = (aa^\#)^*; \\ (xa^+(a^\#)^*)(a^*ax^\#)(xa^+(a^\#)^*) &= (aa^\#)^*xa^+(a^\#)^* = xa^+(a^\#)^*; \end{aligned}$$

and

$$(a^*ax^\#)(xa^+(a^\#)^*)(a^*ax^\#) = (aa^\#)^*(a^*ax^\#) = a^*ax^\#.$$

Hence $(xa^+(a^\#)^*)^\# = a^*ax^\#$;

(2) Noting that $(a^\#)^+ = a^+a^3a^+$. Then

$$\begin{aligned} (a^\#)^+((aa^\#)^*a^\#(aa^\#)^*) &= (a^+a^3a^+)((aa^\#)^*a^\#(aa^\#)^*) = (aa^\#)^*; \\ ((aa^\#)^*a^\#(aa^\#)^*)(a^\#)^+ &= ((aa^\#)^*a^\#(aa^\#)^*)(a^+a^3a^+) = (aa^\#)^*; \\ (a^\#)^+((aa^\#)^*a^\#(aa^\#)^*)(a^\#)^+ &= (aa^\#)^*(a^+a^3a^+) = a^+a^3a^+; \end{aligned}$$

and

$$((aa^\#)^* a^\# (aa^\#)^*) (a^\#)^+ ((aa^\#)^* a^\# (aa^\#)^*) = (aa^\#)^* ((aa^\#)^* a^\# (aa^\#)^*) = (aa^\#)^* a^\# (aa^\#)^*.$$

Hence $((a^\#)^+)^{\#} = (aa^\#)^* a^\# (aa^\#)^*$;

(3) It is an immediate result of (1).

Noting that $a \in R^{SEP}$ if and only if $aa^+ = a^*a$ if and only if $(aa^\#)^* = aa^*$. Hence (4) and (5) follows from (3). \square

Theorem 10.4. Let $a \in R^\# \cap R^+$. Then the following conditions are equivalent:

- (1) $a \in R^{SEP}$;
- (2) $(xa^+(a^\#)^*)(xa^+(a^\#)^*)^+ = a^*a$ for some $x \in \tau_a$;
- (3) $(xa^+(a^\#)^*)(xa^+(a^\#)^*)^+ = aa^*$ for some $x \in \theta_a$.

Proof. First, we have $(xa^+(a^\#)^*)^+ = aa^+a^*(aa^\#)^*x^+$. Hence

$$(xa^+(a^\#)^*)(xa^+(a^\#)^*)^+ = x(aa^\#)^*x^+.$$

Next, using Theorem 10.2, we can complete the proof. \square

Similar to Theorem 10.3, we can show the following theorem.

Theorem 10.5. Let $a \in R^\# \cap R^+$. Then

- (1) $(aa^+x)^+ = x^+(aa^\#)^*$, where $x \in \rho_a$;
- (2) $(aa^+x)^\# = \begin{cases} x^\#aa^\# & ,x \in \tau_a \\ x^+(aa^\#)^* & ,x \in \theta_a \end{cases}$.

The following theorem follows from Theorem 6.1, Theorem 10.3, Theorem 10.4 and Theorem 10.5.

Theorem 10.6. Let $a \in R^\# \cap R^+$. Then the following conditions are equivalent:

- (1) $a \in R^{SEP}$;
- (2) $aa^+a^*(aa^\#)^*x^+ = x^+(aa^\#)^*$ for some $x \in \rho_a$;
- (3) $aa^+a^*(aa^\#)^*x^+ = x^\#aa^\#$ for some $x \in \tau_a$;
- (4) $a^*ax^\# = x^+(aa^\#)^*$ for some $x \in \theta_a$.

References

- [1] W. X. Chen. On EP elements, normal elements and parital isometries in rings with involution. Electron. J. Linear Algebra. 23(2012) 553-561.(DOI: <https://doi.org/10.13001/1081-3810.1540>)
- [2] D. Drivaliaris, S. Karanasios, D. Pappas. Factorizations of EP operators. Linear Algebra Appl. 429(2008) 1555-1567. (DOI: 10.1016/j.laa.2008.04.026)
- [3] R. E. Hartwig. Block generalized inverses. Arch. Retion. Mech. Anal. 61(1976) 197-251. (DOI: 10.1007/BF00281485)
- [4] R. E. Hartwig. Generalized inverses, EP elements and associates. Rev. Roumaine Math. Pures Appl. 23(1978) 57-60.
- [5] R. E. Hartwig, I. J. Katz. Products of EP elements in reflexive semigroups. Linear Algebra Appl. 14(1976) 11-19. (DOI: 10.1016/0024-3795(76)90058-6)
- [6] R. E. Harte, M. Mbekhta. On generalized inverses in C^* -algebras. Studia Math. 103(1992) 71-77.
- [7] S. Karanasios. EP elements in rings and semigroup with involution and C^* -algebras. Serdica Math. J. 41(2015) 83-116.
- [8] J. J. Koliha. The Drazin and Moore-Penrose inverse in C^* -algebras. Math. Proc. R. Ir. Acad. 99A(1999) 17-27.
- [9] J. J. Koliha, D. Cvetković, D. S. Djordjević. Moore-Penrose inverse in rings with involution. Linear Algebra Appl. 426(2007) 371-381. (DOI: 10.1016/j.laa.2007.05.012)
- [10] J. J. Koliha, P. Patrício. Elements of rings with equal spectral idempotents. J. Aust. Math. Soc. 72(2002) 137-152. (DOI: 10.1017/S1446788700003657)
- [11] D. Mosić, Generalized inverses, Faculty of Sciences and Mathematics, University of Niš, Niš, 2018.
- [12] D. Mosić, D. S. Djordjević, J. J. Koliha. EP elements in rings. Linear Algebra Appl. 431(2009) 527-535. (DOI: 10.1016/j.laa.2009.02.032)
- [13] D. Mosić, D. S. Djordjević. New characterizations of EP , generalized normal and generalized Hermitian elements in rings. Appl. Math. Comput. 218(2012) 6702-6710. (DOI: 10.1016/j.amc.2011.12.030)
- [14] D. Mosić, D. S. Djordjević. Further results on partial isometries and EP elements in rings with involution. Math. Comput. Model. 54(2011) 460-465. (DOI: 10.1016/j.mcm.2011.02.035)

- [15] D. Mosić, D. S. Djordjević. Partial isometries and *EP* elements in rings with involution. *Electron. J. Linear Algebra*. 18(2009) 761-722.
- [16] Z. C. Xu, R. J. Chen, J. C. Wei, Strongly *EP* elements in a ring with involution. *Filomat*, 34(6)(2020): 2101-2107.
- [17] D. D. Zhao, J. C. Wei, Strongly *EP* elements in rings with involution. *J. Algebra Appl.*, (2022), 2250088, 10pages, DOI: 10.1142/S0219498822500888.
- [18] D. D. Zhao, J. C. Wei, Some new characterizations of partial isometries in rings with involution. *Intern. Eletron. J. Algebra*, 30(2021): 304-311.
- [19] R. J. Zhao, H. Yao, J. C. Wei, Characterizations of partial isometries and two special kinds of *EP* elements. *Czecho. Math. J.*, 70(2)(2020): 539-551.
- [20] J. M. Yan, X. Chen, J. C. Wei, Torsion invertible elements over $*$ -ring. *Journal of Yangzhou University. Natural Science Edition*, 25(1)(2022): 1-3, 24.
- [21] X. M. Li, X. Y. Zhu, J. C. Wei, Hermite matrix and solutions of matrix equations. *Journal of Yangzhou University. Natural Science Edition*, 25(5)(2022): 1-6.