



Approximations: Modified Szász-Beta operators via general-Appell polynomials

Nadeem Rao^a, Nand Kishor Jha^{b,*}

^aDepartment of Mathematics, University Center for Research and Development, Chandigarh University, Mohali-140413, Panjab, India

^bDepartment of Mathematics, Chandigarh University, Mohali-140413, Panjab, India

Abstract. We present a new sequence of modified Szász-Beta type sequence of operators via general-Appell Polynomials to investigate the approximation properties of Lebesgue integrable functions ($L_1[0, \infty)$). In addition, we study estimates in view of test functions and central moments. Next, convergence rate is discussed using the Korovkin theorem and Voronovskaja type theorem. Moreover, direct approximation results via modulus of continuity of first and second order, Peetre's K-functional, Lipschitz type space, and the r^{th} order Lipschitz type maximal functions are investigated. In subsequent section, weighted approximation results and statistical approximation theorems are discussed.

1. Introduction

Approximation theory has long been a fundamental area of mathematical analysis, influencing pure mathematics, computational science, and practical applications in engineering and computer graphics. In 1885, Weierstrass [1] established a seminal result in this field, the Weierstrass approximation theorem, which states that any continuous function on a closed interval can be uniformly approximated by polynomials. Later, in 1912, Bernstein [2] provided an elegant proof of this theorem using a sequence of polynomials. The advent of fractional calculus has introduced a transformative perspective to operator theory in approximation. The intersection of fractional calculus and operator theory continues to grow as a research domain, with applications extending beyond pure mathematics into interdisciplinary fields such as quantum mechanics, control theory, signal processing, and materials science.

Szász in 1950 [3], constructed a sequence of operators over the infinite length of interval, i.e., $[0, \infty)$ as:

$$P_{\kappa}(\tilde{g}; u) = \sum_{i=0}^{\infty} \tilde{g}\left(\frac{i}{\kappa}\right) p_{\kappa,i}(u), \quad (1)$$

where $u \in [0, \infty)$, $\tilde{g} \in C[0, \infty)$ and $p_{\kappa,i}(u) = e^{-\kappa u} \frac{(\kappa u)^i}{i!}$.

2020 *Mathematics Subject Classification.* Primary 41A10; Secondary 41A25, 41A28, 41A35, 41A36.

Keywords. Modulus of smoothness, Szász operator, Appell Polynomials, Order of approximation; Rate of convergence.

Received: 16 March 2025; Revised: 05 August 2025; Accepted: 08 August 2025

Communicated by Snežana Č. Živković-Zlatanović

* Corresponding author: Nand Kishor Jha

Email addresses: nadeemrao1990@gmail.com (Nadeem Rao), nandkishorjha1982@gmail.com (Nand Kishor Jha)

ORCID iDs: <https://orcid.org/0000-0002-5681-9563> (Nadeem Rao), <https://orcid.org/0009-0008-5297-0486> (Nand Kishor Jha)

In 1962, Schurer [4] constructed a new sequence of Bernstein operators [2] which is denoted as $B_{\kappa+s} : C[0, 1+l] \rightarrow C[0, 1]$ and defined by:

$$B_{\kappa+s}(\tilde{g}; u) = \sum_{j=0}^{\kappa+s} \tilde{g}\left(\frac{j}{\kappa}\right) \binom{\kappa+s}{j} u^j (1-u)^{\kappa+s-j}, \quad u \in [0, 1+l], \quad (2)$$

where $l \in \mathbb{N} \cup \{0\}$ and $\tilde{g} \in C[0, 1+l]$. But these sequences of operators given in (2) are restricted to $C[0, 1+l]$.

The classical modified Szász operators are linear positive operators and approximate continuous functions over the positive semi-axis. Many mathematicians have constructed several generalizations of these sequences of operators to provide flexibility in rate of convergence and order of approximation, e.g., Szász-Schurer and Szász-Integral type operators etc. (see [5], [6]). Several mathematicians, e.g., Raza et al. ([7], [8]), Aslan et al. ([9], [10]), Özger et al. ([11], [12]), Cai et al. ([13], [14]), Braha et al. ([15], [16]), Ayman Mursaleen et al. ([17], [18], [19]), Khursheed et al. ([20], [21]), Mohiuddine et al. ([22], [23]), Mursaleen et al. ([24], [25]), Rao et al. ([26], [27]), Khan et al. [28], Nasiruzzaman [29], Acu et al. ([30], [31]) and Jha et al. [32] provided a number of generalizations for these kinds of sequences to investigate flexibility in approximation properties across several functional spaces.

Recently, Raza et al. [33] provided a class of sequence of operators $G_{\kappa,A}(\cdot; \cdot)$, $\kappa \in \mathbb{N}$, given by the formula

$$\mathcal{G}_{\kappa,A}(\tilde{g}; u) = \frac{e^{-\kappa u}}{\tilde{\Omega}(1)\tilde{\psi}(h, 1)} \sum_{v=0}^{\infty} \frac{A_{p,v}(\kappa u, h)}{v!} \tilde{g}\left(\frac{v}{\kappa}\right), \quad u \in \mathbb{R}_0^+, \quad (3)$$

where $A_{p,v}$ is the two variable Appell polynomials (see [33]).

The sequence presented by (3) are positive and linear. The basic information about positive linear operators, including their modifications and applications can be found in [34].

As the operators described in (3) are limited for continuous function only, we present a sequence of positive linear operators to provide approximations in the larger class of functions, which is termed as modified Szász-Beta operators in context of general Appell Polynomials as:

$$\mathcal{M}_{\kappa+s}^{\mathcal{A}}(\tilde{g}; u) = \sum_{v=0}^{\infty} \mathcal{A}_v^p((\kappa+s)u, h) \int_0^{\infty} Q_{\kappa+s}^v(y) \tilde{g}(y) dy, \quad \text{for } u \in \mathbb{R}_0^+, \quad (4)$$

where

$$\mathcal{A}_v^p((\kappa+s)u, h) = \frac{e^{-(\kappa+s)u}}{\tilde{\Omega}(1)\tilde{\psi}(h, 1)} \frac{A_{p,v}((\kappa+s)u, h)}{v!}$$

$$\text{and } Q_{\kappa+s}^v(y) = \frac{1}{\beta(v+1, \kappa+s)} \left[\frac{y^v}{(1+y)^{v+1+\kappa+s}} \right],$$

with β (Beta) function which is given as:

$$\beta(v+1, \kappa+s) = \int_0^{\infty} \frac{y^v}{(1+y)^{v+1+\kappa+s}} dy.$$

Lemma 1.1. *The sequence of operators introduced in (4) are linear.*

Proof. In view of (4) and for $\tilde{\lambda}_1, \tilde{\lambda}_2 \in \mathbb{R}$, we have

$$\begin{aligned} \mathcal{M}_{\kappa+s}^{\mathcal{A}}(\tilde{\lambda}_1 \tilde{g}_1 + \tilde{\lambda}_2 \tilde{g}_2; u) &= \sum_{v=0}^{\infty} \mathcal{A}_v^p((\kappa+s)u, h) \int_0^{\infty} Q_{\kappa+s}^v(y) (\tilde{\lambda}_1 \tilde{g}_1 + \tilde{\lambda}_2 \tilde{g}_2)(y) dy \\ &= \tilde{\lambda}_1 \sum_{v=0}^{\infty} \mathcal{A}_v^p((\kappa+s)u, h) \int_0^{\infty} Q_{\kappa+s}^v(y) \tilde{g}_1(y) dy \end{aligned}$$

$$\begin{aligned}
& + \tilde{\lambda}_2 \sum_{v=0}^{\infty} \mathcal{A}_v^p((\kappa+s)u, h) \int_0^{\infty} Q_{\kappa+s}^v(y) \tilde{g}_2(y) dy \\
& = \tilde{\lambda}_1 \mathcal{M}_{\kappa+s}^{\mathcal{A}}(\tilde{g}_1; u) + \tilde{\lambda}_2 \mathcal{M}_{\kappa+s}^{\mathcal{A}}(\tilde{g}_2; u),
\end{aligned}$$

where $\tilde{g}_1, \tilde{g}_2 \in L_1[0, \infty)$. $\square$

Lemma 1.2. Let $\tilde{g}_{\theta}(y) = y^{\theta}$, $\theta \in \{0, 1, 2\}$ be test functions. Then, we have

$$\mathcal{M}_{\kappa+s}^{\mathcal{A}}(\tilde{g}_{\theta}; u) = \sum_{v=0}^{\infty} A_v^p((\kappa+s)u, h) \frac{(v+\theta)!(\kappa+s-\theta-1)!}{v!(\kappa+s-1)!}.$$

Proof. We know,

$$\begin{aligned}
\int_0^{\infty} Q_{\kappa+s}^v(y) g_{\theta}(y) dy &= \frac{1}{\beta(v+1, \kappa)} \int_0^{\infty} \frac{y^{v+\theta}}{(1+y)^{v+1+\kappa}} dy \\
&= \frac{1}{\beta(v+1, \kappa+s)} \beta(v+\theta+1, \kappa+s-\theta) \\
&= \frac{(v+\theta)!(\kappa+s-\theta-1)!}{v!(\kappa+s-1)!}.
\end{aligned}$$

From (4), we have

$$\mathcal{M}_{\kappa+s}^{\mathcal{A}}(\tilde{g}_{\theta}; u) = \sum_{v=0}^{\infty} A_v^p((\kappa+s)u, h) \frac{(v+\theta)!(\kappa+s-\theta-1)!}{v!(\kappa+s-1)!}.$$

$\square$

Lemma 1.3. As discussed by Raza et al. in [33], we can have the following equalities:

$$\begin{aligned}
\sum_{v=0}^{\infty} \frac{\mathcal{A}_{p,v}(\kappa u, h)}{v!} &= \tilde{\Omega}(1) e^{\kappa u} \tilde{\psi}(h, 1), \\
\sum_{v=0}^{\infty} v \frac{\mathcal{A}_{p,v}(\kappa u, h)}{v!} &= \left[\kappa u \tilde{\Omega}(1) \tilde{\psi}(h, 1) + \tilde{\Omega}(1) \tilde{\psi}'(h, 1) + \tilde{\Omega}'(1) \tilde{\psi}(h, 1) \right] e^{\kappa u}, \\
\sum_{v=0}^{\infty} v^2 \frac{\mathcal{A}_{p,v}(\kappa u, h)}{v!} &= \left[((\kappa+s)^2 u^2 + \kappa u) \tilde{\Omega}(1) \tilde{\psi}(h, 1) + (2\kappa u + 1) [\tilde{\Omega}(1) \tilde{\psi}'(h, 1) \right. \\
&\quad \left. + \tilde{\Omega}'(1) \tilde{\psi}(h, 1)] + 2\tilde{\psi}'(h, 1) \tilde{\Omega}'(1) + \tilde{\psi}''(h, 1) \tilde{\Omega}(1) \right. \\
&\quad \left. + \tilde{\psi}(h, 1) \tilde{\Omega}''(1) \right] e^{\kappa u}.
\end{aligned}$$

Lemma 1.4. Let $\tilde{g}_{\theta}(y) = y^{\theta}$, $\theta \in \{0, 1, 2\}$ be the test functions by (3). Then, we have

$$\begin{aligned}
\mathcal{M}_{\kappa+s}^{\mathcal{A}}(1; u) &= 1, \\
\mathcal{M}_{\kappa+s}^{\mathcal{A}}(y; u) &= u + \frac{1}{\kappa+s-1} \left[u + \frac{\tilde{\psi}'(h, 1)}{\tilde{\psi}(h, 1)} + \frac{\tilde{\Omega}'(1)}{\tilde{\Omega}(1)} + 1 \right]; \quad \kappa+s > 1, \\
\mathcal{M}_{\kappa+s}^{\mathcal{A}}(y^2; u) &= u^2 + \frac{1}{(\kappa+s-1)(\kappa+s-2)} \left[(3(\kappa+s)-2)u^2 + 4(\kappa+s)u \right. \\
&\quad \left. + (2(\kappa+s)u + 4) \left\{ \frac{\tilde{\psi}'(h, 1)}{\tilde{\psi}(h, 1)} + \frac{\tilde{\Omega}'(1)}{\tilde{\Omega}(1)} \right\} + 2 \frac{\tilde{\psi}'(h, 1) \tilde{\Omega}'(1)}{\tilde{\psi}(h, 1) \tilde{\Omega}(1)} \right]
\end{aligned}$$

$$+ \frac{\tilde{\psi}''(h, 1)}{\tilde{\psi}(h, 1)} + \frac{\tilde{\Omega}''(1)}{\tilde{\Omega}(1)} + 2 \Big]; \quad \kappa + s > 2,$$

for each $u \in \mathbb{R}_0^+$.

Proof. In the direction of (4), we have

$$\mathcal{M}_{\kappa+s}^{\mathcal{A}}(\tilde{g}_\theta; u) = \sum_{v=0}^{\infty} \mathcal{A}_v^p((\kappa+s)u, h) \int_0^{\infty} Q_{\kappa+s}^v(y) \tilde{g}_\theta(y) dy.$$

Now, for $\theta = 0$,

$$\mathcal{M}_{\kappa+s}^{\mathcal{A}}(1; u) = \sum_{v=0}^{\infty} A_v^p((\kappa+s)u, h) \frac{v!(\kappa+s-1)!}{v!(\kappa+s-1)!} = 1.$$

For $\theta = 1$ and $\kappa + s > 1$,

$$\begin{aligned} \mathcal{M}_{\kappa+s}^{\mathcal{A}}(y; u) &= \frac{e^{-(\kappa+s)u}}{\tilde{\Omega}(1)\tilde{\psi}(h, 1)} \sum_{v=0}^{\infty} \frac{\mathcal{A}_{p,v}((\kappa+s)u, h)}{v!} \frac{(v+1)!(\kappa+s-2)!}{v!(\kappa+s-1)!} \\ &= \frac{e^{-(\kappa+s)u}}{\tilde{\Omega}(1)\tilde{\psi}(h, 1)} \sum_{v=0}^{\infty} \frac{\mathcal{A}_{p,v}((\kappa+s)u, h)}{v!} \frac{v+1}{\kappa+s-1} \\ &= \frac{e^{-(\kappa+s)u}}{(\kappa+s-1)\tilde{\Omega}(1)\tilde{\psi}(h, 1)} \left[\tilde{\Omega}(1)\tilde{\psi}'(h, 1) + \tilde{\Omega}'(1)\tilde{\psi}(h, 1) \right. \\ &\quad \left. + ((\kappa+s)u+1)\tilde{\Omega}(1)\tilde{\psi}(h, 1) \right] e^{(\kappa+s)u} \\ &= \frac{1}{\kappa+s-1} \left[(\kappa+s)u+1 + \frac{\tilde{\psi}'(h, 1)}{\tilde{\psi}(h, 1)} + \frac{\tilde{\Omega}'(1)}{\tilde{\Omega}(1)} \right] \\ &= u + \frac{1}{\kappa+s-1} \left[u + \frac{\tilde{\psi}'(h, 1)}{\tilde{\psi}(h, 1)} + \frac{\tilde{\Omega}'(1)}{\tilde{\Omega}(1)} + 1 \right] \\ &:= \psi_{\kappa+s}. \quad (\text{Say}) \end{aligned}$$

For $\theta = 2$

$$\begin{aligned} \mathcal{M}_{\kappa+s}^{\mathcal{A}}(y^2; u) &= \frac{e^{-(\kappa+s)u}}{\tilde{\Omega}(1)\tilde{\psi}(h, 1)} \sum_{v=0}^{\infty} \frac{\mathcal{A}_{p,v}((\kappa+s)u, h)}{v!} \frac{(v+2)!(\kappa+s-3)!}{v!(\kappa+s-1)!} \\ &= \frac{e^{-(\kappa+s)u}}{\tilde{\Omega}(1)\tilde{\psi}(h, 1)} \sum_{v=0}^{\infty} \frac{\mathcal{A}_{p,v}((\kappa+s)u, h)}{v!} \frac{((\kappa+s)u+1)(v+2)}{(\kappa+s-1)(\kappa+s-2)} \\ &= \frac{1}{(\kappa+s-1)(\kappa+s-2)\tilde{\Omega}(1)\tilde{\psi}(h, 1)} \left[(2(\kappa+s)u)\tilde{\Omega}(1)\tilde{\psi}'(h, 1) \right. \\ &\quad \left. + 4(\tilde{\Omega}(1)\tilde{\psi}'(h, 1) + \tilde{\Omega}'(1)\tilde{\psi}(h, 1)) + 2\tilde{\psi}'(h, 1)\tilde{\Omega}'(1) + \tilde{\psi}''(h, 1)\tilde{\Omega}(1) \right. \\ &\quad \left. + \tilde{\psi}(h, 1)\tilde{\Omega}''(1) + ((\kappa+s)^2u^2 + 4(\kappa+s)u + 2)\tilde{\Omega}(1)\tilde{\psi}(h, 1) \right] \\ &= u^2 + \frac{1}{(\kappa+s-1)(\kappa+s-2)} \left[(3(\kappa+s)-2)u^2 + 4(\kappa+s)u \right. \\ &\quad \left. + (2(\kappa+s)u+4) \left[\frac{\tilde{\psi}'(h, 1)}{\tilde{\psi}(h, 1)} + \frac{\tilde{\Omega}'(1)}{\tilde{\Omega}(1)} \right] + 2 \frac{\tilde{\psi}'(h, 1)\tilde{\Omega}'(1)}{\tilde{\Omega}(1)\tilde{\psi}(h, 1)} \right] \end{aligned}$$

$$+ \frac{\tilde{\Omega}''(1)}{\tilde{\Omega}(1)} + \frac{\tilde{\psi}''(h, 1)}{\tilde{\psi}(h, 1)} + 2]; \quad \kappa + s > 2.$$

Hence, proof of Lemma 1.4 is completed. $\square$

Lemma 1.5. Let $\gamma_\theta(y) = (y - u)^\theta$, $\theta = 0, 1, 2$. Then, for the operators (4), we have central moments $\mathcal{M}_{\kappa+s}^{\mathcal{A}}(\gamma_\theta(y), u)$ as:

$$\begin{aligned} \mathcal{M}_{\kappa+s}^{\mathcal{A}}(\gamma_0(y), u) &= 1; \\ \mathcal{M}_{\kappa+s}^{\mathcal{A}}(\gamma_1(y), u) &= \frac{1}{\kappa + s - 1} \left[u + \frac{\tilde{\psi}'(h, 1)}{\tilde{\psi}(h, 1)} + \frac{\tilde{\Omega}'(1)}{\tilde{\Omega}(1)} + 1 \right]; \quad \kappa + s > 1, \\ \mathcal{M}_{\kappa+s}^{\mathcal{A}}(\gamma_2(y), u) &= \frac{1}{(\kappa + s - 1)(\kappa + s - 2)} \left[(\kappa + s + 2)u^2 + 2u(\kappa + s + 2) \right. \\ &\quad + 4(u + 1) \left[\frac{\tilde{\psi}'(h, 1)}{\tilde{\psi}(h, 1)} + \frac{\tilde{\Omega}'(1)}{\tilde{\Omega}(1)} \right] + 2 \frac{\tilde{\psi}'(h, 1)\tilde{\Omega}'(1)}{\tilde{\Omega}(1)\tilde{\psi}(h, 1)} + \frac{\tilde{\Omega}''(1)}{\tilde{\Omega}(1)} \\ &\quad \left. + \frac{\tilde{\psi}''(h, 1)}{\tilde{\psi}(h, 1)} + 2 \right]; \quad \kappa + s > 2, \end{aligned}$$

for each $u \in \mathbb{R}_0^+$.

Proof. With the aid of the Lemma 1.4 and linearity property, we can easily complete the proof of Lemma 1.5. $\square$

In the following sections, we examine the convergence rate of operators and their approximation order. Specifically, we discuss direct results both globally and locally in several spaces. In the final section, we explore some results of the A-Statistical approximation in various functional spaces.

2. Uniform Rate of Convergence and Order of Approximation

Definition 2.1. [34] The modulus of smoothness for $\tilde{g} \in C[0, \infty)$ is given by

$$\omega(\tilde{g}; \delta) = \sup_{|u_1 - u_2| \leq \delta} |\tilde{g}(u_1) - \tilde{g}(u_2)|, \quad u_1, u_2 \in [0, \infty).$$

Theorem 2.2. Let $\mathcal{M}_{\kappa+s}^{\mathcal{A}}(\cdot; \cdot)$ be a operators described in Eq. (4). Then, on each bounded and closed interval of $[0, \infty)$, $\mathcal{M}_{\kappa+s}^{\mathcal{A}}(\tilde{g}; \cdot) \rightrightarrows \tilde{g}$, for all $\tilde{g} \in C_B[0, \infty)$, where the symbol $\rightrightarrows$ denotes uniform convergence.

Proof. Considering the classical Korovkin theorem [36], which characterizes the uniform convergence for the sequence of positive linear operators, it is enough to note that

$$\lim_{\kappa \rightarrow \infty} \mathcal{M}_{\kappa+s}^{\mathcal{A}}(\tilde{g}_\theta; u) = u^\theta, \quad \theta = 0, 1, 2,$$

uniformly on all bounded and closed subsets of $[0, \infty)$. We can easily establish this result with the help of Lemma 1.4. $\square$

Now, we show that Voronovskaja-type asymptotic approximation theorem for the $\mathcal{M}_{\kappa+s}^{\mathcal{A}}(\cdot; \cdot)$ given in (4).

Theorem 2.3. Let $\tilde{g} \in C_B[0, \infty)$ and $\tilde{g}', \tilde{g}''$ exist at a fixed point $u \in [0, \infty)$. Then, one has

$$\begin{aligned} \lim_{\kappa \rightarrow \infty} (\kappa + s) \left(\mathcal{M}_{\kappa+s}^{\mathcal{A}}(\tilde{g}; u) - \tilde{g}(u) \right) &= \tilde{g}'(u) \left[u + \frac{\tilde{\psi}'(h, 1)}{\tilde{\psi}(h, 1)} + \frac{\tilde{\Omega}'(1)}{\tilde{\Omega}(1)} + 1 \right] \\ &\quad + \frac{1}{2} (u^2 + 2u) \tilde{g}''(u). \end{aligned}$$

Proof. In accordance with Taylor's formula for the function $\tilde{g}$, we get

$$\tilde{g}(y) = \tilde{g}(u) + (y - u)\tilde{g}'(u) + \frac{1}{2}(y - u)^2\tilde{g}''(u) + t(y, u)(y - u)^2, \quad (5)$$

where $t(y, u)$ is Peano remainder and

$$\lim_{y \rightarrow u} t(y, u) = 0.$$

Applying operators on both the sides in (5), we yield

$$\begin{aligned} (\mathcal{M}_{\kappa+s}^{\mathcal{A}}(\tilde{g}; u) - \tilde{g}(u)) &= \tilde{g}'(u)\mathcal{M}_{\kappa+s}^{\mathcal{A}}((y - u); u) + \frac{1}{2}\tilde{g}''(u)\mathcal{M}_{\kappa+s}^{\mathcal{A}}((y - u)^2; u) \\ &+ \mathcal{M}_{\kappa+s}^{\mathcal{A}}(t(y, u)(y - u)^2; u). \end{aligned}$$

In view of Lemma 1.5

$$\begin{aligned} (\kappa + s)(\mathcal{M}_{\kappa+s}^{\mathcal{A}}(\tilde{g}; u) - \tilde{g}(u)) &= \frac{(\kappa + s)\tilde{g}'(u)}{\kappa + s - 1} \left[u + \frac{\tilde{\psi}'(h, 1)}{\tilde{\psi}(h, 1)} + \frac{\tilde{\Omega}'(1)}{\tilde{\Omega}(1)} + 1 \right] \\ &+ \frac{1}{2} \frac{(\kappa + s)\tilde{g}''(u)}{(\kappa + s - 1)(\kappa + s - 2)} \left[(\kappa + s + 2)u^2 + 2u(\kappa + s + 2) \right. \\ &+ 4(u + 1) \left(\frac{\tilde{\psi}'(h, 1)}{\tilde{\psi}(h, 1)} + \frac{\tilde{\Omega}'(1)}{\tilde{\Omega}(1)} \right) + 2 \frac{\tilde{\psi}'(h, 1)\tilde{\Omega}'(1)}{\tilde{\Omega}(1)\tilde{\psi}(h, 1)} + \frac{\tilde{\Omega}''(1)}{\tilde{\Omega}(1)} \\ &\left. + \frac{\tilde{\psi}''(h, 1)}{\tilde{\psi}(h, 1)} + 2 \right] + (\kappa + s)\mathcal{M}_{\kappa+s}^{\mathcal{A}}(t(y, u)(y - u)^2; u). \end{aligned}$$

Operate the limits on both the sides of the above expression, we get

$$\begin{aligned} \lim_{\kappa \rightarrow \infty} (\kappa + s)(\mathcal{M}_{\kappa+s}^{\mathcal{A}}(\tilde{g}; u) - \tilde{g}(u)) &= \tilde{g}'(u) \left[u + \frac{\tilde{\psi}'(h, 1)}{\tilde{\psi}(h, 1)} + \frac{\tilde{\Omega}'(1)}{\tilde{\Omega}(1)} + 1 \right] \\ &+ \frac{1}{2}(u^2 + 2u)\tilde{g}''(u) \\ &+ \lim_{\kappa \rightarrow \infty} (\kappa + s)\mathcal{M}_{\kappa+s}^{\mathcal{A}}(t(y, u)(y - u)^2; u). \end{aligned}$$

Now, we need to show that

$$\lim_{\kappa \rightarrow \infty} (\kappa + s)\mathcal{M}_{\kappa+s}^{\mathcal{A}}(t(y, u)(y - u)^2; u) = 0.$$

In view of Cauchy-Schwarz inequality, we calculate last term of above expression can be written as:

$$\begin{aligned} (\kappa + s)\mathcal{M}_{\kappa+s}^{\mathcal{A}}(t(y, u)(y - u)^2; u) &\leq \sqrt{\mathcal{M}_{\kappa+s}^{\mathcal{A}}(t^2(y, u); u)} \\ &\times \sqrt{(\kappa + s)^2\mathcal{M}_{\kappa+s}^{\mathcal{A}}((y - u)^4; u)}. \end{aligned} \quad (6)$$

We see that $t^2(u, u) = 0$ and $t^2(y, u) \in C_B[0, \infty)$. Thus we have

$$\lim_{\kappa \rightarrow \infty} \mathcal{M}_{\kappa+s}^{\mathcal{A}}(t^2(y, u); u) = t^2(u, u) = 0. \quad (7)$$

From (6) and (7) it follows that

$$\lim_{\kappa \rightarrow \infty} (\kappa + s)\mathcal{M}_{\kappa+s}^{\mathcal{A}}(t(y, u)(y - u)^2; u) = 0.$$

Hence, the proof is completed. $\square$

According to Shisha et al. [35], order of convergence relative to Ditzian-Totik modulus of continuity can easily be proved.

Theorem 2.4. Consider $\tilde{g} \in C_B[0, \infty)$ and for the operators $\mathcal{M}_{\kappa+s}^{\mathcal{A}}(., .)$ presented in Eq. (4), we acquire

$$|\mathcal{M}_{\kappa+s}^{\mathcal{A}}(\tilde{g}; u) - \tilde{g}(u)| \leq 2\omega(\tilde{g}; \delta),$$

where $\delta = \sqrt{\mathcal{M}_{\kappa+s}^{\mathcal{A}}((y-u)^2; u)}$.

Proof. In accordance with Lemma 1.4, 1.5 and Cauchy-Schwartz inequality, we have

$$\begin{aligned} |\mathcal{M}_{\kappa+s}^{\mathcal{A}}(\tilde{g}; u) - \tilde{g}(u)| &\leq \mathcal{M}_{\kappa+s}^{\mathcal{A}}(|\tilde{g}(y) - \tilde{g}(u)|; u) \\ &\leq \mathcal{M}_{\kappa+s}^{\mathcal{A}}\left(\left(1 + \frac{|y-u|}{\delta}\right)\omega(\tilde{g}, \delta); u\right) \\ &\leq \left(1 + \frac{1}{\delta} \mathcal{M}_{\kappa+s}^{\mathcal{A}}(|y-u|; u)\right)\omega(\tilde{g}, \delta) \\ &\leq \left(1 + \frac{1}{\delta} \sqrt{\mathcal{M}_{\kappa+s}^{\mathcal{A}}(\gamma_2(y); u)}\right)\omega(\tilde{g}, \delta). \end{aligned}$$

By selecting $\delta = \sqrt{\mathcal{M}_{\kappa+s}^{\mathcal{A}}(\gamma_2(y); u)}$, we obtained the desired proof. $\square$

3. Locally Approximation Results

We recall a few functional spaces and functional relations in this part as: $C_B[0, \infty)$: Denotes a real valued functional space which acquires bounded and continuous functions. Now, Peetre's K-functional [34] is defined as

$$K_2(\tilde{g}, \delta) = \inf_{\tilde{h} \in C_B^2[0, \infty)} \left\{ \|\tilde{g} - \tilde{h}\|_{C_B[0, \infty)} + \delta \|\tilde{h}''\|_{C_B^2[0, \infty)} \right\},$$

where $C_B^2[0, \infty) = \{\tilde{h} \in C_B[0, \infty) : \tilde{h}', \tilde{h}'' \in C_B[0, \infty)\}$ associated with norm $\|\tilde{g}\| = \sup_{0 \leq y < \infty} |\tilde{g}(y)|$ and second order Ditzian-Totik modulus of smoothness is presented by

$$\omega_2(\tilde{g}; \sqrt{\delta}) = \sup_{0 < k \leq \sqrt{\delta}} \sup_{y \in [0, \infty)} |\tilde{g}(y+2k) - 2\tilde{g}(y+k) + \tilde{g}(y)|.$$

As described in ([34] by DeVore and Lorentz on page no. 177, Theorem 2.4) as:

$$K_2(\tilde{g}; \delta) \leq \tilde{C}\omega_2(\tilde{g}; \sqrt{\delta}), \quad (8)$$

where $\tilde{C}$ is an absolute constant. To establish the next result, we consider the auxiliary operator defined as:

$$\begin{aligned} \widehat{\mathcal{M}}_{\kappa+s}^{\mathcal{A}}(\tilde{g}; u) &= \mathcal{M}_{\kappa+s}^{\mathcal{A}}(\tilde{g}; u) + \tilde{g}(u) \\ &\quad - \tilde{g}\left(\frac{1}{\kappa+s-1}\left[(\kappa+s)u + \frac{\tilde{\psi}'(h, 1)}{\tilde{\psi}(h, 1)} + \frac{\tilde{\Omega}'(1)}{\tilde{\Omega}(1)} + 1\right]\right). \end{aligned} \quad (9)$$

where $\tilde{g} \in C_B[0, \infty)$, $u \geq 0$ and $\kappa > 1$. From Eq. (9), one can yield

$$\widehat{\mathcal{M}}_{\kappa+s}^{\mathcal{A}}(1; u) = 1, \quad \widehat{\mathcal{M}}_{\kappa+s}^{\mathcal{A}}(\gamma_1(y); u) = 0 \text{ and } |\widehat{\mathcal{M}}_{\kappa+s}^{\mathcal{A}}(\tilde{g}; u)| \leq 3\|\tilde{g}\|. \quad (10)$$

Lemma 3.1. If $u \geq 0$ and $\kappa + s > 2$, one has

$$|\widehat{\mathcal{M}}_{\kappa+s}^{\mathcal{A}}(\tilde{g}; u) - \tilde{g}(u)| \leq \Theta(u)\|\tilde{g}''\|,$$

where $\tilde{g} \in C_B^2[0, \infty)$ and $\Theta(u) = \widehat{\mathcal{M}}_{\kappa+s}^{\mathcal{A}}(\gamma_1(y); u) + (\widehat{\mathcal{M}}_{\kappa+s}^{\mathcal{A}}(\gamma_1(y); u))^2$.

Proof. For $\tilde{g} \in C_B^2[0, \infty)$ and by Taylor expansion, we get

$$\tilde{g}(y) = \tilde{g}(u) + (y - u)\tilde{g}'(u) + \int_u^y (y - v)\tilde{g}''(v)dv. \quad (11)$$

Implementing the auxiliary operators $\widehat{\mathcal{M}}_{\kappa+s}^{\mathcal{A}}(.;.)$ introduced in Eq.(9) to Eq. (11), we get

$$\widehat{\mathcal{M}}_{\kappa+s}^{\mathcal{A}}(\tilde{g}; u) - \tilde{g}(u) = \tilde{g}'(u)\widehat{\mathcal{M}}_{\kappa+s}^{\mathcal{A}}(\gamma_1(y); u) + \widehat{\mathcal{M}}_{\kappa+s}^{\mathcal{A}}\left(\int_u^y (y - v)\tilde{g}''(v)dv; u\right).$$

Using the Eqs. (10) and (11), one yield

$$\begin{aligned} \widehat{\mathcal{M}}_{\kappa+s}^{\mathcal{A}}(\tilde{g}; u) - \tilde{g}(u) &= \widehat{\mathcal{M}}_{\kappa+s}^{\mathcal{A}}\left(\int_u^y (y - v)\tilde{g}''(v)dv; u\right) \\ &= \mathcal{M}_{\kappa+s}^{\mathcal{A}}\left(\int_u^y (y - v)\tilde{g}''(v)dv; u\right) \\ &\quad - \int_u^{\psi_{\kappa+s}} \left(\frac{1}{\kappa + s - 1} \left[(\kappa + s)u + \frac{\tilde{\psi}'(h, 1)}{\tilde{\psi}(h, 1)} + \frac{\tilde{\Omega}'(1)}{\tilde{\Omega}(1)} + 1 \right] - v \right) \tilde{g}''(v)dv, \\ |\widehat{\mathcal{M}}_{\kappa+s}^{\mathcal{A}}(\tilde{g}; u) - \tilde{g}(u)| &\leq \left| \widehat{\mathcal{M}}_{\kappa+s}^{\mathcal{A}}\left(\int_u^y (y - v)\tilde{g}''(v)dv; u\right) \right| \\ &\quad + \left| \int_u^{\psi_{\kappa+s}} \left(\frac{1}{\kappa + s - 1} \left[(\kappa + s)u + \frac{\tilde{\psi}'(h, 1)}{\tilde{\psi}(h, 1)} + \frac{\tilde{\Omega}'(1)}{\tilde{\Omega}(1)} + 1 \right] - v \right) \tilde{g}''(v)dv \right|. \end{aligned} \quad (12)$$

Since,

$$\left| \int_u^y (y - v)\tilde{g}''(v)dv \right| \leq (y - u)^2 \|\tilde{g}''\|, \quad (13)$$

then

$$\begin{aligned} &\left| \int_u^{\psi_{\kappa+s}} \left(\frac{1}{\kappa + s - 1} \left[(\kappa + s)u + \frac{\tilde{\psi}'(h, 1)}{\tilde{\psi}(h, 1)} + \frac{\tilde{\Omega}'(1)}{\tilde{\Omega}(1)} + 1 \right] - v \right) \tilde{g}''(v)dv \right| \\ &\leq \left(\frac{1}{\kappa + s - 1} \left[\frac{\tilde{\psi}'(h, 1)}{\tilde{\psi}(h, 1)} + \frac{\tilde{\Omega}'(1)}{\tilde{\Omega}(1)} + 1 \right] \right)^2 \|\tilde{g}''\|. \end{aligned} \quad (14)$$

In accordance with (12), (13) and (14), we acquire

$$\begin{aligned} |\widehat{\mathcal{M}}_{\kappa+s}^{\mathcal{A}}(\tilde{g}; u) - \tilde{g}(u)| &\leq \left\{ \widehat{\mathcal{M}}_{\kappa+s}^{\mathcal{A}}(\gamma_2(y); u) \right. \\ &\quad \left. + \left(\frac{1}{\kappa + s - 1} \left[\frac{\tilde{\psi}'(h, 1)}{\tilde{\psi}(h, 1)} + \frac{\tilde{\Omega}'(1)}{\tilde{\Omega}(1)} + 1 \right] \right)^2 \right\} \|\tilde{g}''\| \end{aligned}$$

$$= \Theta(u) \|\tilde{g}''\|.$$

This proves the required result. $\square$

Theorem 3.2. For $\tilde{g} \in C_B^2[0, \infty)$, there corresponds a non-negative constant $\tilde{C} > 0$ such that

$$|\mathcal{M}_{\kappa+s}^{\mathcal{A}}(\tilde{g}; u) - \tilde{g}(u)| \leq \tilde{C} \omega_2(\tilde{g}; \sqrt{\Theta(u)}) + \omega(\tilde{g}; \mathcal{M}_{\kappa+s}^{\mathcal{A}}(\gamma_1(y); u),$$

where $\Theta(u)$ is given by in Lemma 3.1.

Proof. Let $\tilde{h} \in C_B^2[0, \infty)$ and $\tilde{g} \in C_B[0, \infty)$. Then, by definition of $\widehat{\mathcal{M}}_{\kappa+s}^{\mathcal{A}}(.;.)$ given in (9), we get

$$\begin{aligned} |\mathcal{M}_{\kappa+s}^{\mathcal{A}}(\tilde{g}; u) - \tilde{g}(u)| &\leq |\widehat{\mathcal{M}}_{\kappa+s}^{\mathcal{A}}(\tilde{g} - \tilde{h}; u)| + |(\tilde{g} - \tilde{h})(u)| + |\widehat{\mathcal{M}}_{\kappa+s}^{\mathcal{A}}(\tilde{g}; u) - \tilde{g}(u)| \\ &\quad + \left| \tilde{g} \left(\frac{1}{\kappa+s-1} \left[(\kappa+s)u + \frac{\tilde{\psi}'(h, 1)}{\tilde{\psi}(h, 1)} + \frac{\tilde{\Omega}'(1)}{\tilde{\Omega}(1)} + 1 \right] \right) - \tilde{g}(u) \right|. \end{aligned}$$

According to Lemma 3.1 and inequalities mentioned in Eq. (10), we acquire

$$\begin{aligned} |\mathcal{M}_{\kappa+s}^{\mathcal{A}}(\tilde{g}; u) - \tilde{g}(u)| &\leq 4\|\tilde{g} - \tilde{h}\| + |\widehat{\mathcal{M}}_{\kappa+s}^{\mathcal{A}}(\tilde{g}; u) - \tilde{g}(u)| \\ &\quad + \left| \tilde{g} \left(\frac{1}{\kappa+s-1} \left[(\kappa+s) + \frac{\tilde{\psi}'(h, 1)}{\tilde{\psi}(h, 1)} + \frac{\tilde{\Omega}'(1)}{\tilde{\Omega}(1)} + 1 \right] \right) - \tilde{g}(u) \right| \\ &\leq 4\|\tilde{g} - \tilde{h}\| + \Theta(u) \|\tilde{h}''\| + \omega(\tilde{g}; \mathcal{M}_{\kappa+s}^{\mathcal{A}}(y - u); u). \end{aligned}$$

Using Eq. (8), we established the required result. $\square$

Further, we address the next result in Lipschitz type space presented by [37] as:

$$Lip_M^{\zeta_1, \zeta_2}(\eta) := \left\{ \tilde{g} \in C_B[0, \infty) : |\tilde{g}(y) - \tilde{g}(u)| \leq \tilde{M} \frac{|y - u|^\eta}{(y + \zeta_1 u + \zeta_2 u^2)^{\frac{\eta}{2}}} : u, y \in (0, \infty) \right\},$$

where $\tilde{M} > 0$, $0 < \eta \leq 1$ and $\zeta_1, \zeta_2 > 0$.

Theorem 3.3. Consider sequence of linear positive operators in (4) and $\tilde{g} \in Lip_M^{\zeta_1, \zeta_2}(\eta)$, one obtain

$$|\mathcal{M}_{\kappa+s}^{\mathcal{A}}(\tilde{g}; u) - \tilde{g}(u)| \leq \tilde{M} \left(\frac{\lambda(u)}{\zeta_1 u + \zeta_2 u^2} \right)^{\frac{\eta}{2}}, \quad (15)$$

where $0 < \eta \leq 1$, $\zeta_1, \zeta_2 \in (0, \infty)$ and $\lambda(y) = \mathcal{M}_{\kappa+s}^{\mathcal{A}}(\gamma_2(y); u)$.

Proof. For $\eta = 1$ and $u \geq 0$, one get

$$\begin{aligned} |\mathcal{M}_{\kappa+s}^{\mathcal{A}}(\tilde{g}; u) - \tilde{g}(u)| &\leq \mathcal{M}_{\kappa+s}^{\mathcal{A}}(|\tilde{g}(y) - \tilde{g}(u)|; u) \\ &\leq \tilde{M} \mathcal{M}_{\kappa+s}^{\mathcal{A}} \left(\frac{|y - u|}{(y + \zeta_1 u + \zeta_2 u^2)^{\frac{1}{2}}}; u \right). \end{aligned}$$

Since $\frac{1}{y + \zeta_1 u + \zeta_2 u^2} < \frac{1}{\zeta_1 u + \zeta_2 u^2}$, for each $u \in (0, \infty)$, we acquire

$$\begin{aligned} |\mathcal{M}_{\kappa+s}^{\mathcal{A}}(\tilde{g}; u) - \tilde{g}(u)| &\leq \frac{\tilde{M}}{(\zeta_1 u + \zeta_2 u^2)^{\frac{1}{2}}} (\mathcal{M}_{\kappa+s}^{\mathcal{A}}(\gamma_2(y); u))^{\frac{1}{2}} \\ &\leq \tilde{M} \left(\frac{\lambda(u)}{\zeta_1 u + \zeta_2 u^2} \right)^{\frac{1}{2}}, \end{aligned}$$

which indicates that the Theorem 3.3 is valid for $\eta = 1$. Next, we examine the case where $\eta \in (0, 1)$ and in accordance with Hölder's inequality by selecting $p = \frac{2}{\eta}$ and $q = \frac{2}{2-\eta}$, we obtain

$$\begin{aligned} |\mathcal{M}_{\kappa+s}^{\mathcal{A}}(\tilde{g}; u) - \tilde{g}(u)| &\leq \left(|\mathcal{M}_{\kappa+s}^{\mathcal{A}}(|\tilde{g}(y) - \tilde{g}(u)|^{\frac{2}{\eta}}; u) \right)^{\frac{\eta}{2}} \\ &\leq \tilde{M} \left(\mathcal{M}_{\kappa+s}^{\mathcal{A}} \left(\frac{|y - u|^2}{(y + \zeta_1 u + \zeta_2 u^2)}; u \right) \right)^{\frac{\eta}{2}}. \end{aligned}$$

Since $\frac{1}{y + \zeta_1 u + \zeta_2 u^2} < \frac{1}{\zeta_1 u + \zeta_2 u^2}$, for all $u \in (0, \infty)$, one get

$$\begin{aligned} |\mathcal{M}_{\kappa+s}^{\mathcal{A}}(\tilde{g}; u) - \tilde{g}(u)| &\leq \tilde{M} \left(\frac{|\mathcal{M}_{\kappa+s}^{\mathcal{A}}(\tilde{g}; u) - \tilde{g}(u)| (|y - u|^2; u)}{\zeta_1 u + \zeta_2 u^2} \right)^{\frac{\eta}{2}} \\ &\leq \tilde{M} \left(\frac{\lambda(u)}{\zeta_1 u + \zeta_2 u^2} \right)^{\frac{\eta}{2}}. \end{aligned}$$

Thus, we obtain the desired result. $\square$

Further, we address the local approximation in terms of the r^{th} order modulus of smoothness, followed by the Lipschitz-type function introduced by Lenze [37] as:

$$\tilde{\omega}_r(\tilde{g}; u) = \sup_{y \neq u, y \in (0, \infty)} \frac{|\tilde{g}(y) - \tilde{g}(u)|}{|y - u|^r}, \quad u \in [0, \infty) \text{ and } r \in (0, 1]. \quad (16)$$

Theorem 3.4. Assume $\tilde{g} \in C_B[0, \infty)$ and $r \in (0, 1]$. Then, for every $u \in [0, \infty)$, we have

$$|\mathcal{M}_{\kappa+s}^{\mathcal{A}}(\tilde{g}; u) - \tilde{g}(u)| \leq \tilde{\omega}_r(\tilde{g}; u) \left(\lambda(u) \right)^{\frac{r}{2}}.$$

Proof. It can be observed that

$$|\mathcal{M}_{\kappa+s}^{\mathcal{A}}(\tilde{g}; u) - \tilde{g}(u)| \leq \mathcal{M}_{\kappa+s}^{\mathcal{A}}(|\tilde{g}(y) - \tilde{g}(u)|; u).$$

Using Eq. (16), one get

$$|\mathcal{M}_{\kappa+s}^{\mathcal{A}}(\tilde{g}; u) - \tilde{g}(u)| \leq \tilde{\omega}_s(\tilde{g}; u) \mathcal{M}_{\kappa+s}^{\mathcal{A}}(|y - u|^r; u).$$

Then by employing Hölder's inequality with $p = \frac{2}{r}$ and $q = \frac{2}{2-r}$, we obtain

$$|\mathcal{M}_{\kappa+s}^{\mathcal{A}}(\tilde{g}; u) - \tilde{g}(u)| \leq \tilde{\omega}_r(\tilde{g}; u) \left(\mathcal{M}_{\kappa+s}^{\mathcal{A}}(|y - u|^2; u) \right)^{\frac{r}{2}}.$$

Thus, we conclude the proof. $\square$

4. Approximation Properties Globally

Consider $v(u) = 1 + u^2, 0 \leq u < \infty$ as weight function. Then, $B_v[0, \infty) = \{\tilde{g}(u) : |\tilde{g}(u)| \leq \tilde{M}_{\tilde{g}}(1 + u^2)\}$, here the constant $\tilde{M}_{\tilde{g}}$ depends on $\tilde{g}$ and $C_v[0, \infty)$ represents the continuous functional space in $B_v[0, \infty)$ along with the norm $\|\tilde{g}\|_v = \sup_{u \in [0, \infty)} \frac{|\tilde{g}(u)|}{v(u)}$ and $C_v^{\tilde{k}}[0, \infty) = \{\tilde{g} \in C_v[0, \infty) : \lim_{u \rightarrow \infty} \frac{\tilde{g}(u)}{v(u)} = \tilde{k}, \text{ where constant } \tilde{k} \text{ depends on } \tilde{g}\}$.

If $\tilde{g}$ is a function defined on $[0, b]$ where $b > 0$. Then, Ditzian-Totik modulus of continuity is given by

$$\omega_b(\tilde{g}, \delta) = \sup_{|y-u| \leq \delta} \sup_{u, y \in [0, b]} |\tilde{g}(y) - \tilde{g}(u)|. \quad (17)$$

It is straightforward to observe that for $\tilde{g} \in C_v[0, \infty)$, the modulus of continuity defined in Eq. (17) tends to zero

Remark 4.1. In this article, we employ the test function defined by

$$\tilde{g}_\theta(y) = y^\theta, \quad \theta \in \{0, 1, 2\}.$$

Theorem 4.2. Let $\tilde{g} \in C_v[0, \infty)$ and $\omega_{b+1}(\tilde{g}; \delta)$ denote the modulus of smoothness defined on $[0, b+1] \subset [0, \infty)$. Then, for $y \in [0, b]$, we obtain

$$\|\mathcal{M}_{\kappa+s}^{\mathcal{A}}(\cdot; \cdot) - \tilde{g}\|_{C[0,b]} \leq 4\tilde{M}_{\tilde{g}}(1+b^2)\delta_s(b) + 2\omega_{b+1}(\tilde{g}; \sqrt{\delta_s(b)}),$$

where $\delta_s(b) = \max_{u \in [0,b]} \mathcal{M}_{\kappa+s}^{\mathcal{A}}(\gamma_2; u)$.

Proof. For any $u \in [0, b]$ and $y \in [0, \infty)$, we have

$$|\tilde{g}(y) - \tilde{g}(u)| \leq 4\tilde{M}_{\tilde{g}}(1+b^2)(y-u)^2 + \left(1 + \frac{|y-u|}{\delta}\right)\omega_{b+1}(\tilde{g}; \delta).$$

Implementing operator $\mathcal{M}_{\kappa+s}^{\mathcal{A}}(\cdot; \cdot)$ on both the sides, we acquire

$$\begin{aligned} |\mathcal{M}_{\kappa+s}^{\mathcal{A}}(\tilde{g}; u) - \tilde{g}(u)| &\leq 4\tilde{M}_{\tilde{g}}(1+b^2)\mathcal{M}_{\kappa+s}^{\mathcal{A}}(\gamma_2; u) \\ &\quad + \left(1 + \frac{\mathcal{M}_{\kappa+s}^{\mathcal{A}}(|y-u|; u)}{\delta}\right)\omega_{b+1}(\tilde{g}; \delta). \end{aligned}$$

Now, in accordance with Lemma 1.5 and $x \in [0, b]$, one has

$$|\mathcal{M}_{\kappa+s}^{\mathcal{A}}(\cdot; \cdot) - \tilde{g}| \leq 4\tilde{M}_{\tilde{g}}(1+b^2)\delta_s(b) + \left(1 + \frac{\sqrt{\delta_s(b)}}{\delta}\right)\omega_{b+1}(\tilde{g}; \delta).$$

By selecting $\delta = \delta_s(b)$, desired result can easily be obtained. $\square$

Theorem 4.3. ([38], [39]) Assume that the sequence of linear positive operators $(L_{\kappa+s})_{\kappa \geq 1}$ mapping from $C_v[0, \infty)$ to $B_v[0, \infty)$ meets the conditions

$$\lim_{\kappa \rightarrow \infty} \|L_{\kappa+s}(\tilde{g}_\theta; \cdot) - \tilde{g}_\theta\|_v = 0, \quad \text{where } \theta = 0, 1, 2,$$

thus, for $\tilde{g} \in C_v^k[0, \infty)$, we get

$$\lim_{\kappa \rightarrow \infty} \|L_{\kappa+s}(\tilde{g}; \cdot) - \tilde{g}\|_v = 0.$$

Theorem 4.4. Let $\tilde{g} \in C_v^k[0, \infty)$. Then, we obtain

$$\lim_{\kappa \rightarrow \infty} \|\mathcal{M}_{\kappa+s}^{\mathcal{A}}(\tilde{g}; \cdot) - \tilde{g}\|_v = 0.$$

Proof. To prove the result of Theorem 4.4, it is enough to verify that

$$\lim_{\kappa \rightarrow \infty} \|\mathcal{M}_{\kappa+s}^{\mathcal{A}}(\tilde{g}_\theta; \cdot) - \tilde{g}_\theta\|_v = 0, \quad \text{for } \theta = 0, 1, 2.$$

Considering the Lemma 1.4, one can see $\|\mathcal{M}_{\kappa+s}^{\mathcal{A}}(\tilde{g}_0; \cdot) - 1\|_v = 0$, here $\kappa \rightarrow \infty$ and restrict with $\kappa > 2$, also

$$\begin{aligned} \|\mathcal{M}_{\kappa+s}^{\mathcal{A}}(\tilde{g}_1; \cdot) - \tilde{g}_1\|_{v(u)} &= \sup_{u \in [0, \infty)} \frac{1}{v(u)} \left| \frac{1}{\kappa+s-1} \left[\frac{\tilde{\psi}'(h, 1)}{\tilde{\psi}(h, 1)} + \frac{\tilde{\Omega}'(1)}{\tilde{\Omega}(1)} + 1 \right] \right| \\ &= \frac{1}{\kappa+s-1} \left[\frac{\tilde{\psi}'(h, 1)}{\tilde{\psi}(h, 1)} + \frac{\tilde{\Omega}'(1)}{\tilde{\Omega}(1)} + 1 \right] \sup_{u \in [0, \infty)} \frac{1}{1+u^2}. \end{aligned}$$

For a large value of κ , we get $\|\mathcal{M}_{\kappa+s}^{\mathcal{A}}(\tilde{g}_1; \cdot) - \tilde{g}_1\|_v \rightarrow 0$.

Now,

$$\begin{aligned} \|\mathcal{M}_{\kappa+s}^{\mathcal{A}}(\tilde{g}_2; \cdot) - \tilde{g}_2\|_v &\leq \left(\frac{3(\kappa+s)-2}{(\kappa+s-2)(\kappa+s-1)} \right) \sup_{u \in [0, \infty)} \frac{u^2}{1+u^2} \\ &\quad + \left(\frac{4(\kappa+s)}{(\kappa+s-2)(\kappa+s-1)} \right) \sup_{u \in [0, \infty)} \frac{u}{1+u^2} \\ &\quad + \left(\frac{2(\kappa+s)u+4}{(\kappa+s-2)(\kappa+s-1)} \left[\frac{\tilde{\psi}'(h,1)}{\tilde{\psi}(h,1)} + \frac{\tilde{\Omega}'(1)}{\tilde{\Omega}(1)} \right] \right) \sup_{u \in [0, \infty)} \frac{1}{1+u^2} \\ &\quad + \left(\frac{1}{(\kappa+s-2)(\kappa+s-1)} \left(2 \frac{\tilde{\psi}'(h,1)\tilde{\Omega}'(1)}{\tilde{\Omega}(1)\tilde{\psi}(h,1)} + \frac{\tilde{\psi}''(h,1)}{\tilde{\psi}(h,1)} \right. \right. \\ &\quad \left. \left. + \frac{\tilde{\Omega}''(1)}{\tilde{\Omega}(1)} + 2 \right) \right) \sup_{y \in [0, \infty)} \frac{1}{1+u^2}. \end{aligned}$$

Which implies $\|\mathcal{M}_{\kappa+s}^{\mathcal{A}}(\tilde{g}_2; \cdot) - \tilde{g}_2\|_v \rightarrow 0$ as $\kappa \rightarrow \infty$. Thus, we conclude the proof of the Theorem 4.4 $\square$

Theorem 4.5. Let $\tilde{g} \in C_v^{\tilde{\kappa}}[0, \infty)$ and $\zeta > 0$. Then,

$$\lim_{\kappa \rightarrow \infty} \sup_{u \in [0, \infty)} \frac{\mathcal{M}_{\kappa+s}^{\mathcal{A}}(\tilde{g}; u) - \tilde{g}(u)}{(1+u^2)^{1+\zeta}} = 0.$$

Proof. Since $|\tilde{g}(x)| \leq \|\tilde{g}\|_v(1+u^2)$, for any real fixed number $u_0 > 0$, we get

$$\begin{aligned} \sup_{u \in [0, \infty)} \frac{\mathcal{M}_{\kappa+s}^{\mathcal{A}}(\tilde{g}; u) - \tilde{g}(u)}{(1+u^2)^{1+\zeta}} &\leq \sup_{u \leq u_0} \frac{\mathcal{M}_{\kappa+s}^{\mathcal{A}}(\tilde{g}; u) - \tilde{g}(u)}{(1+u^2)^{1+\zeta}} + \sup_{u \geq u_0} \frac{\mathcal{M}_{\kappa+s}^{\mathcal{A}}(\tilde{g}; u) - \tilde{g}(u)}{(1+u^2)^{1+\zeta}} \\ &\leq \|\mathcal{M}_{\kappa+s}^{\mathcal{A}}(\tilde{g}; u) - \tilde{g}(u)\|_{C[0, u_0]} \\ &\quad + \|\tilde{g}\|_v \sup_{u \geq u_0} \frac{\mathcal{M}_{\kappa+s}^{\mathcal{A}}(1+y^2; u)}{(1+u^2)^{1+\zeta}} + \sup_{u \geq u_0} \frac{|\tilde{g}(u)|}{(1+u^2)^{1+\zeta}} \\ &= \tilde{R}_1 + \tilde{R}_2 + \tilde{R}_3, \quad \text{say.} \end{aligned} \tag{18}$$

Now,

$$\tilde{R}_3 = \sup_{u \geq u_0} \frac{|\tilde{g}(u)|}{(1+u^2)^{1+\zeta}} \leq \sup_{u \geq u_0} \frac{\|\tilde{g}\|_v(1+u^2)}{(1+u^2)^{1+\zeta}} \leq \frac{\|\tilde{g}\|_v}{(1+u_0^2)^{\zeta}}.$$

In view of Lemma 1.4, it gives

$$\lim_{\kappa \rightarrow \infty} \sup_{u \in [u_0, \infty)} \frac{\mathcal{M}_{\kappa+s}^{\mathcal{A}}(1+y^2; u)}{1+u^2} = 1.$$

Therefore, for any arbitrary $\epsilon > 0$, there corresponds $\kappa_1 \in \mathbb{N}$ with

$$\sup_{u \in [u_0, \infty)} \frac{\mathcal{M}_{\kappa+s}^{\mathcal{A}}(1+y^2; u)}{1+u^2} \leq \frac{(1+u_0^2)^{\zeta}}{\|\tilde{g}\|_v} \frac{\epsilon}{3} + 1, \text{ for all } \kappa \geq \kappa_1.$$

Therefore

$$\tilde{R}_2 = \|\tilde{g}\|_v \sup_{u \in [u_0, \infty)} \frac{\mathcal{M}_{\kappa+s}^{\mathcal{A}}(1+y^2; u)}{(1+u^2)^{1+\zeta}} \leq \frac{\|\tilde{g}\|_v}{(1+u_0^2)^{\zeta}} + \frac{\epsilon}{3}, \text{ for all } \kappa \geq \kappa_1. \tag{19}$$

Hence, we get

$$\tilde{R}_2 + \tilde{R}_3 < 2 \frac{\|\tilde{g}\|_v}{(1+u^2)^\zeta} + \frac{\epsilon}{3}.$$

If we take u_0 to be so large that $\frac{\|\tilde{g}\|_v}{(1+u^2)^\zeta} < \frac{\epsilon}{6}$, then, we have

$$\tilde{R}_2 + \tilde{R}_3 < \frac{2\epsilon}{3} \text{ for all } \kappa \geq \kappa_1. \quad (20)$$

Now, from Theorem 4.2, there corresponds $\kappa_2 > \kappa$ with

$$\tilde{R}_1 = \|\mathcal{M}_{\kappa+s}^{\mathcal{A}}(\tilde{g}; \cdot) - \tilde{g}\|_{C[0, u_0]} < \frac{\epsilon}{3} \text{ for all } \kappa_2 \geq \kappa. \quad (21)$$

Let $\kappa_3 = \max(\kappa_1, \kappa_2)$. Then, using the Eqs. (18), (20) and (21), we get

$$\sup_{u \in [0, \infty)} \frac{|\mathcal{M}_{\kappa+s}^{\mathcal{A}}(\tilde{g}; u) - \tilde{g}(u)|}{(1+u^2)^{1+\zeta}} < \epsilon,$$

which, completes the proof. $\square$

5. A-Statistical Approximation

We revisit some notation from [40]. Suppose that $B = (b_{(\kappa+s)\mu})$ is an infinite, non-negative summability matrix. A sequence $u := (u_\mu)$ is A-statistically convergent to L , denoted as $st_B - \lim u = L$, if for each $\epsilon > 0$

$$\lim_{\kappa} \sum_{\mu: |u_\mu - L| \geq \epsilon} b_{(\kappa+s)\mu} = 0.$$

Let $q = (q_{(\kappa+s)})$ be a sequence such that the following assertions are true

$$st_B - \lim_{\kappa} q_{(\kappa+s)} = 1 \text{ and } st_B - \lim_{\kappa} q_{(\kappa+s)}^\kappa = b, \quad 0 \leq b < 1. \quad (22)$$

Theorem 5.1. Consider $B = (b_{(\kappa+s)\mu})$ be a non-negative regular summability matrix and sequence $q = (q_{\kappa+s})$ along with condition (22), $q_{\kappa+s} \in (0, 1)$, $\kappa \in \mathbb{N}$. Then, for each $\tilde{g} \in C_v^0[0, \infty)$, $st_B - \lim_{\kappa} \|\mathcal{M}_{\kappa+s}^{\mathcal{A}}(\tilde{g}; \cdot) - \tilde{g}\|_v = 0$.

Proof. In accordance with Lemma 1.4, one has

$$st_B - \lim_{\kappa} \|\mathcal{M}_{\kappa+s}^{\mathcal{A}}(\tilde{g}_0; \cdot) - \tilde{g}_0\|_v = 0.$$

Also,

$$\begin{aligned} \|\mathcal{M}_{\kappa+s}^{\mathcal{A}}(\tilde{g}_1; \cdot) - \tilde{g}_1\|_v &= \sup_{u \in [0, \infty)} \frac{1}{1+u^2} \left| \frac{1}{\kappa+s-1} \left[(\kappa+s)u + 1 + \frac{\tilde{\psi}'(h, 1)}{\tilde{\psi}(h, 1)} + \frac{\tilde{\Omega}'(1)}{\tilde{\Omega}(1)} \right] - u \right| \\ &= \frac{1}{\kappa+s-1} \sup_{u \in [0, \infty)} \frac{u}{1+u^2} \\ &\quad + \frac{1}{\kappa+s-1} \left[1 + \frac{\tilde{\psi}'(h, 1)}{\tilde{\psi}(h, 1)} + \frac{\tilde{\Omega}'(1)}{\tilde{\Omega}(1)} \right] \sup_{u \in [0, \infty)} \frac{1}{1+u^2}. \end{aligned}$$

Now

$$\tilde{K}_1 := \left\{ \kappa : \|\mathcal{M}_{\kappa+s}^{\mathcal{A}}(\tilde{g}_1; \cdot) - \tilde{g}_1\| \geq \epsilon \right\},$$

$$\tilde{K}_2 := \left\{ \kappa : \frac{1}{\kappa + s - 1} \geq \frac{\epsilon}{2} \right\},$$

$$\tilde{K}_3 := \left\{ \kappa : \frac{1}{\kappa + s - 1} \left[1 + \frac{\tilde{\psi}'(h, 1)}{\tilde{\psi}(h, 1)} + \frac{\tilde{\Omega}'(1)}{\tilde{\Omega}(1)} \right] \geq \frac{\epsilon}{2} \right\}.$$

Which implies that $\tilde{K}_1 \subseteq \tilde{K}_2 \cup \tilde{K}_3$, this shows that

$$\sum_{\mu \in \tilde{K}_1} b_{(\kappa+s)\mu} \leq \sum_{\mu \in \tilde{K}_2} b_{(\kappa+s)\mu} + \sum_{\mu \in \tilde{K}_3} b_{(\kappa+s)\mu}.$$

Therefore, we get

$$st_B - \lim_{\kappa} \|\mathcal{M}_{\kappa+s}^{\mathcal{A}}(\bar{g}_1; \cdot) - \bar{g}_1\|_v = 0. \quad (23)$$

Now by using Lemma 1.4, we have

$$\begin{aligned} \|\mathcal{M}_{\kappa+s}^{\mathcal{A}}(\bar{g}_2; \cdot) - \bar{g}_2\|_{1+u^2} &\leq \sup_{u \in [0, \infty)} \frac{1}{v(u)} \left| \frac{1}{(\kappa + s - 1)(\kappa + s - 2)} \left[(3(\kappa + s) - 2)u^2 \right. \right. \\ &\quad \left. \left. + 4(\kappa + s)u + (2(\kappa + s)u + 4) \left(\frac{\tilde{\psi}'(h, 1)}{\tilde{\psi}(h, 1)} + \frac{\tilde{\Omega}'(1)}{\tilde{\Omega}(1)} \right) \right. \right. \\ &\quad \left. \left. + 2 \frac{\tilde{\psi}'(h, 1)\tilde{\Omega}'(1)}{\tilde{\psi}(h, 1)\tilde{\Omega}(1)} + \frac{\tilde{\psi}''(h, 1)}{\tilde{\psi}(h, 1)} + \frac{\tilde{\Omega}''(1)}{\tilde{\Omega}(1)} + 2 \right] \right|. \end{aligned}$$

For a given $\epsilon > 0$, we have the following sets

$$\begin{aligned} \tilde{H}_1 &:= \left\{ \kappa : \|\mathcal{M}_{\kappa+s}^{\mathcal{A}}(\bar{g}_2; \cdot) - \bar{g}_2\|_v \geq \epsilon \right\}, \\ \tilde{H}_2 &:= \left\{ \kappa : \frac{3(\kappa + s) - 2}{(\kappa + s - 2)(\kappa + s - 1)} \geq \frac{\epsilon}{4} \right\}, \\ \tilde{H}_3 &:= \left\{ \kappa : \frac{4(\kappa + s)}{(\kappa + s - 2)(\kappa + s - 1)} \geq \frac{\epsilon}{4} \right\}, \\ \tilde{H}_4 &:= \left\{ \kappa : \frac{1}{(\kappa + s - 2)(\kappa + s - 1)} \left[\frac{\tilde{\psi}'(h, 1)}{\tilde{\psi}(h, 1)} + \frac{\tilde{\Omega}'(1)}{\tilde{\Omega}(1)} \right] \geq \frac{\epsilon}{4} \right\}, \\ \tilde{H}_5 &:= \left\{ \kappa : \frac{1}{(\kappa + s - 2)(\kappa + s - 1)} \left[2 \frac{\tilde{\psi}'(h, 1)\tilde{\Omega}'(1)}{\tilde{\psi}(h, 1)\tilde{\Omega}(1)} + \frac{\tilde{\psi}''(h, 1)}{\tilde{\psi}(h, 1)} + \frac{\tilde{\Omega}''(1)}{\tilde{\Omega}(1)} + 2 \right] \geq \frac{\epsilon}{4} \right\}. \end{aligned}$$

It can be seen that $\tilde{H}_1 \subseteq \tilde{H}_2 \cup \tilde{H}_3 \cup \tilde{H}_4 \cup \tilde{H}_5$. Therefore, we acquire

$$\sum_{\mu \in \tilde{H}_1} b_{(\kappa+s)\mu} \leq \sum_{\mu \in \tilde{H}_2} b_{(\kappa+s)\mu} + \sum_{\mu \in \tilde{H}_3} b_{(\kappa+s)\mu} + \sum_{\mu \in \tilde{H}_4} b_{(\kappa+s)\mu} + \sum_{\mu \in \tilde{H}_5} b_{(\kappa+s)\mu}.$$

As $\kappa \rightarrow \infty$, we have

$$st_B - \lim_{\kappa} \|\mathcal{M}_{\kappa+s}^{\mathcal{A}}(\bar{g}_2; u) - \bar{g}_2\|_v = 0. \quad (24)$$

Thus, we concludes the proof of the Theorem 5.1. $\square$

Next, we will examine the convergence rate of A-Statistical approximation with respect to Peetre's K-functional for the operators $\mathcal{M}_{\kappa+s}^{\mathcal{A}}(\cdot; \cdot)$.

Theorem 5.2. Let $\tilde{g} \in C_B^2[0, \infty)$. Then,

$$st_B - \lim_{\kappa} \|\mathcal{M}_{\kappa+s}^{\mathcal{A}}(\tilde{g}; \cdot) - \tilde{g}\|_{C_B[0, \infty)} = 0.$$

Proof. Considering Taylor's result, we have

$$\tilde{g}(y) = \tilde{g}(u) + \tilde{g}'(u)(y - u) + \frac{1}{2}\tilde{g}''(\eta)(y - u)^2,$$

where $v \leq \eta \leq u$. Operating $\mathcal{M}_{\kappa+s}^{\mathcal{A}}(\cdot; \cdot)$, on both sides in above equation, one get

$$\mathcal{M}_{\kappa+s}^{\mathcal{A}}(\tilde{g}; u) - \tilde{g}(u) = \tilde{g}'(u)\mathcal{M}_{\kappa+s}^{\mathcal{A}}(\eta_1; u) + \frac{1}{2}\tilde{g}''(\eta)\mathcal{M}_{\kappa+s}^{\mathcal{A}}(\eta_2; u),$$

which yields that

$$\begin{aligned} \|\mathcal{M}_{\kappa+s}^{\mathcal{A}}(\tilde{g}; \cdot) - \tilde{g}\|_{C_B[0, \infty)} &\leq \|\tilde{g}'\|_{C_B[0, \infty)}\|\mathcal{M}_{\kappa+s}^{\mathcal{A}}(\tilde{g}_1 - \cdot; \cdot)\|_{C_B[0, \infty)} \\ &\quad + \|\tilde{g}''\|_{C_B[0, \infty)}\|\mathcal{M}_{\kappa+s}^{\mathcal{A}}(\tilde{g}_1 - \cdot; \cdot)^2\|_{C_B[0, \infty)} \\ &= \tilde{W}_1 + \tilde{W}_2, \quad \text{say.} \end{aligned} \quad (25)$$

Based on Eqs. (23) and (24), it follows that

$$\begin{aligned} \lim_{\kappa} \sum_{\mu \in \mathbb{N}: \tilde{W}_1 \geq \frac{\epsilon}{2}} b_{(\kappa+s)\mu} &= 0, \\ \lim_{\kappa} \sum_{\mu \in \mathbb{N}: \tilde{W}_2 \geq \frac{\epsilon}{2}} b_{(\kappa+s)\mu} &= 0. \end{aligned}$$

From Eq. (25), we have

$$\lim_{\kappa} \sum_{\mu \in \mathbb{N}: \|\mathcal{M}_{\kappa+s}^{\mathcal{A}}(\tilde{g}; \cdot) - \tilde{g}\|_{C_B[0, \infty)} \geq \epsilon} b_{(\kappa+s)\mu} \leq \lim_{\kappa} \sum_{\mu \in \mathbb{N}: \tilde{W}_1 \geq \frac{\epsilon}{2}} b_{(\kappa+s)\mu} + \lim_{\kappa} \sum_{\mu \in \mathbb{N}: \tilde{W}_2 \geq \frac{\epsilon}{2}} b_{(\kappa+s)\mu}.$$

Thus, $st_B - \lim_{\kappa} \|\mathcal{M}_{\kappa+s}^{\mathcal{A}}(\tilde{g}; \cdot) - \tilde{g}\|_{C_B[0, \infty)} \rightarrow 0$ as $\kappa \rightarrow \infty$.

Hence, we arrive the proof. $\square$

Theorem 5.3. For $\tilde{g} \in C_B^2[0, \infty)$,

$$\|\mathcal{M}_{\kappa+s}^{\mathcal{A}}(\tilde{g}; \cdot) - \tilde{g}\|_{C_B[0, \infty)} \leq \tilde{M}\omega_2(\tilde{g}; \sqrt{\delta}),$$

where $\delta = \|\mathcal{M}_{\kappa+s}^{\mathcal{A}}(\tilde{g}_1 - \cdot; \cdot)\|_{C_B[0, \infty)} + \|\mathcal{M}_{\kappa+s}^{\mathcal{A}}(\tilde{g}_1 - \cdot)^2; \cdot\|_{C_B[0, \infty)}$ and $\|\tilde{g}\|_{C_B^2[0, \infty)} = \|\tilde{g}\|_{C_B[0, \infty)} + \|\tilde{g}'\|_{C_B[0, \infty)} + \|\tilde{g}''\|_{C_B[0, \infty)}$.

Proof. Let $\tilde{f} \in C_B^2[0, \infty)$. Using Eq. (25), one obtain

$$\begin{aligned} \|\mathcal{M}_{\kappa+s}^{\mathcal{A}}(\tilde{f}) - \tilde{f}\|_{C_B[0, \infty)} &\leq \|\tilde{f}'\|_{C_B[0, \infty)}\|\mathcal{M}_{\kappa+s}^{\mathcal{A}}(\tilde{g}_1 - \cdot; \cdot)\|_{C_B[0, \infty)} \\ &\quad + \frac{1}{2}\|\tilde{f}''\|_{C_B[0, \infty)}\|\mathcal{M}_{\kappa+s}^{\mathcal{A}}(\tilde{g}_1 - \cdot)^2; \cdot\|_{C_B[0, \infty)} \\ &\leq \delta\|\tilde{f}\|_{C_B^2[0, \infty)}. \end{aligned} \quad (26)$$

For each $\tilde{g} \in C_B[0, \infty)$ and $\tilde{f} \in C_B^2[0, \infty)$, Using Eq. (26), we acquire

$$\begin{aligned} \|\mathcal{M}_{\kappa+s}^{\mathcal{A}}(\tilde{g}; \cdot) - \tilde{g}\|_{C_B[0, \infty)} &\leq \|\mathcal{M}_{\kappa+s}^{\mathcal{A}}(\tilde{g}; \cdot) - \mathcal{M}_{\kappa+s}^{\mathcal{A}}(\tilde{f}; \cdot)\|_{C_B[0, \infty)} \\ &\quad + \|\mathcal{M}_{\kappa+s}^{\mathcal{A}}(\tilde{f}; \cdot) - \tilde{f}\|_{C_B[0, \infty)} + \|\tilde{f} - \tilde{g}\|_{C_B[0, \infty)} \\ &\leq 2\|\tilde{f} - \tilde{g}\|_{C_B[0, \infty)} + \|\mathcal{M}_{\kappa+s}^{\mathcal{A}}(\tilde{f}; \cdot) - \tilde{f}\|_{C_B[0, \infty)} \\ &\leq 2\|\tilde{f} - \tilde{g}\|_{C_B[0, \infty)} + \delta\|\tilde{f}\|_{C_B^2[0, \infty)}. \end{aligned}$$

Considering Peetre's K-functional, we obtain

$$\begin{aligned}\|\mathcal{M}_{k+s}^{\mathcal{A}}(\tilde{g}; \cdot) - \tilde{g}\|_{C_B[0,\infty)} &\leq 2K_2(\tilde{g}; \delta) \\ &\leq 2\tilde{C}\omega_2(\tilde{g}; \sqrt{\delta}) \\ &= \tilde{M}\omega_2(\tilde{g}; \sqrt{\delta})\end{aligned}$$

where $\tilde{M} = 2\tilde{C}$ is an absolute constant. Therefore,

$$\|\mathcal{M}_{k+s}^{\mathcal{A}}(\tilde{g}; \cdot) - \tilde{g}\|_{C_B[0,\infty)} \leq \tilde{M}\omega_2(\tilde{g}; \sqrt{\delta}).$$

which concludes the proof of desired result. $\square$

6. Conclusion

In this article, we introduce a sequence of positive linear operators using generalized Appell polynomials in Beta form. These operators are designed to approximate functions defined on a Lebesgue measurable space and are known as modified Szász-Beta type operators introduced in (4). Moreover, we derive estimates to establish the rate of convergence and the accuracy of approximation. Further, we explore various aspects of approximation, including local and global results, as well as A-statistical approximation, utilizing these operators to obtain enhanced approximations across different functional spaces. These operators provide better flexibility in approximations compared to the operators defined in (3). In addition, these sequence of operators discuss approximations in wider class in comparison of operators defined in (3).

Conflicts of interest

There are no conflicts of interest, according to the authors.

Data availability

No data were used to support this study.

References

- [1] K. Weierstrass, 'Über die analytische Darstellbarkeit sogenannter willkürlicher Functionen einer reellen Veränderlichen, Sitzungsberichte der Königlich Preussischen Akademie der Wissenschaften zu Berlin, **2**, 633-639, (1885).
- [2] S. N. Bernstein, *Démonstration du théorème de Weierstrass fondée sur le calcul de probabilités*, Commun. Soc. Math. Kharkow., **13** (2), 1-2, (1912/1913).
- [3] O. Szász, *Generalization of S. Bernstein's polynomials to the infinite interval*, J. Research Nat. Bur. Standards, **45**, 239-245, (1950).
- [4] F. Schurer, *Linear positive operators in approximation theory*, Math. Inst. Techn. Univ. Delft Report, (1962).
- [5] N. Rao, P. Kumar, N. K. Jha, and Md. Heshamuddin, *Quantitative estimators of extended Beta type Szász-Schurer operators*, Palestine Journal of Mathematics, **14** (2), 463-473, (2025).
- [6] N. Rao, M. Farid, and N. K. Jha, *Szász-Integral operators linking general-Appell Polynomials and approximation*, AIMS Math., **10** (6), 13836-13854, (2025).
- [7] N. Raza and M. Kumar, *Approximation with Chlodowsky variant of Kantorovich-Stancu-operators employing associated λ -polynomials*, J. Inequal. Appl., **2024** (1), 1-19, (2024).
- [8] N. Raza, M. Kumar, and M. Mursaleen, *On approximation operators involving Tricomi function*, Bull. Malaysian Math. Sci. Soc., **47** (5), 154, (2024).
- [9] R. Aslan, *Some approximation properties of Riemann-Liouville type fractional Bernstein-Stancu-Kantorovich operators with order of α* , Iran. J. Sci., 1-14, (2024).
- [10] R. Aslan, *Rate of approximation of blending type modified univariate and bivariate λ -Schurer-Kantorovich operators*, Kuwait J. Sci., **51** (1), 100168, (2024).
- [11] F. Özger, *On new Bézier bases with Schurer polynomials and corresponding results in approximation theory*, Commun. Fac. Sci. Univ. Ank. Ser. Al. Math. Stat, **69** (1), 376-393, (2020).
- [12] F. Özger, M. T. Ersoy, and O. Z. Özger, *Existence of solutions: Investigating Fredholm integral equations via a fixed-point theorem*, **13**, 261, (2024).
- [13] Q. B. Cai and R. Aslan, *Note on a new construction of Kantorovich form q -Bernstein operators related to shape parameter λ* , Comput. Model. Eng. Sci, **130** (3), 1479-1493, (2022).

- [14] Q. B. Cai, R. Aslan, F. Özger, and H. M. Srivastava, *Approximation by a new Stancu variant of generalized (λ, μ) -Bernstein operators*, Alex. Eng. J., **107**, 205-214, (2024).
- [15] N. L. Braha, *Some summability methods for Dunkl–Gamma-type operators including Appell polynomials*, Math. Methods Appl. Sci., (2024).
- [16] N. L. Braha, T. Mansour, and M. Mursaleen, *Some Approximation Properties of Parametric Baskakov–Schurer–Szász Operators Through a Power Series Summability Method*, Complex Anal. Oper. Theory, **18** (3), pages- 71, (2024).
- [17] M. A. Mursaleen, M. Heshamuddin, N. Rao, B. K. Sinha, and A. K. Yadav, *Hermite polynomials linking Szász–Durrmeyer operators*, Comp. Appl. Math., **43** (223), 407-421, (2024).
- [18] M. A. Mursaleen, M. Nasiruzzaman, N. Rao, M. Dilshad, and K. S. Nisar, *Approximation by the modified λ -Bernstein-polynomial in terms of basis function*, AIMS Math., **9** (2), 4409-4426, (2024).
- [19] M. A. Mursaleen, M. Nasiruzzaman, and N. Rao, *On the Approximation of Szász–Jakimovski–Leviatan beta type integral operators enhanced by Appell polynomials*, Iran. J. Sci., 1-10, (2025).
- [20] K. J. Ansari, M. Mursaleen, and S. Rahman Özger, *Approximation by Jakimovski–Leviatan operators of Durrmeyer type involving multiple Appell polynomials*, Rev. R. Acad. Cienc. Exactas Fis. Nat. Ser. A Math. RACSAM, **113**, 1007-1024, (2019).
- [21] K. J. Ansari, F. Özger, and O. Z. Özger, *Numerical and theoretical approximation results for Schurer–Stancu operators with shape parameter λ* , Comput. Appl. Math., **41** (4), 181, (2022).
- [22] S. A. Mohiuddine, T. Acar, and A. Alotaibi, *Construction of a new family of Bernstein–Kantorovich operators*, Math. Methods Appl. Sci., **40**, 7749 – 7759, (2017).
- [23] S. A. Mohiuddine, N. Ahmad, F. Özger, A. Alotaibi, and B. Hazarika, *Approximation by the parametric generalization of Baskakov–Kantorovich operators linking with Stancu operators*, Iran. J. Sci. Technol. Trans., **45** (2), 593-605, (2021).
- [24] M. Mursaleen, K.J. Ansari, and A. Khan, *Approximation properties and error estimation of q -Bernstein shifted operators*, Numer. Algorithms, **84**, 207-227, (2020).
- [25] M. Mursaleen, A. Naaz, and A. Khan, *Improved approximation and error estimations by King type (p, q) -Szász–Mirakjan Kantorovich operators*, Appl. Math. Comput, **348**, 2175-185, (2019).
- [26] N. Rao, M. Shahzad, and N. K. Jha, *Study of two dimensional α -modified Bernstein bi-variate operators*, Filomat, **39** (5), 1509-1522, (2025).
- [27] N. Rao, N. K. Jha, and M. Shahzad, *On a new sequence of Bernstein–Chlodowsky operators and approximation properties*, J. Math. Sci., (2025).
- [28] A. Khan, M. Mansoori, K. Khan, and M. Mursaleen, *Phillips-type q -Bernstein operators on triangles*, J. Funct. Spaces, **2021**, 1-13, (2021).
- [29] Md. Nasiruzzaman, *Approximation properties by Szász–Mirakjan operators to bivariate functions via Dunkl analogue*, Iran. J. Sci. Technol. Trans., **45**, 259-269, (2021).
- [30] A. M. Acu, T. Acar, and V. A. Radu, *Approximation by modified U_n^p operators*, Rev. R. Acad. Cienc. Exactas Fis. Nat. Ser. A Math. RACSAM, **113**, 2715-2729, (2019).
- [31] A. M. Acu, P. Agrawal, and D. Kumar, *Approximation properties of modified q -Bernstein–Kantorovich operators*, Commun. Fac. Sci. Univ. Ank. Ser. A1 Math. Stat. **68** (2), 2170-2197, (2019).
- [32] N. K. Jha and N. Rao, *On generalized Bernstein Kantorovich Schurer type operators and its approximation behaviour*, Filomat, **39** (10), 3249-3262, (2025).
- [33] N. Raza, M. Kumar, and M. Mursaleen, *Approximation with Szász–Chlodowsky operators employing general-Appell polynomials*, J. Inequal. Appl., **26** (2024), (2024).
- [34] R. A. DeVore and G. G. Lorentz, *Constructive Approximation*, Grudlehren der Mathematischen Wissenschaften Fundamental principales of Mathematical Sciences, Springer-Verlag, Berlin, (1993).
- [35] O. Shisha and B. Mond, *The degree of convergence of linear positive operators*, Proc. Nat. Acad. Sci. USA, **60**, 1196-1200, (1968).
- [36] Korovkin, P.P., *On convergence of linear positive operators in the space of continuous functions*, Dokl. Akad. Nauk. SSSR, **90**, 961–964, (1953).
- [37] B. Lenze, *On Lipschitz type maximal functions and their smoothness spaces*, Nederl Akad Indag Math. **50**, 53-63, (1988).
- [38] A. D. Gadjiev, *The convergence problem for a sequence of positive linear operators on bounded sets and theorems analogous to that of P. P. Korovkin*, Dokl. Akad. Nauk SSSR **218** (5), (1974).
- [39] A. D. Gadjiev, *On P. P. Korovkin type theorems*, Mat. Zametki, **20**, 781-786, (1976); Transl. in Math. Notes (5 – 6), 995-998, (1978).
- [40] A. D. Gadjiev and C. Orhan, *Some approximation theorems via statistical convergence*, Rocky Mountain J. Math., **32** (1), 129-138, (2007).